

## Erratum to: Association of cardiovascular disease risk factors with left ventricular mass, biventricular function, and the presence of silent myocardial infarction on cardiac MRI in an asymptomatic population

Eliel Nham<sup>1</sup> · Sung Mok Kim<sup>2,3</sup> · Sang-Chol Lee<sup>3,4</sup> · Sung-A Chang<sup>3,4</sup> · Jidong Sung<sup>4,5</sup> · Soo Jin Cho<sup>5</sup> · Shin Yi Jang<sup>3</sup> · Yeon Hyeon Choe<sup>2,3</sup>

Published online: 4 July 2016  
© Springer Science+Business Media Dordrecht 2016

**Erratum to: Int J Cardiovasc Imaging**  
**DOI 10.1007/s10554-016-0885-1**

Unfortunately in the original publication of this article, Fig. 1 has been incorrectly published. Correct Fig. 1 appears in this erratum.

---

The online version of the original article can be found under  
doi:[10.1007/s10554-016-0885-1](https://doi.org/10.1007/s10554-016-0885-1).

---

✉ Sang-Chol Lee  
sc.lea@samsung.com

✉ Yeon Hyeon Choe  
yhchoe@skku.edu


<sup>1</sup> Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea

<sup>2</sup> Department of Radiology and Center for Imaging Science, Samsung Medical Center, Sungkyunkwan University School of Medicine, 81 Ilwon-ro, Gangnam-gu, Seoul 06351, South Korea

<sup>3</sup> HVSI Imaging Center, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea

<sup>4</sup> Division of Cardiology, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea

<sup>5</sup> Center for Health Promotion, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea



**Fig. 1** Flow chart of the study population. *CMR* cardiac MRI, *MS* metabolic syndrome, *SMI* silent myocardial infarction, + presence, – absence. Silent myocardial infarction was present in 7 patients with diabetes mellitus (9.1% of 77 diabetic males and 6.9% of all diabetics)