

Erratum to: Effects of climate variability and climate change on the fishing conditions for grey mullet (*Mugil cephalus* L.) in the Taiwan Strait

**Kuo-Wei Lan · Ming-An Lee · Chang Ik Zhang ·
Pei-Yuan Wang · Long-Jing Wu · Kuo-Tien Lee**

Published online: 20 November 2014
© Springer Science+Business Media Dordrecht 2014

**Erratum to: Climatic Change (2014) 126:189–202
DOI 10.1007/s10584-014-1208-y**

The captions of Figs. 3 and 4 are in the correct position, but the actual figures should be interchanged.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10584-014-1208-y>.

K.-W. Lan · M.-A. Lee (✉) · P.-Y. Wang · K.-T. Lee
Department of Environmental Biology Fisheries Science, National Taiwan Ocean University,
2 Pei-Ning Rd., Keelung 20224, Taiwan, Republic of China
e-mail: malee@mail.ntou.edu.tw

K.-W. Lan · M.-A. Lee
Center of Excellence for Oceans, National Taiwan Ocean University, 2 Pei-Ning Rd, Keelung 20224,
Taiwan, Republic of China

C. I. Zhang
Department of Marine Production System Management, Pukyong National University, 559-1, Daeyeon,
Busan 608-737, South Korea

L.-J. Wu
Fisheries Research Institute, COA, Taiwan, Coastal and Offshore Resources Research Center,
1-1 North 1 st Road, Kaohsiung 80672, Taiwan, Republic of China

M.-A. Lee
Taiwan Group On Earth Observations, 13F-3, No.32, Gaotie 2nd Road, Hsinchu 30274, Taiwan,
Republic of China




Fig. 3 Cross-wavelet coherence between climatic indices and log10 (CPUE of grey mullet caught using purse seiners) (a–g), and climatic indices and winter SSTs (h–i): (a) annual PDO, (b) winter PDO, (c) autumn ONI, (d) annual sunspots, (e) winter Nino 3.4, (f) winter WPO, and (g) winter SST; (h) annual PDO (i) annual ONI with winter SST. The solid-black contour encloses regions of >95 % confidence, and the black line indicates the cone of influence where edge effects become important. The phase relationship is shown by arrows, with in-phase relationships pointing to the right, out-of-phase relationships pointing to the left




Fig. 4 Annual trends in (a) the CPUE of grey mullet caught using purse seiners (red line) and gill nets (blue line) with the PDO (black line) and (b) winter sea surface temperature anomaly (black dotted line) from 1967 to 2009