

## Erratum to: The role of theory building in the teaching of secondary geometry

Michael Weiss<sup>1</sup> · Patricio Herbst<sup>2</sup>

Published online: 1 April 2015  
© Springer Science+Business Media Dordrecht 2015

**Erratum to: Educ Stud Math (2015)**  
**DOI 10.1007/s10649-015-9599-x**

The original version of this article contained an error in the presentation of Theorem 2 (Ceva's Theorem). The corrected presentation is given below.

**Theorem 2 (Ceva's Theorem)** Consider triangle  $ABC$  with points  $P$  on  $\overleftrightarrow{AB}$ ,  $Q$  on  $\overleftrightarrow{BC}$ , and  $R$  on  $\overleftrightarrow{CA}$ . Then lines  $\overleftrightarrow{AQ}$ ,  $\overleftrightarrow{BR}$ , and  $\overleftrightarrow{CP}$  are concurrent if and only if

$$\frac{\langle AP \rangle}{\langle PB \rangle} \frac{\langle BQ \rangle}{\langle QC \rangle} \frac{\langle CR \rangle}{\langle RA \rangle} = 1$$

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10649-015-9599-x>.

✉ Patricio Herbst  
pgherbst@umich.edu  
Michael Weiss  
mweiss@msu.edu

<sup>1</sup> College of Education, Michigan State University, East Lansing, MI, USA

<sup>2</sup> School of Education, University of Michigan, Ann Arbor, MI, USA