




Correction to: Brief Report: Maternal Opioid Prescription from Preconception Through Pregnancy and the Odds of Autism Spectrum Disorder and Autism Features in Children

Eric Rubenstein¹  · Jessica C. Young² · Lisa A. Croen³ · Carolyn DiGuseppi⁴ · Nicole F. Dowling⁵ · Li-Ching Lee⁶ · Laura Schieve⁵ · Lisa D. Wiggins⁵ · Julie Daniels²

Published online: 20 September 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Correction to:
Journal of Autism and Developmental Disorders
<https://doi.org/10.1007/s10803-018-3721-8>

The original version of this article unfortunately contained a mistake in Table 2. The “Time of use” should be the column header with the corresponding row with “N = 126” belonging to the “Peri-pregnancy” row. The “Preconception” row should be the row with “N = 17”, “Trimester 1” should be the “N = 29” row, “Trimester 2” should be the “N = 25” row, and “Trimester 3” should be the “N = 47” row.

The corrected Table 2 is given below.

The original article can be found online at <https://doi.org/10.1007/s10803-018-3721-8>.

✉ Eric Rubenstein
Erubenstein2@wisc.edu

¹ Waisman Center, University of Wisconsin-Madison, Rm 529, 1500 Highland Avenue, Madison, WI 53705, USA

² Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina Chapel Hill, 135 Dauer Dr, Chapel Hill, NC 27599, USA

³ Kaiser Permanente Division of Research, 2000 Broadway, Oakland, CA 94612, USA

⁴ Department of Epidemiology, Colorado School of Public Health, University of Colorado Anschutz Medical Campus School of Public Health, 13001 E. 17th Place, Aurora, CO 80045, USA

⁵ National Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention, 1600 Clifton Road, Atlanta, GA 30329, USA

⁶ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe Street, Baltimore, MD 21205, USA

Table 2 Unadjusted and adjusted odds ratios and 95% confidence intervals for the association between maternal opioid prescriptions in the peri-pregnancy period and child neurodevelopmental outcomes in children born from 2003 to 2012 enrolled in the Study to Explore Early Development

Time of use	Unadjusted								
	ASD N = 1369			DD-only N = 938			ASD/DD with autism features ^a N = 1845		
	N	OR ^b	95% CI	N	OR	95% CI	N	OR	95% CI
Peri-pregnancy ^c	126	1.56	1.19, 2.06	66	1.17	0.84, 1.62	174	1.61	1.24, 2.08
Preconception ^d	17	2.82	1.27, 6.82	8	1.92	0.70, 5.34	23	2.83	1.21, 6.62
Trimester 1	29	1.99	1.09, 3.63	12	1.19	0.57, 2.50	36	1.83	1.02, 3.26
Trimester 2	25	1.52	0.84, 2.78	21	1.88	1.00, 3.52	38	1.72	0.99, 3.00
Trimester 3	47	1.16	0.77, 1.75	24	0.85	0.52, 1.41	63	1.15	0.78, 1.69
Time of use	Adjusted ^e								
	ASD N = 1359			DD-Only N = 927			ASD/DD with autism features N = 1834		
	N	aOR	95% CI	N	aOR	95% CI	N	aOR	95% CI
Peri-pregnancy	126	1.26	0.94, 1.68	66	1.06	0.76, 1.49	174	1.30	0.99, 1.71
Preconception	17	2.43	0.99, 6.02	8	1.94	0.68, 5.52	23	2.64	1.10, 6.31
Trimester 1	29	1.58	0.85, 2.94	12	1.06	0.50, 2.28	36	1.48	0.81, 2.69
Trimester 2	25	1.13	0.61, 2.12	21	1.64	0.87, 3.12	38	1.24	0.69, 2.22
Trimester 3	47	1.02	0.66, 1.57	24	0.85	0.52, 1.41	63	0.96	0.64, 1.45

ASD Autism spectrum disorder, *DD-only* developmental delay/disorder without features of ASD, *OR* odds ratio, *aOR* adjusted odds ratio, *CI* confidence interval

^aASD/DD with autism features group includes children with autism spectrum disorder or non-ASD developmental delay/disorders with autism features

^bExposure for case groups were compared to exposure for population controls at each time of use

^c3 months preconception to 1 day prior to childbirth

^d3 months preconception to conception

^eAdjusted for maternal education, race/ethnicity, smoking during pregnancy, psychiatric condition prior to childbirth, and SEED study period (2003–2006 or 2007–2012). Ns are smaller due to missing covariate data