

## **Erratum to: Factors Controlling Pre-Columbian and Early Historic Maize Productivity in the American Southwest, Part 2: The Chaco Halo, Mesa Verde, Pajarito Plateau/Bandelier, and Zuni Archaeological Regions**

**Larry V. Benson**

Published online: 19 June 2010  
© Springer Science+Business Media, LLC 2010

### **Erratum to: J Archaeol Method Theory DOI 10.1007/s10816-010-9083-y**

The original version of this article unfortunately contained a mistake.

p. nnn: Columns labeled ‘bu/ac’ should read ‘g NO<sub>3</sub>’ in Table 10.

In the explanation written below Table 10, ‘bu/ac indicates the number of bushels of maize that could be supported in the first year of plant growth’ should read ‘g NO<sub>3</sub> indicates the amount of nitrate produced within the root zone during the first year of plant growth.’

p. mmm: Columns labeled ‘bu/ac’ should read ‘g NO<sub>3</sub>’ in Table 11.

The first sentence of the explanation written below Table 11 should read ‘6.6 g of NO<sub>3</sub>-N is needed to produce 7 bu/ac of maize planted at a hill spacing of 2.4 m and an average hill density of 2 plants.’

In the explanation written below Table 11, ‘bu/ac indicates the number of bushels of maize that could be supported in the first year of plant growth’ should read ‘g NO<sub>3</sub> indicates the amount of nitrate produced within the root zone during the first year of plant growth.’

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10816-010-9083-y>.

L. V. Benson (✉)

National Research Program, U.S. Geological Survey, 3215 Marine St., Boulder, CO 80303, USA  
e-mail: lbenson@usgs.gov