

Erratum to: Olfactory Cues From Pathogenic Fungus Affect the Direct of Motion of Termites, *Coptotermes formosanus*

Aya Yanagawa¹ · Tomoya Imai¹ · Toshiharu Akino² · Yoshihiro Toh³ · Tsuyoshi Yoshimura¹

Published online: 13 September 2016
© Springer Science+Business Media New York 2016

Erratum to: J Chem Ecol (2015) 41:1118–1126
DOI 10.1007/s10886-015-0649-8

On page 1120, the unit associated with the external standard was mistyped. The solution of external standard was ‘10 ng/μl’, not ‘ng/ml’.

On page 1120 we should have reported that ‘The GC injections were 1 μl of sample or standard for these analyses’.

On page 1120 the standards used for Y tube tests are not adequately described. We should correct the current text as follows: ‘Dilutions were made by carrying out a serial dilution of 1 μl /ml solutions for each compound of interest (for 3-octanone, 1 μl /ml = 8.2 mg/ml; for 1-octene-3-ol, 1 μl /ml = 8.4 mg/ml). The solutions were diluted to concentrations as low as 8×10^{-18} ng/ml, which was the lower limit of detection of 3-octanone with our GC-MS method.’

On page 1122 the units were again mistyped. The correct concentration of 1-octen-3-ol was 0.052 ± 0.049 ng/μl in 10^7 conidia, and 3-octanone was 0.044 ng/μl in 10^7 conidia.

The online version of the original article can be found at <http://10.1007/s10886-015-0649-8>.

✉ Aya Yanagawa
ayanagawa@rsh.kyoto-u.ac.jp

¹ Research Institute for Sustainable Humanosphere, Kyoto University, Uji 611-0011, Japan

² Laboratory of Applied Entomology, Kyoto Institute of Technology, Ukyo-ku, Kyoto 616-8354, Japan

³ Department of Biology, Faculty of Science, Kyushu University, Higashi-ku, Fukuoka 812-8581, Japan