

Erratum to: Hypofractionated radiosurgery has a better safety profile than single fraction radiosurgery for large resected brain metastases

Bree R. Eaton^{1,4} · Michael J. LaRiviere¹ · Sungjin Kim^{2,5} · Roshan S. Prabhu^{1,6} · Kirtesh Patel¹ · Shravan Kandula¹ · Nelson Oyesiku³ · Jeffrey Olson³ · Walter Curran¹ · Hui-Kuo Shu¹ · Ian Crocker¹

Published online: 23 April 2015
© Springer Science+Business Media New York 2015

Erratum to: J Neurooncol
DOI 10.1007/s11060-015-1767-4

The second author's name was incorrect in the original publication. It is correct in this erratum.

The online version of the original article can be found under
doi:[10.1007/s11060-015-1767-4](https://doi.org/10.1007/s11060-015-1767-4).

✉ Bree R. Eaton
brupper@emory.edu

¹ Department of Radiation Oncology, Winship Cancer Institute of Emory University, Atlanta, GA, USA

² Department of Biostatistics and Bioinformatics, Winship Cancer Institute of Emory University, Atlanta, GA, USA

³ Department of Neurosurgery, Winship Cancer Institute of Emory University, Atlanta, GA, USA

⁴ Massachusetts General Hospital, Harvard Medical School, 55 Fruit Street, LL2, Boston, MA 02114, USA

⁵ Biostatistics and Bioinformatics Research Center, Cedars-Sinai Medical Center, Los Angeles, CA, USA

⁶ Southeastern Radiation Oncology Group, Levine Cancer Institute, Charlotte, NC, USA