

Erratum to: Rostrocaudal Dynamics of CSF Biomarkers

Andrew Tarnaris · Ahmed K. Toma ·
Miles D. Chapman · Axel Petzold · Geoff Keir ·
Neil D. Kitchen · Laurence D. Watkins

Published online: 2 October 2011
© Springer Science+Business Media, LLC 2011

Erratum to: Neurochem Res (2011) 36:528–532
DOI 10.1007/s11064-010-0374-1

The original publication of the article contains few lines of incorrect text. The corrected version of the paragraph and the references cited are given below for your reading.

There is a rostro-caudal tissue gradient of the neurofilament (Nf) protein concentration in the brain, with the lowest Nf protein concentration found in cortical neurons and the highest Nf protein concentration found in spinal cord axons, particularly within the large fibre motor axons of the pyramidal tracts [33–35]. Not surprisingly, the anatomical distribution of neurofilament heavy chain (NfH) is also reflected in the CSF concentration. Extending on Reiber's theory, we hypothesise that the RCG should be <1 , also for other protein biomarkers specific for the axon.

References

- Petzold A, Gveric D, Groves M, Schmierer K, Grant D, Chapman M, Keir G, Cuzner L, Thompson EJ (2008) Phosphorylation and compactness of neurofilaments in multiple sclerosis: indicators of axonal pathology. *Exp Neurol* 213:326–335
- Anderson KJ, Scheff SW, Miller KM, Roberts KN, Gilmer LK, Yang C, Shaw G (2008) The phosphorylated axonal form of the neurofilament subunit NF-H (pNF-H) as a blood biomarker of traumatic brain injury. *J Neurotrauma* 25:1079–1085
- Marszalek JR, Williamson TL, Lee MK, Xu Z, Hoffman PN, Becher MW, Crawford TO, Cleveland DW (1996) Neurofilament subunit nf-h modulates axonal diameter by selectively slowing neurofilament transport. *J Cell Biol* 135(3):711–724

The online version of the original article can be found under doi:[10.1007/s11064-010-0374-1](https://doi.org/10.1007/s11064-010-0374-1).

A. Tarnaris (✉) · A. K. Toma · N. D. Kitchen · L. D. Watkins
The Victor Horsley Department of Neurosurgery,
National Hospital for Neurology and Neurosurgery,
Box 32, Queen Square, London WC 1N 3BG, UK
e-mail: andrewtarnaris@gmail.com

M. D. Chapman · A. Petzold · G. Keir
Department of Neuroimmunology, Institute of Neurology,
Queen Square, London WC 1N 3BG, UK

A. Petzold
Department of Neurology, Free University Amsterdam,
Amsterdam, The Netherlands