

Identification of obligate C₃ photosynthesis in *Dendrobium*

S. QIU*, S. SULTANA*, Z.D. LIU*, L.Y. YIN**, and C.Y. WANG*,†

Key Laboratory for Biology of Horticultural Plants, Ministry of Education, Huazhong Agricultural University, Wuhan 430070, China*

Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresources, Hainan University, HaiKou 570228, China**

Abstract

Dendrobium is one of the three largest genera in the Orchidaceae and is distributed throughout various habitats. We investigated photosynthesis in seven *Dendrobium* species and cultivars by comparing their leaf $\delta^{13}\text{C}$ values, titratable acidity, and CO₂ exchange in well-watered and drought-stressed conditions. In addition, the leaf thickness and mesophyll succulence index (S_m) were measured in well-watered conditions. Our results indicate that *Dendrobium loddigesii* is a typical obligate (or constitutive) CAM plant because the leaf $\delta^{13}\text{C}$ values were -14.47 and -14.66‰ in both conditions, respectively. Others showed the leaf thickness of $0.31\text{--}0.89$ mm and their $\delta^{13}\text{C}$ values ranged from -25.68 to -28.37‰ . These are not the CAM plants but they could not be classified as obligate C₃ or C₃/CAM intermediate plants. *Dendrobium crepidatum* and *Dendrobium fimbriatum* were further identified as the obligate C₃ plants because the net CO₂ uptake was positive during daytime and negative during nighttime in both conditions. In contrast, *Dendrobium chrysotoxum*, *Dendrobium nobile*, and *D. nobile* 'V1' and 'V4', showed no positive net CO₂ uptake and low ΔH^+ values during nighttime under well-watered conditions, indicating C₃ photosynthesis. However, they showed the positive net CO₂ uptake and large ΔH^+ values during nighttime after drought-stress (21 or 28 days without H₂O), indicating CAM photosynthesis. Therefore, these four species and cultivars were identified as C₃/CAM intermediate (inducible or facultative) plants. In brief, obligate CAM, C₃/CAM intermediate, and obligate C₃ plant types all exist in the section of *Dendrobium*. To the best of our knowledge, this is the first report of the obligate C₃ plants in *Dendrobium*, and these diverse photosynthetic pathways may explain their varied environmental adaptations.

Additional key words: C₃/CAM intermediate plant; CO₂ exchange; crassulacean acid metabolism; titratable acidity; $\delta^{13}\text{C}$ value.

Introduction

The diverse photosynthetic pathways observed in plants are usually considered to be evolutionary adaptations to environmental variation (Ehleringer and Monson 1993, Ackerly *et al.* 2000, Caruso *et al.* 2005). C₃, C₄, and crassulacean acid metabolism (CAM) are three types of photosynthesis observed in vascular plants. The C₃ pathway was the first photosynthetic pathway that was biochemically characterized; C₄ and CAM photosynthetic pathways are evolutionarily derived from C₃ photosynthesis (Ehleringer and Monson 1993, Vaasen *et al.* 2006, Silvera 2010). CAM is a plastic photosynthetic

adaptation to arid environments (Cushman 2001). In contrast to C₃ and C₄ metabolism, where CO₂ uptake occurs during daytime, CAM is characterized by CO₂ uptake at nighttime with depressed CO₂ uptake and stomatal closure during daytime. Several C₃ plants are able to switch between C₃ and CAM modes, which are known as C₃/CAM intermediate (inducible or facultative) plants. For example, *Mesembryanthemum crystallinum* (the ice plant) is an annual plant, which switches its photosynthetic mode in response to water and/or salt stress (Winter and Holtum 2007). In contrast, most C₃ plants use only C₃

Received 14 January 2014, accepted 3 October 2014.

†Corresponding author; phone: +86 27 87282010, fax: +86 27 87282010, e-mail: wangcy@mail.hzau.edu.cn

Abbreviations: CAM – crassulacean acid metabolism; Chl – chlorophyll; DM – dry mass; DS – drought-stressed; FM – fresh mass; ΔH^+ – titratable acidity accumulated during nighttime; LWC – leaf water content; S_m – mesophyll succulence index; WW – well-watered; $\delta^{13}\text{C}$ – carbone isotope ratio $^{13}\text{C}/^{12}\text{C}$.

Acknowledgements: The authors would like to thank Enago (www.enago.cn) and Dr. Jing Luo of Huazhong Agricultural University for the English language review. This work was funded by the National Natural Science Foundation of China (31170657) and the “Twelfth Five-Year” National Science and Technology Project in Rural Areas (2011BAD12B02).

© The Author(s). This article is published with open access at link.springer.com

photosynthesis for CO₂ uptake throughout their life cycle, e.g., rice and wheat, and are classified as obligate C₃ plants. Plants that predominantly employ CAM photosynthesis to absorb CO₂ (Ting 1985), such as *Opuntia basilaris* (Hanscom and Ting 1978), are known as obligate CAM (or constitutive CAM) plants. CAM plants are widely distributed within the plant kingdom, i.e., 343 genera in 34 families (Holtum *et al.* 2007), but only few families, such as the Clusiaceae and Orchidaceae, include obligate C₃, C₃/CAM intermediate, and obligate CAM plants within a single family (Vaasen *et al.* 2006, Silvera 2010). Photosynthetic pathways have consistently demonstrated progress from C₃ ancestors to CAM photosynthesis (Ehleringer and Monson 1993, Vaasen *et al.* 2006, Silvera 2010). However, the genetic changes required for this progression (and reversion) remain unclear (Silvera 2010). Thus, it is important to study the multiple independent evolutionary origins of CAM in genera that contains the obligate C₃, C₃/CAM intermediate, and the obligate CAM plants. However, only a few genera, such as *Clusia* in the Clusiaceae and *Cymbidium* in the Orchidaceae, are known to contain the obligate C₃, C₃/CAM intermediate, and obligate CAM plants in a single genus (Vaasen *et al.* 2006, Silvera *et al.* 2010).

Orchids are model plants for studying the evolution of CAM. Orchidaceae is the largest family of flowering plants and CAM photosynthesis is widespread in this family (Winter *et al.* 1983, Silvera *et al.* 2005), i.e., approx. 9,000 species (Lüttge 2004, Silvera *et al.* 2010). *Dendrobium* is one of the three largest orchid genera, with approx. 1,500 wild species and several more hybrids (Cribb and Govaerts 2005, Lavarack *et al.* 2000, Schuiteman 2011). Most have high ornamental and/or medicinal value, such as *D. nobile*. Section *Dendrobium* is one of 11 *Dendrobium* sections, comprising 54 species that are mostly distributed in the Himalayas from India through to China as well as in Japan, with 27 species in China (Zhu *et al.* 2009). In the study of Gehrig *et al.* (2001), all *Dendrobium* species, which were investigated, were CAM plants. In *Dendrobium*, 25 wild species and some cultivars (mainly *Phalaenopsis* cane-type cultivars) are known to utilize exclusively the CAM photosynthetic pathway (i.e., obligate CAM plants) (Hew and Khoo 1980, Ando 1982, Fu and Hew 1982, Winter *et al.* 1983, He *et al.* 1998, Sayed 2001). However, several species in this genus appear to be very flexible with respect to their photosynthetic pathways; they switch between C₃ and CAM photosynthesis (Winter *et al.* 1983, Su *et al.* 2003, Ren 2008). However, no cases of obligate C₃ have been reported in *Dendrobium* in previous studies.

Several methods have been developed to identify the photosynthetic pathways in plants, such as leaf succulence or thickness and the leaf $\delta^{13}\text{C}$ value. In most species, there

is a close correlation between greater leaf succulence or thickness and increasing magnitude of CAM (Silvera 2010), because CAM is typically associated with epiphytic or succulent life forms in tropical species. However, it is difficult to identify whether a plant is obligate C₃ based only on the leaf thickness. For example, a few plants with thinner leaves fix CO₂ using CAM, e.g., the leaf thickness of *D. bigibbum* is only 0.79 mm, but it yields -11.9‰ of $\delta^{13}\text{C}$, which is another characteristics of CAM plants (Winter *et al.* 1983).

C₃ and CAM plants can be distinguished by their leaf $\delta^{13}\text{C}$ value measurements ranging from -20 to -35‰ in C₃ and from -10 to -22‰ in CAM (O'Leary 1981, Sternberg *et al.* 1984, Farquhar *et al.* 1989). Twelve *Dendrobium* species (including *D. nobile*) were characterized as C₃ plants based on their leaf $\delta^{13}\text{C}$ values, which ranged from -20 to -35‰ (Winter *et al.* 1983). However, this analysis showed that *D. nobile* was the C₃ plant with the $\delta^{13}\text{C}$ value of 25.1‰, although the CO₂ exchange pattern could be induced to switch from the C₃ to the CAM mode by drought stress (Winter *et al.* 1983, Su *et al.* 2003, Ren 2008). Thus, *D. nobile* was shown to be the C₃/CAM intermediate plant. The $\delta^{13}\text{C}$ measurements are less time consuming and more convenient for large species surveys, but the $\delta^{13}\text{C}$ value alone cannot distinguish obligate C₃ and C₃/CAM intermediate species if CAM gives a small contribution to the total carbon gain of the C₃/CAM intermediate pathway (Borland *et al.* 1993, Holtum and Winter 1999, Winter and Holtum 2002).

Careful measurements of diel changes of titratable acidity or dark CO₂ uptake have been recently used to distinguish obligate C₃ and C₃/CAM intermediate species, although these two methods are highly time-consuming (Borland *et al.* 1993, Holtum and Winter 1999, Winter and Holtum 2002). In C₃/CAM intermediate plants, environmental stress (such as drought) can induce or enhance the expression of CAM; therefore, comparing diel changes of titratable acidity or dark CO₂ uptake under optimal conditions and environmental stress are useful for characterizing obligate C₃ and C₃/CAM intermediate plants.

In this study, based on leaf thickness differences, we selected five species from the section of *Dendrobium* (including *D. nobile*), one species from the section of *Chrysotoxae*, and two cultivars of *D. nobile* and identified their photosynthesis patterns by comparing the leaf thickness, mesophyll succulence index (S_m), $\delta^{13}\text{C}$ values, titratable acidity, and CO₂ exchange pattern. We observed a new type in *Dendrobium*, which was characterized by obligate C₃ photosynthesis. The discovery that three types of photosynthesis occur in plants within the single section of *Dendrobium* could help to understand the diversity of photosynthetic pathways and the evolutionary and ecological importance of CAM photosynthesis.

Materials and methods

Plant material and treatments: Seven *Dendrobium* species (cultivars) were selected as accessions in this study, *i.e.*, four species from the section of *Dendrobium* (including *D. nobile*), one species from the section of *Chrysotoxae* (*D. chrysotoxum*), and two *D. nobile* cultivars (Table 1). *D. nobile* and *D. chrysotoxum* were selected for the comparison because they are known to be the C₃/CAM intermediate plants. These species and cultivars were grown and maintained in the greenhouse at Huazhong Agricultural University, Wuhan, China. The potting medium used was moss and coconut blocks at the ratio of 3:1 (v/v); the diameter of the coconut block was 1.2 cm. All plants were watered with 1/2 HS nutrient solution (Hoagland and Snyder 1933) once a week from March to October during the growing season. From September to November, well-developed plants were selected when they developed more than four pairs of leaves; they were assigned to well-watered (WW) and drought-stressed (DS) treatments. The WW plants were placed into a climate chamber with light intensity at the level of leaves kept at about 250–350 $\mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD, a 55/75% (day/night) of relative air humidity, a 12/12 h (light/dark) photoperiod, and a temperature regime of 28/18°C (day/night) for at least two weeks. The plants were watered every other day. The volume of the climate chamber was 400 L. These conditions satisfied the requirements of the C₃/CAM intermediate plants to switch to the C₃ photosynthesis mode. The DS plants were placed into other climate chambers with light at about 250–350 $\mu\text{mol}(\text{photon}) \text{m}^{-2} \text{s}^{-1}$, 55/75% (day/night) humidity, and a 12/12 h (light/dark) photoperiod. The plants were kept without watering for at least 21 d to induce the CAM photosynthesis mode in the C₃/CAM intermediate plants. Normally, the potting medium contains water and is very difficult to dry, thus in order to dry plants easier, a temperature regime of 30/22°C (day/night) was applied. The middle leaves of the pseudobulbs were selected for measurements when they were fully expanded. All experiments were performed in triplicate.

Leaf succulence: The leaf succulence of the WW plants was determined by measuring the leaf thickness and the mesophyll succulence index (S_m). Six to ten leaves were measured with precision calipers to determine the leaf thickness. The leaf water content (LWC) was determined as the difference between the fresh mass (FM) and the mass after drying for 72 h at 75°C [dry mass (DM)], *i.e.*, $\text{LWC} = (\text{FM} - \text{DM})/\text{FM}$. The total Chl content was spectrophotometrically determined at wavelengths of 663 nm (Chl *a*) and 645 nm (Chl *b*) (TU-1810, Beijing Purkinje General Instrument Co. Ltd., Beijing, China) (Hao *et al.* 2013) using 0.2-g leaf sections, which were collected at 18:00 h and weighed before cutting into smaller pieces. Chl was extracted from these samples for 12 h using 20 ml

of acetone and ethyl alcohol (1:1, v/v). The formula used to calculate the Chl content was: $\text{Chl} = (8.05 \text{ OD}_{663} + 20.29 \text{ OD}_{645}) 10^{-1}$, where the results were expressed in $\text{mg g}^{-1}(\text{FM})$. S_m was calculated using the following formula: $S_m = \text{g}(\text{H}_2\text{O}) \text{mg}^{-1}(\text{Chl})$, *i.e.*, $S_m = \text{LWC}/\text{Chl}$ content, which is the improved version of the method proposed by Kluge and Ting (1978).

Leaf carbon isotope ratio ($\delta^{13}\text{C}$): Leaf samples were harvested from the WW and DS-treated plants, frozen in liquid nitrogen, and stored at -70°C before measuring. Furthermore, the carbon isotope ratios ($^{13}\text{C}/^{12}\text{C}$) were determined using freezing-dried samples with their mass approx. 3.0 mg (Winter and Holtum 2002). Samples were analyzed using an organic elemental analyzer (*Delta V Advantage Isotope Ratio MS*, Thermo Fisher Scientific Inc., Bremen, Germany) and compared with the PDB standard (*Belemnite americana*). The stable carbon isotope composition was expressed in conventional delta (δ) notation as the $^{13}\text{C}/^{12}\text{C}$ ratio relative to the standard: $\delta^{13}\text{C} [\text{‰}] = [(\delta^{13}\text{C} \text{ sample})/(\delta^{13}\text{C} \text{ standard}) - 1] \times 1,000$. The precision of the mass spectrometer used for the $\delta^{13}\text{C}$ analyses was 0.05‰.

Leaf titratable acidity: Leaf samples were harvested from the WW and DS plants at the end (18:00 h) and beginning (6:00 h) of the photoperiod, where 1.0 g of leaf samples were weighed, frozen in liquid nitrogen, and stored at -70°C before measuring their titratable acidity (Eastmond and Ross 1997). Samples were ground in a prechilled pestle and mortar with 5 ml of distilled water, and the crude extract was boiled for 20 min and then allowed to cool to room temperature. The extract was centrifuged at $5,000 \times g$ for 15 min and 2 ml of the supernatant was diluted to 25 ml with distilled water before titration with 0.01 M KOH to the end point of pH 7.2, using bromothymol blue as an indicator. The titratable acidity was expressed in $\mu\text{mol}(\text{H}^+) \text{g}^{-1}(\text{FM})$. The titratable acidity accumulated during nighttime (ΔH^+) was calculated as: $\Delta\text{H}^+ = \text{titratable acidity at dawn} - \text{titratable acidity at dusk}$.

Leaf CO₂ exchange: The net CO₂ uptake was measured every 2 h for 24 h using an infrared gas-exchange system equipped with a LED light source (LI-6400-02B) (LI-6400, LI-COR Inc., Lincoln, NE, USA) (Pierce *et al.* 2002). Each leaf portion was enclosed in a chamber (2 cm \times 3 cm), where the environmental conditions were controlled. A CO₂ cylinder was used to maintain a constant CO₂ concentration of 380 $\mu\text{mol mol}^{-1}$. The light intensity at the chamber was 300 $\mu\text{mol}(\text{photon}) \text{m}^{-2} \text{s}^{-1}$ supplied by the LED, the photoperiod of 12/12 h (light/dark), and humidity of 55/75% (day/night) for both treatments. The temperature regime was 28/18°C (day/night) for the WW plants and 30/22°C for the DS plants, respectively. On the

same time, the chamber of LI-6400 was also placed during measurements in the bigger climate chamber (400 L), where the environmental conditions were also controlled as above. The net CO₂ uptake was analyzed for each treatment; the fully expanded leaves from the middle parts of the pseudobulbs were measured from six randomly selected plants.

Statistical analyses: A completely randomized design was used in the experiments and repeated three times. The means were compared by analysis of variance (ANOVA,

Results

Leaf thickness and S_m: *D. loddigesii* showed the highest leaf thickness of 0.99 mm, which was significantly different from the other species or cultivars (Table 1). The leaf thickness of *D. fimbriatum* was the lowest, only 0.31 mm, although the leaf thickness of *D. crepidatum* was also low (0.38 mm). The leaf thickness of the other four species or cultivars (*i.e.*, *D. chrysotoxum*, *D. nobile*, and *D. nobile* ‘V1’ and ‘V4’) ranged from 0.58 mm to 0.89 mm. The highest S_m value was obtained with *D. loddigesii*, *i.e.*, 2.55 g(H₂O) mg⁻¹(Chl), which was significantly higher than those of the others. The lowest S_m value was obtained with *D. crepidatum*, *i.e.*, 0.53 g(H₂O) mg⁻¹(Chl). The S_m values of the other five species and cultivars ranged from 0.57 to 1.22 g(H₂O) mg⁻¹(Chl).

Leaf δ¹³C values: The δ¹³C values of *D. loddigesii* were within -15‰, *i.e.*, -14.47‰ and -14.66‰ under the WW and DS conditions, respectively (Table 2). This is a typical value for the obligate CAM plants. In the other species or cultivars, the leaf δ¹³C values ranged from -25.68 to -28.37‰, which were lower than those of *D. loddigesii*. These values are typical for the C₃ plants, but we could not distinguish the obligate C₃ plants and the C₃/CAM intermediate plants based on this value alone. In addition, except *D. nobile*, no significant differences were in the δ¹³C values of the WW and DS plants in other six species or cultivars.

Titrate acidity: For *D. loddigesii*, there was a

three means) using SAS v. 8.0 (SAS Institute Inc., USA). Significant differences in leaf thickness and the mesophyll succulence index (S_m) among different species were analyzed by ANOVA with Duncan’s multiple range test; the *t*-test was used to test for differences in leaf δ¹³C values between WW and DS plants and to test for differences between dawn and dusk of WW plants and that of DS plants within each species. The *t*-test was used to test for differences in ΔH⁺ between WW and DS plants. The *P*<0.05 was considered as statistically significant.

significant difference between dawn and dusk in WW and DS conditions (Table 3). The highest ΔH⁺ was obtained in *D. loddigesii*, *i.e.*, 355.5 and 425.9 μmol(H⁺) g⁻¹(FM) in WW and DS conditions, respectively. For the other six species and cultivars, the ΔH⁺ showed no significant difference between dawn and dusk in WW condition, which indicated that plants use C₃ photosynthetic pathway to uptake CO₂ under such condition. After DS for 21 d, the ΔH⁺ was not significantly different between dawn and dusk in *D. crepidatum* and *D. fimbriatum*, but the ΔH⁺ value of *D. fimbriatum* significantly increased to 48.0 μmol(H⁺) g⁻¹(FM) compared with that of WW conditions. In contrast, for the other four species and cultivars, including *D. chrysotoxum*, *D. nobile*, and *D. nobile* ‘V1’ and ‘V4’, the ΔH⁺ was significantly different between dawn and dusk in DS condition. In addition, the ΔH⁺ values were 116.2, 184.8, 166.5, and 192.8 μmol(H⁺) g⁻¹(FM), respectively, which clearly showed the significant increase compared with those of WW condition.

Leaf CO₂ exchange in plants with CAM photosynthesis: In the WW plants, net CO₂ uptake occurred only from 6:00 to 8:00 h, and the values were negative during most of the daytime (10:00 – 18:00 h) and nighttime (Fig. 1). After 14-d drought stress, there were significantly negative values during daytime and positive during nighttime. However, if the drought stress continued for another 14 d, the net CO₂ uptake was more or less constant at a low value (approx. 0 μmol m⁻² s⁻¹) throughout the day,

Table 1. The leaf thickness and mesophyll succulence index (S_m) of seven *Dendrobium* species and cultivars in well-watered conditions. Results followed by the same letter within each species were not significantly different (*p*<0.05). The results represent the mean ± SE (*n* = 3).

Species or cultivar	Section	Thickness [mm]	S _m [g(H ₂ O) mg ⁻¹ (Chl)]
<i>D. crepidatum</i> Lindl.	<i>Dendrobium</i>	0.38 ± 0.02 ^f	0.53 ± 0.05 ^d
<i>D. fimbriatum</i> Hook.	<i>Dendrobium</i>	0.31 ± 0.01 ^f	0.58 ± 0.03 ^d
<i>D. nobile</i> Lindl.	<i>Dendrobium</i>	0.63 ± 0.03 ^{de}	1.22 ± 0.08 ^b
<i>D. nobile</i> ‘V1’	cultivar	0.76 ± 0.02 ^c	1.10 ± 0.07 ^{bc}
<i>D. nobile</i> ‘V4’	cultivar	0.58 ± 0.01 ^e	0.90 ± 0.09 ^c
<i>D. chrysotoxum</i> Lindl.	<i>Chrysotoxae</i>	0.89 ± 0.04 ^b	1.05 ± 0.02 ^{bc}
<i>D. loddigesii</i> Rolf.	<i>Dendrobium</i>	0.99 ± 0.04 ^a	2.55 ± 0.15 ^a

Table 2. Leaf $\delta^{13}\text{C}$ ($^{13}\text{C}/^{12}\text{C}$) values for seven species of *Dendrobium*. Results followed by the same letter within each species were not significantly different ($p < 0.05$). The results represent the mean \pm SE ($n = 3$).

Species or cultivar	Leaves/well-watered [%]	Leaves/drought-stressed [%]
<i>D. crepidatum</i>	-27.63 ± 0.27^a	-28.37 ± 0.55^a
<i>D. fimbriatum</i>	-27.62 ± 0.15^a	-27.49 ± 0.38^a
<i>D. nobile</i>	-27.07 ± 0.33^a	-25.68 ± 0.21^b
<i>D. nobile</i> 'V1'	-26.34 ± 0.73^a	-26.73 ± 0.23^a
<i>D. nobile</i> 'V4'	-28.09 ± 0.20^a	-26.26 ± 0.06^a
<i>D. chrysotoxum</i>	-27.55 ± 0.16^a	-26.08 ± 0.76^a
<i>D. loddigesii</i>	-14.47 ± 0.28^a	-14.66 ± 0.43^a

Table 3. Dawn and dusk analysis of the leaf titratable acidity in seven *Dendrobium* species and cultivars. Results followed by the same small letter of dawn and dusk under well-watered conditions within each species were not significantly different ($p < 0.05$); results followed by the same capital letter of dawn and dusk under drought-stressed treatment within each species were not significantly different ($p < 0.05$); results followed by the same number of stars of dawn-dusk between well-watered and drought-stressed treatment within each species were not significantly different ($p < 0.05$). The results represent the mean \pm SE ($n = 3$).

Species or cultivar	Well-watered [$\mu\text{mol}(\text{H}^+) \text{g}^{-1}(\text{FM})$]			Drought-stressed [$\mu\text{mol}(\text{H}^+) \text{g}^{-1}(\text{FM})$]		
	Dawn	Dusk	Dawn–Dusk	Dawn	Dusk	Dawn–Dusk
<i>D. crepidatum</i>	175.4 ± 19.2^a	192.1 ± 50.5^a	$-16.7 \pm 31.2^*$	241.7 ± 50.5^A	242.3 ± 46.8^A	$-0.6 \pm 3.7^*$
<i>D. fimbriatum</i>	194.3 ± 15.2^a	228.4 ± 16.3^a	$-34.1 \pm 1.1^{**}$	222.9 ± 14.5^A	174.9 ± 15.9^A	$48.0 \pm 11.4^*$
<i>D. nobile</i>	289.9 ± 57.6^a	291.3 ± 20.7^a	$-1.3 \pm 36.9^{**}$	359.1 ± 41.4	174.3 ± 10.7^B	$184.8 \pm 30.7^*$
<i>D. nobile</i> 'V1'	330.3 ± 43.2^a	278.5 ± 11.4^a	$51.8 \pm 32.2^{**}$	331.2 ± 16.5^A	164.7 ± 19.0^B	$166.5 \pm 2.5^*$
<i>D. nobile</i> 'V4'	241.1 ± 30.6^a	228.2 ± 21.5^a	$12.9 \pm 9.1^{**}$	425.7 ± 51.2^A	232.9 ± 37.5^B	$192.8 \pm 13.6^*$
<i>D. chrysotoxum</i>	311.5 ± 23.7^a	275.8 ± 15.3^a	$35.9 \pm 8.4^{**}$	312.1 ± 13.5^A	195.9 ± 27.9^B	$116.2 \pm 14.3^*$
<i>D. loddigesii</i>	651.3 ± 36.3^a	295.8 ± 36.0^b	$355.5 \pm 34.6^{**}$	705.5 ± 25.5^A	279.6 ± 43.0^B	$425.9 \pm 17.5^*$




Fig. 1. Net CO_2 exchange by *Dendrobium loddigesii* in well-watered conditions (T1), 14 days without water (T2), and 28 days without water (T3). The results represent the means where $n = 6$. Open bar represents light period, closed bar represents dark period.

excluding 0.03 (2:00 h) and 0.02 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (4:00 h) in the evening (Fig. 1). Thus, *D. loddigesii* utilized the CAM photosynthesis mode under both conditions. In addition, the stomatal conductance of *D. loddigesii* in WW conditions, after 14 d and 28 d without H_2O showed values more or less about 0 $\text{mmol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, except they were 0.045 and 0.085 $\text{mmol m}^{-2} \text{s}^{-1}$ at 18:00 and 20:00 h after 14 days without H_2O (Fig. 1S, supplement available online).

Leaf CO_2 exchange of plants with C_3 photosynthesis:

The net CO_2 uptake during daytime was positive in WW conditions (0.44 to 3.52 $\mu\text{mol m}^{-2} \text{s}^{-1}$), whereas no net CO_2 uptake occurred during nighttime (-0.5 to $-2.92 \mu\text{mol m}^{-2} \text{s}^{-1}$) (Fig. 2A). After 28 d without H_2O , the net CO_2 uptake during daytime decreased to a small positive value (approx. 0.5 $\mu\text{mol m}^{-2} \text{s}^{-1}$), but there was still no net CO_2 uptake during nighttime. The results were similar for




Fig. 2. Net CO₂ exchange by *Dendrobium crepidatum* (A) in well-watered conditions (T1) and 28 days without water (T2); and *Dendrobium fimbriatum* (B) in well-watered conditions (T1), 21 days without water (T2), and 28 days without water (T3). The results represent the means where $n = 6$. Open bars represent light periods, closed bars represent dark periods.




Fig. 3. Net CO₂ exchange by *Dendrobium nobile* (A) and *Dendrobium nobile* 'V1' (B) in well-watered conditions (T1) and 28 days without water (T2), respectively. The results represent the means where $n = 6$. Open bars represent light periods, closed bars represent dark periods.

D. fimbriatum (Fig. 2B). In addition, the leaf CO₂-exchange patterns of *D. fimbriatum* were measured after it was stressed for 35 d; the net CO₂ uptake decreased to a smaller positive value during daytime (from 0.032 to 0.489 μmol m⁻² s⁻¹) and negative CO₂ uptake during nighttime

(from -0.126 to -0.027 μmol m⁻² s⁻¹) (Fig. 2B). Thus, there was no net CO₂ uptake by the two species during nighttime even under severe drought stress. The g_s of the two species was about 0–0.03 mmol m⁻² s⁻¹ (Fig. 2S, supplement available online).




Fig. 4. Net CO₂ exchange by *Dendrobium nobile* 'V4' (A) and *Dendrobium chrysotoxum* (B) in well-watered conditions (T1), 21 days without water (T2), and 28 days without water (T3), respectively. The results represent the means where $n = 6$. Open bars represent light periods, closed bars represent dark periods.

Leaf CO₂ exchange in plants with C₃/CAM intermediate photosynthesis: For *D. nobile*, *D. nobile* 'V1', *D. nobile* 'V4', and *D. chrysotoxum*, the net CO₂ uptake during daytime in WW condition was more or less constant with positive rates (Figs. 3, 4), whereas almost no nocturnal net CO₂ uptake was observed (except there was some positive at 18:00 h in *D. nobile*). For *D. nobile* 'V4' (Fig. 4A) and *D. chrysotoxum* (Fig. 4B), the net CO₂ uptake began to show positive values during nighttime after DS for 21 d. Nevertheless, *D. nobile* and *D. nobile* 'V1' began to switch from C₃ photosynthesis to the CAM photosynthesis mode after 28 d without water, with approx.

0 μmol m⁻² s⁻¹ throughout the day and only a small net CO₂ uptake during nighttime (Fig. 3). The daily CO₂ exchange patterns of *D. nobile* 'V4' (Fig. 4A) and *D. chrysotoxum* (Fig. 4B) in DS conditions (28 d without H₂O) are also shown in Fig. 4. The net CO₂ uptake was more or less constant at a low value (approx. 0 μmol m⁻² s⁻¹) throughout the day, while the CO₂ exchange values were occasionally positive during nighttime and negative during daytime although they were very low; it was different from that of *D. fimbriatum* under drought stress for 35 d. In addition, the g_s of the four species or cultivars are shown in Figs. 3S, 4S (*supplements available online*).

Discussion

The results in this study demonstrated that obligate C₃ photosynthesis occurred in *Dendrobium*. For *D. crepidatum* and *D. fimbriatum*, net CO₂ uptake was positive during the day and negative during the night under the WW and DS conditions, which indicated they only use C₃ photosynthesis to fix CO₂. In addition, all of them yielded the δ¹³C values lower than -27‰ under WW and DS conditions. Their leaf thickness was only 0.38 and 0.31 mm, respectively; and their S_m were 0.53 and 0.58 g(H₂O) mg⁻¹(Chl), which were also far below 1 g(H₂O) mg⁻¹(Chl). Thus, results were also consistent with other C₃ plants with lower leaf succulence and lower δ¹³C values. Winter *et al.* (1983) and Silvera *et al.* (2005) investigated leaf thickness and δ¹³C values of about 400 plants and found 105 plants with leaf thickness lower than 0.40 mm. All 105 plants did not yield the δ¹³C values higher than -22‰, which indicated the plants with leaf thickness thinner than 0.40 mm that may only use C₃ photosynthesis to take up CO₂. However, the ΔH⁺ of *D. fimbriatum* increased signifi-

cantly under DS conditions, which might show that they use CAM photosynthesis to take up CO₂, but we did not find any significant change in ΔH⁺ (dawn-dusk) of malic acid and citrate between WW and DS plants measured by HPLC (Liu 2014). Mao and Zhang (1985) also found obligate C₃ plant, *Oxalis corniculata*, with ΔH⁺ of 120 μmol(H⁺) g⁻¹(FM). Here, we were not sure what caused the increment of ΔH⁺ values in *D. fimbriatum*. In 2001, Gehrig *et al.* discussed all *Dendrobium* species might be CAM plants for they have succulence organs. But we now know that not all succulent plants are CAM plants. And Ando (1982) identified some *D. nobile* hybrids as not being CAM plants. Thus, based on these analyses, we identified the two species only use C₃ photosynthesis no matter they were under stress condition or not.

Obligate CAM and C₃/CAM intermediate species were also found in the present study. *D. loddigesii* is the typical obligate (or constitutive) CAM plant because the leaf δ¹³C values were -14.47 and -14.66‰ in both conditions. In

addition, the net CO₂ exchange pattern was significantly negative during daytime and positive during nighttime after DS treatment for 14 d without water (Fig. 1). However, we observed the net CO₂ uptake of *D. loddigesii* was positive only between 6:00 and 10:00 h in WW conditions, which could be classified as “CAM cycling”. After 28-d drought stress, it maintained constant at a low value (approx. 0 μmol m⁻² s⁻¹) throughout the day, though there were 0.03 (2:00 h) and 0.02 μmol m⁻² s⁻¹ (4:00 h) in the evening (Fig. 1); it could be classified as “CAM idling”. The phenomenon of “CAM cycling” and “CAM idling” belongs to different permutations of CAM (Szarek *et al.* 1973, Osmond 1978, Ting 1985). It was reported that the obligate CAM plant, *Agave deserti*, could switch from a strong CAM type to a C₃ type pattern when plants were watered extensively (Hartsock and Nobel 1976). However, the day/night pattern of CO₂ exchange in the obligate CAM plant, *A. angustifolia*, did not shift towards the C₃ pattern completely when the supply of water was effectively unlimited (Winter *et al.* 2014). In contrast, *D. chrysotoxum*, *D. nobile*, and *D. nobile* ‘V1’ and ‘V4’ were identified as the C₃/CAM intermediate (inducible or facultative) plants, because there was no positive net CO₂ uptake during nighttime and low ΔH⁺ values in WW conditions, but there was some degree of positive net CO₂ uptake during nighttime and large ΔH⁺ values in DS conditions (21 or 28 d without water). However, the net CO₂ uptake of *D. nobile* ‘V4’ and *D. chrysotoxum* did not show any significant positive values during nighttime in DS condition (28 d without water); there was only 0.002–0.003 μmol m⁻² s⁻¹ (0:00–4:00 h) and 0.008–0.136 μmol m⁻² s⁻¹ (4:00–6:00 h), respectively. We also consider this the phenomenon of “CAM idling”. In addition, Ando (1982) identified some *D. nobile* hybrids as not being CAM plants in WW environment. In this study, we further investigated and found that photosynthesis pathway of *D. nobile* ‘V1’ and ‘V4’ could be induced from the C₃ mode to the CAM mode by stress; thus, they belong to the

C₃/CAM intermediate plants.

This comparative study of the single diverse section *Dendrobium* in one genus might be particularly useful for understanding the evolution and ecological importance of CAM. The molecular biology, biochemistry, and ecophysiology of CAM are well understood, but little is known about the evolutionary origins of CAM, particularly the genetic changes required for its evolution (Ehleringer and Monson 1993, Silvera *et al.* 2009). Thus, the section *Dendrobium*, where plants show three types of photosynthesis, suggests that the evolution towards CAM photosynthesis is still ongoing in the genus *Dendrobium*. This provides an opportunity to understand the adaptive and evolutionary significance of photosynthetic characteristics from an evolutionary view.

Many *Dendrobium* species are grown for their high ornamental or medicinal value. Understanding the photosynthetic pathways of plants is important for developing and improving cultivation techniques, and the results of the present study might be useful for local orchid growers who need to optimize the supply of water or light to plants. In addition, *Dendrobium* species have been the backbone of the cut-flower orchid business for several years. Since the first hybrid was reported in 1855, more than 8,000 novel *Dendrobium* cultivars have been bred *via* interspecific hybridization (Lavarack *et al.* 2000). However, several cultivars exhibit slow growth and/or they require high energy inputs. Most obligate C₃ plants grow more rapidly than obligate CAM plants, and C₃/CAM intermediate plants can use C₃ photosynthesis to improve their production in well-watered conditions, whereas they switch to CAM to survive in stressful environments. Thus, we can select obligate C₃ plants as parents to breed rapid-growing cultivars and use C₃/CAM intermediate or obligate CAM plants to breed new cultivars, more tolerant to adverse environments. Therefore, the results of this research might help select parents for cross-breeding of *Dendrobium* plants.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Ackerly D.D., Dudley S.A., Sultan S. E. *et al.*: The evolution of plant ecophysiological traits: recent advances and future directions. – *Bioscience* **50**: 979-995, 2000.
- Ando T.: Occurrence of two different modes of photosynthesis in *Dendrobium* cultivars. – *Sci. Hortic.-Amsterdam* **17**: 169-175, 1982.
- Borland A.M., Griffiths H., Broadmeadow M.S.J. *et al.*: Short-term changes in carbon isotope discrimination in the C₃-CAM intermediate *Clusia minor* L. growing in Trinidad. – *Oecologia* **95**: 444-453, 1993.
- Caruso C.M., Maherali H., Mikulyuk A. *et al.*: Genetic variance and covariance for physiological traits in lobelia: are there constraints on adaptive evolution? – *Evolution* **59**: 826-837, 2005.
- Cribb P., Govaerts R.: Just how many orchids are there? – In: Raynal-Roques A., Rogeueuant A., Prat D. (ed.): *Proceedings of the 18th World Orchid Conference*. Pp. 161-172. Naturalia Publications, Dijon 2005.
- Cushman J.C.: Crassulacean acid metabolism. A plastic photosynthetic adaptation to arid environments. – *Plant Physiol.* **127**: 1439-1448, 2001.
- Eastmond P.J., Ross J.D.: Evidence that the induction of crassulacean acid metabolism by water stress in *Mesembryanthemum crystallinum* (L.) involves root signaling. – *Plant Cell Environ.* **20**: 1559-1565, 1997.
- Ehleringer J.R., Monson R.K.: Evolutionary and ecological aspects of photosynthetic pathway variation. – *Annu. Rev.*

- Ecol. Syst. **24**: 411-439, 1993.
- Farquhar G.D., Ehleringer J.R., Hubick K.T.: Carbon isotope discrimination and photosynthesis. – *Annu. Rev. Plant Phys.* **40**: 503-537, 1989.
- Fu C.F., Hew C.S.: Crassulacean acid metabolism in orchids under water stress. – *Bot. Gaz.* **143**: 294-297, 1982.
- Gehrig H., Heute V., Kluge M.: New partial sequences of phosphoenolpyruvate carboxylase as molecular phylogenetic markers. – *Mol. Phylogenet. Evol.* **20**: 262-274, 2001.
- Hanscom Z., Ting I.P.: Irrigation magnifies CAM-photosynthesis in *Opuntia basilaris* (Cactaceae). – *Oecologia* **33**: 1-5, 1978.
- Hartsock T.L., Nobel P.S.: Watering converts a CAM plant to daytime CO₂ uptake. – *Nature* **262**: 574-576, 1976.
- Hao Y.H., Wu C.F., Zhao D.W. *et al.*: Proteomic analysis of glucohexaose induced resistance to downy mildew in *Cucumis sativus*. – *Aust. J. Crop Sci.* **7**: 1242-1251, 2013.
- He J., Khoo G.H., Hew C.S.: Susceptibility of CAM *Dendrobium* leaves and flowers to high light and high temperature under natural tropical conditions. – *Environ. Exp. Bot.* **40**: 255-264, 1998.
- Hew C.S., Khoo S.I.: Photosynthesis of young Orchid seedlings. – *New Phytol.* **86**: 349-357, 1980.
- Hoagland D.R., Snyder W.C.: Nutrition of strawberry plants under controlled conditions: (a) Effects of deficiencies of boron and certain other elements: (b) Susceptibility to injury from sodium salts. – *P. Am. Soc. Hortic. Sci.* **30**: 288-294, 1933.
- Holtum J.A.M., Winter K.: Degrees of crassulacean acid metabolism in tropical epiphytic and lithophytic ferns. – *Aust. J. Plant Physiol.* **26**: 749-757, 1999.
- Holtum J.A.M., Winter K., Weeks M.A. *et al.*: Crassulacean acid metabolism in the ZZ plant, *Zamioculcas zamiifolia* (Araceae). – *Am. J. Bot.* **94**: 1670-1676, 2007.
- Kluge M., Ting I.P.: Crassulacean acid metabolism. Analysis of an Ecological Adaptation. Springer-Verlag, Berlin – Heidelberg – New York 1978.
- Lavarack B., Harris W., Stocker G.: *Dendrobium* and its Relatives. Timber Press, Portland 2000.
- Liu Z.D.: [The study of transformation mechanism of morphology and physiology in the switch from C₃-photosynthesis to Crassulacean acid metabolism of *Dendrobium*]. Pp. 15-41. Huazhong Agricultural University, Wuhan 2014. [In Chinese]
- Lüttge U.: Ecophysiology of crassulacean acid metabolism (CAM). – *Ann. Bot.-London* **93**: 629-652, 2004.
- Mao Z.Y., Zhang W.J.: [A simple method for determining CAM plant.] – *Plant Physiol. Commun.* **35**: 55-56, 1985. [In Chinese]
- O'Leary M.H.: Carbon isotope fraction in plants. – *Phytochemistry* **20**: 553-567, 1981.
- Osmond C. B.: Crassulacean acid metabolism: a curiosity in context. – *Annu. Rev. Plant Phys.* **29**: 379-414, 1978.
- Pierce S., Winter K., Griffiths H.: The role of CAM in high rainfall cloud forests: an in situ comparison of photosynthetic pathways in Bromeliaceae. – *Plant Cell Environ.* **25**: 1181-1189, 2002.
- Ren J.W.: [Study on dynamics of electron transport and switching between different carbon assimilation pathways in three species of *Dendrobium*.] – Ph.D. Thesis, Chin. Acad. Forest. Pp. 61-65. Beijing 2008. [In Chinese]
- Sayed O.H.: Crassulacean acid metabolism 1975-2000, a check list. – *Photosynthetica* **39**: 339-352, 2001.
- Schuiteman A.: *Dendrobium* (Orchidaceae): To split or not to split? – *Garden. Bull. Singapore* **63**: 245-257, 2011.
- Silvera K., Santiago L.S., Cushman J.C. *et al.*: The incidence of crassulacean acid metabolism in Orchidaceae derived from carbon isotope ratios: a checklist of the flora of Panama and Costa Rica. – *Bot. J. Linn. Soc.* **163**: 194-222, 2010.
- Silvera K., Santiago L.S., Cushman J.C. *et al.*: Crassulacean acid metabolism and epiphytism linked to adaptive radiations in the Orchidaceae. – *Plant Physiol.* **149**: 1838-1847, 2009.
- Silvera K., Santiago L.S., Winter K.: Distribution of crassulacean acid metabolism in orchids of Panama: evidence of selection of weak and strong modes. – *Funct. Plant Biol.* **32**: 397-407, 2005
- Silvera K.I.: Crassulacean acid metabolism in tropical orchids: integrating phylogenetic, ecophysiological and molecular genetic approaches. Pp. 1-253. University of Nevada, Reno 2010.
- Sternberg L.O., Deniro M.J., Johnson H.B.: Isotope ratios of cellulose from plants having different photosynthetic pathways. – *Plant Physiol.* **74**: 557-561, 1984.
- Su W.H., Zhang G.F.: [Primary study on photosynthetic characteristics of *Dendrobium nobile*.] – *J. Chin. Med. Mat.* **26**: 157-159, 2003. [In Chinese]
- Szarek S.R., Johnson H.B., Ting I.P.: Drought adaptation in *Opuntia basilaris*. Significance of recycling carbon through crassulacean acid metabolism. – *Plant Physiol.* **52**: 539-541, 1973.
- Ting I.P.: Crassulacean acid metabolism. – *Annu. Rev. Plant Phys.* **36**: 595-622, 1985.
- Vaasen A., Begerow D., Hampp R.: Phosphoenolpyruvate carboxylase genes in C₃, crassulacean acid metabolism (CAM) and C₃/CAM intermediate species of the genus *Chusia*: rapid reversible C₃/CAM switches are based on the C₃ housekeeping gene. – *Plant Cell Environ.* **29**: 2113-2123, 2006.
- Winter K., Holtum J.A.M.: Environment or development? Lifetime net CO₂ exchange and control of the expression of crassulacean acid metabolism in *Mesembryanthemum crystallinum*. – *Plant Physiol.* **143**: 98-107, 2007.
- Winter K., Holtum J.A.M.: How closely do the δ¹³C values of crassulacean acid metabolism plants reflect the proportion of CO₂ fixed during day and night? – *Plant Physiol.* **129**: 1843-1851, 2002.
- Winter K., Wallace B., Stocker G. *et al.*: Crassulacean acid metabolism in Australian vascular epiphytes and some related species. – *Oecologia* **57**: 129-141, 1983.
- Winter K., Garcia M., Holtum J.A.M.: Nocturnal versus diurnal CO₂ uptake: how flexible is *Agave angustifolia*? – *J. Exp. Bot.* **65**: 1695-1703, 2014.
- Zhu G.H., Ji Z.H., Wood J.J. *et al.*: *Dendrobium* Swartz, Nova Acta Regiae Soc. Sci. Upsal., ser. 2, 6: 82. 1799, nom. cons. – In: *Flora of China* **25**: 367-397. 2009.