

Warming alters competition for organic and inorganic nitrogen between co-existing grassland plant species

Thomas M. Kuster · Anna Wilkinson · Paul W. Hill ·
Davey L. Jones · Richard D. Bardgett

Received: 13 September 2015 / Accepted: 7 March 2016 / Published online: 29 March 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract

Introduction Grass species may acquire different forms of nitrogen (N) to reduce competition for the same resources. Climate change influences the availability of soil N and is therefore likely to cause shifts in N forms acquired by plants, thereby affecting their competitive interactions.

Methods We investigated the effects of warming on the uptake of different N forms and competitive interactions of *Festuca ovina* and *Anthoxanthum odoratum* in a pot experiment. The plants were grown either in monocultures or mixture, and at ambient or elevated temperature (+10 °C), and supplied with ¹³C and ¹⁵N isotopes to test for treatment effects on the relative uptake of ammonium, alanine or tri-alanine.

Results Both grass species took up relatively more N supplied as ammonium than as alanine or tri-alanine

when grown under ambient conditions in monoculture. In contrast, when grown in mixtures, *F. ovina* took up the three supplied N forms in equal amounts, whereas *A. odoratum* switched to tri-alanine as the main N form. Under warmed conditions, both species took up the N forms equally, irrespective of competition treatments.

Conclusions We have shown that grass species grown in mixture and under ambient conditions reduce competition by acquiring different N forms. Warming increased the availability of inorganic N in the soil and therefore deregulated the need for differential uptake of N forms.

Keywords Amino acid · Peptide · Nutrient · Coexistence · Niche differentiation

Responsible Editor: Ad C. Borstlap.

Electronic supplementary material The online version of this article (doi:10.1007/s11104-016-2856-7) contains supplementary material, which is available to authorized users.

T. M. Kuster · A. Wilkinson · R. D. Bardgett
Faculty of Life Sciences, Michael Smith Building, The University of Manchester, Oxford Road, Manchester M13 9PT, UK

T. M. Kuster (✉)
Institute for Plant Production Sciences, Agroscope, Schloss 1,
8820 Wädenswil, Switzerland
e-mail: thomaskuster@gmx.ch

P. W. Hill · D. L. Jones
School of Environment, Natural Resources and Geography,
College of Natural Sciences, Bangor University, Gwynedd LL57
2UW, UK

Introduction

Soil nitrogen (N) availability is one of the most important growth-limiting factors in natural or semi-natural grasslands (Vitousek and Howarth 1991). There is growing evidence that increasing temperatures due to global warming will accelerate rates of soil N turnover in these and other temperature-limited ecosystems (Bai et al. 2013; IPCC 2013; Prescott 2010; Zhang et al. 2008), leading to increased soil N availability and a shift in the dominant N form from dissolved organic N (DON) to soluble inorganic N (DIN) (Bai et al. 2013; Rennenberg et al. 2009; Saxe et al. 2001). In addition to changing climate, changes in grassland land use, such as shifts in management intensity or grazing density, also modify microbial communities and rates of soil N turnover, causing shifts in the

availability of different N forms (de Vries et al. 2012; Medina-Roldan et al. 2012), with the amount of DON relative to DIN being greater in low than in high productivity, intensively managed grasslands (Bardgett et al. 2003; Christou et al. 2005; Schimel and Bennett 2004).

It is well established that plant species are able to take up soil N in a range of forms, either as inorganic N, in the form of ammonium (NH_4^+) and nitrate (NO_3^-), or as organic N, in the form of urea, amino acids and peptides (Näsholm et al. 2009; Näsholm and Persson 2001; Sauheitl et al. 2009b; Soper et al. 2011). Although grasses are relatively plastic with regard to their use of different N forms (Falkengren-Grerup et al. 2000; Sauheitl et al. 2009b), it has been suggested that under N limiting conditions grass species acquire contrasting forms of N, which appear to be linked to their growth strategies (Kahmen et al. 2006; Weigelt et al. 2005). This plasticity in acquiring different N forms has been proposed to be a strategy for co-existing plant species to reduce niche overlap, and therefore to avoid competition for the same limiting resource (Ashton et al. 2010; McKane et al. 2002). Results from studies testing for niche partitioning based on chemical forms of N in grasslands, however, are mixed: some report differences in N forms taken up by co-existing grassland plant species (Ashton et al. 2010; Kahmen et al. 2006), whereas others do not (Ashton et al. 2008; Harrison et al. 2007).

Given that climatic conditions are known to regulate the availability of different N forms, it is likely that modified N availability due to warming will also lead to a shift in N forms taken up by plants. Indeed, Warren (2009) reported that *Eucalyptus pauciflora* Sieber ex Spreng. took up more glycine than nitrate at low temperatures, whereas the opposite was true when temperatures were higher due to changed N pool turnover rates. Similarly, in arctic tundra, glycine uptake by herbs was reduced by long-term warming (Sorensen et al. 2008), whereas glycine acquisition by the grass *Deschampsia flexuosa* (L.) was found to increase with warming (Andresen et al. 2009). Given this, our goal was to test how warming impacts the uptake of different N forms by grass species with contrasting life history strategies, and whether this influences their competitive interactions. We focused on two grass species that co-exist in low productivity, semi-natural temperate grassland: the slower-growing species *Festuca ovina* L. and the faster-growing species *Anthoxanthum odoratum* L. (Elberse and Berendse 1993; Ryser and Wahl 2001; Schippers and Olf 2000; Schippers et al. 1999). These species

have previously been shown to differ in their acquisition of organic and inorganic N forms in monoculture. *Festuca rubra* L., as a close relative to *F. ovina*, displays a selective placement in nutrient-rich patches with shorter roots and has been reported to take up relatively more inorganic than organic N, whereas *A. odoratum*, with its longer roots spread more evenly in the soil, relies equally on both forms (Elberse and Berendse 1993; Harrison et al. 2007; Harrison et al. 2008; Mommer et al. 2011; Schippers and Olf 2000; Weigelt et al. 2005).

We hypothesised that: (i) the two grass species preferentially take up N supplied in different forms reflecting their differing life history strategies, and this difference is greater in mixture to avoid competition for soil N; and (ii) at warmer temperatures preferences for N supplied in different forms become less important for *F. ovina* and *A. odoratum* due to increased availability of DIN compared to ambient temperatures. To test these hypotheses, we conducted a factorial pot experiment, in which *F. ovina* and *A. odoratum* were grown either in monocultures or mixtures at both ambient or elevated temperature, and were supplied with ^{13}C and ^{15}N labelled compounds to test the relative uptake of ammonium (as a representative form of inorganic N), alanine (amino acid) or tri-alanine (peptide).

Materials and methods

Experimental setup

We established a pot experiment using field soil collected from a grassland site at Abergwyngregyn, Gwynedd, North Wales, UK (53° 13' 27" N, 4° 00' 50" W, 320 m a.s.l.), as described by Farrell et al. (2011a). Briefly, the selected site is classified as a semi-natural *Agrostis-Festuca* grassland, based on the UK National Vegetation Classification (Rodwell 1992), and is dominated by the grasses *Agrostis canina* L., *Agrostis capillaris* L., *A. odoratum* and *F. ovina*, and the herbs *Potentilla erecta* (L.) Raeusch. and *Galium saxatile* L. The soil is an organic matter rich Cambic Podzol with an acidic pH (4.8) and is representative of a typical semi-natural, sheep-grazed upland grassland in the western United Kingdom (Bardgett et al. 2001). The dissolved N pool is rich in organic N ($301 \pm 74 \text{ mg m}^{-2}$), whereas concentrations of $\text{NH}_4^+\text{-N}$ ($73.4 \pm 36.8 \text{ mg m}^{-2}$) and NO_3^-N ($0.6 \pm 0.5 \text{ mg m}^{-2}$) are lower (data refer to a

depth of 15 cm, published in Wilkinson et al. (2015)). The climate, measured at sea level at a distance of ca. 1 km from the sampling site, is cool and wet with a mean annual air temperature of 10.7 °C, soil temperature of 11 °C (at 10 cm depth) and rainfall of 1250 mm. In spring 2013, soil from the field site was excavated from the rooting zone down to 15 cm depth. Soil was transported back to the laboratory where stones and roots were removed. After passing through a 4 mm sieve, the soil was thoroughly mixed and stored afterwards at 4 °C until the start of the experiment.

We selected two grass species: *A. odoratum* and *F. ovina*. Both species co-exist at the site, although *A. odoratum* is generally more abundant in more productive grasslands, and *F. ovina* is more abundant in lower productivity grasslands (Grime et al. 2007). In April 2013, seeds (Emorsgate Seeds, King's Lynn, UK) of *A. odoratum* and *F. ovina* were germinated in a 1:1 mixture (v:v) of a low fertility compost (No 1; John Innes Manufacturers Association, Reading, UK) and horticulture sand (Keith Singleton Horticulture, Egremont, UK) at ambient temperatures in a greenhouse at The University of Manchester. Due to differences in germination and establishment rates, *A. odoratum* was sown 2 weeks later than *F. ovina* in order to produce uniformly sized seedlings. Trays were watered every second day with tap water without using any additional fertiliser. After 32 (*A. odoratum*) and 46 (*F. ovina*) days, seedlings with an average height of 9 cm were allocated to 3 intra- and interspecific planting treatments, each with two individual plants: i) *F. ovina* monoculture; ii) *A. odoratum* monoculture; and iii) *F. ovina* and *A. odoratum* mixture. Care was taken to ensure that the height of individuals in each of the 192 pots (side length = 9 cm, used height = 7 cm, average soil volume = 0.567 l) was similar. Immediately after planting, pots of each treatment were randomly assigned to two temperatures in controlled growth cabinets (day length 16 h), namely: 12 °C, representing ambient growing season temperature, and 22 °C, representing warming. The ambient temperature refers to an average temperature during growing seasons at the field site (13.7 °C at sea level, implying approximately 12 °C at the field site). Warming of 10 °C was used as an approach to extrapolate the climate sensitivity of N availability and uptake in a model ecosystem. Pots were randomly relocated within cabinets twice per week.

Pots were irrigated with tap water bi-weekly (ambient: 50 ml pot⁻¹ week⁻¹; warming: 100 ml pot⁻¹ week⁻¹,

total dissolved N in tap water <0.4 mg l⁻¹), with differences in irrigation between the two treatments accounting for estimated greater evapotranspiration due to increased temperature and plant biomass in the warmed compared to ambient treatment. The difference in N input through irrigation between the treatments due to the different amount of water (ambient: <0.16 mg pot⁻¹; warming: <0.32 mg pot⁻¹) was negligible compared to total N per pot (approximately 2 g N pot⁻¹). The height of each seedling (longest shoot) was measured weekly.

Isotope labelling and harvest of plant biomass

Labelling of soils to measure uptake of different N forms was performed after 71 days, at a period when shoot height had remained stable for several weeks. Twelve replicate pots of each planting × temperature treatment were randomly allocated to the following three labelling treatments (72 out of 192 pots): i) ¹⁵NH₄Cl (98 % ¹⁵N, Cambridge Isotope Laboratories, Andover, MA, USA); ii) alanine (97–99 % U-¹³C, 97–99 % ¹⁵N, Cambridge Isotope Laboratories); and iii) tri-alanine, (97–99 % U-¹³C, 97–99 % ¹⁵N, CK Gas Products, Ibstock, UK). Nitrate concentration in the original field soil was negligible compared to DON and ammonium (Wilkinson et al. 2015), and therefore, nitrate was not used for labelling. There were 4 replicates for each treatment-labelling combination. The other 120 pots were treated with an unlabelled N solution (18 μmol N pot⁻¹), from which 8 pots were analysed for natural abundance assessments. Each labelling solution (18 μmol N pot⁻¹) was made up of equal concentrations (6 μmol N pot⁻¹ for each N form) of ammonium, alanine and tri-alanine, in which one of the three N forms was isotopically labelled. This enabled us to test for preferential uptake by individual plant species and soil microbes (Harrison et al. 2007; Weigelt et al. 2005). The use of dual-labelled ¹³C¹⁵N compounds is generally, but not unequivocally, considered to be a good indication whether amino acids and peptides such as alanine and tri-alanine are taken up by plants directly as organic N, or as inorganic N after microbial mineralisation, as confirmed by enrichment of plant tissue with both ¹³C and ¹⁵N (Näsholm et al. 1998). The amount of N added to each pot was considered to be sufficient to allow for detection of ¹³C and ¹⁵N within plant and microbial biomass, but keeping the possible N fertilisation effect on plant growth to a minimum (18 μmol N pot⁻¹ (0.3 kg N ha⁻¹) < N_{H20} = 490 μmol N pot⁻¹). Within

each pot, the labelling solution (20 ml) was injected at 5 different locations, equally distributed over the soil depth, using a glass syringe (S Murray & Co, Surrey, UK). Pots were randomly labelled over a period of 4 days.

Three hours after labelling, pots were destructively harvested and plants were separated from the soil. A chase period of 3 h was chosen to reduce plant uptake of recycled mineralised organic N, but to provide sufficient time to detect $^{13}\text{C}^{15}\text{N}$ in roots and shoots (Warren 2012). Roots were first washed with deionised water and then rinsed with 0.5 M CaCl_2 to remove ^{13}C and ^{15}N in the apoplast and sorbed to the cell wall. Roots of the two species in the mixed treatment were distinguished from each other by their colour. Root, shoot and soil samples were dried at 65 °C for 2 days prior to grinding (MM 400, Retsch, Haan, Germany). Root and shoot samples of the two individuals grown in monocultures were pooled prior to grinding, whereas for mixed treatments both individuals were analysed separately.

Root and, shoot extract samples were analysed for $^{12/13}\text{C}$ and $^{14/15}\text{N}$ concentrations at the NERC Life Sciences Mass Spectrometer Facility, Centre for Ecology and Hydrology, Lancaster, UK, (precision for working standards better than 0.46 ‰ (^{13}C) and 6.92 ‰ (^{15}N)). Samples were combusted in a Carlo Erba NA1500 elemental analyser (Thermo Scientific, Waltham, MA, USA). The resultant CO_2/N_2 from combustion and reduction was analysed for $\delta^{13}\text{C}/^{15}\text{N}$ using an isotope ratio mass spectrometer (IRMS; Dennis Leigh Technologies, Sandbach, UK). $^{13}\text{C}^{15}\text{N}$ excess values were calculated by using formulas (1) and (2).

$$R_{\text{sample}} = [(\delta^{13}\text{C}/1000) + 1] * R_{\text{PDB}} \quad (1)$$

where R is the ratio of ^{13}C /of ^{15}N to ^{12}C /to ^{14}N and R_{PDB} is the natural abundance standard for C and N.

$$\text{Atom}\% = (R/R + 1) * 100 \quad (2)$$

Atom % excess values were calculated by subtracting control atom % values from treatment atom % values. Natural abundance levels of ^{13}C in our samples were highly variable. We therefore used the lowest natural abundance atom % value to calculate ^{13}C excess values.

Soil nutrients, microbial biomass and root length

Immediately after plants were harvested, fresh soil samples were extracted with deionised water (1:7.1 w/v

soil:extractant; extraction time = 10 min) to measure total dissolved N and inorganic N as either nitrate (NO_3^-) or ammonium (NH_4^+). Extracts were measured with an AutoAnalyzer 3 (SEAL Analytical, Fareham, UK). DON was calculated after subtracting water-soluble inorganic N from total water-soluble N. Dissolved organic carbon (DOC) was measured in water extracts using a TOC-L analyser (Shimadzu, Kyoto, Japan). For determining microbial C (C_{mic}) and N (N_{mic}), chloroform-fumigated (fumigation time = 24 h, amylene-stabilised CHCl_3 , Fisher Scientific, Waltham, MA, USA) and non-fumigated soil samples were extracted with 0.5 M K_2SO_4 (1:2.5 w/v soil:extractant; extraction time = 60 min) (Brookes et al. 1985), and total soluble organic carbon and N in K_2SO_4 -extracts were measured with a TOC-L (Shimadzu, Kyoto, Japan) and an AutoAnalyzer 3 (SEAL Analytical, Fareham, UK), respectively. Microbial biomass was calculated as the respective differences between fumigated and non-fumigated samples. The differences were divided by the corrections factors $k_{\text{EC}} = 0.35$ and $k_{\text{EN}} = 0.50$ to estimate C_{mic} and N_{mic} (Carter 2008). In labelled soil samples, pH was measured in 0.01 M CaCl_2 (FE20, Mettler-Toledo, Schwerzenbach, Switzerland). Root samples from pots that were not used for the labelling experiment were analysed for their diameter and length using an Epson Expression 11,000 XL, scanner (Nagano, Japan) and WinRHIZO Pro 2013a (Regent Instruments Inc., Quebec, CA).

Statistical analysis

Data were analysed after log-transformation by ANOVA using a linear model (significant at $P < 0.05$) in R 3.02 (R Development Core Team, Vienna, AT). The initial shoot height (analysis of root and shoot biomass) and final biomass ($^{12/13}\text{C}^{14/15}\text{N}$ values) were included in the models to account for differences between pots. Selected differences between treatments and soils were pair-wise tested using contrasts based on *t*-tests (significant at $P < 0.05$).

Results

Soil N availability and microbial biomass

Concentrations of inorganic N (NH_4^+ and NO_3^-) were influenced by planting and warming treatments (Table 1). Concentrations of NH_4^+ ($F_{(2,181)} = 36.6$,

Table 1 Soil properties at the end of the experiment

	Ambient			Warming			F-values		
	<i>F. ovina</i> monoculture	<i>A. odoratum</i> monoculture	Mixture	<i>F. ovina</i> monoculture	<i>A. odoratum</i> monoculture	Mixture	T	P	T × P
SWC	*41.1 ± 1.0 ^a	39.2 ± 1.2 ^a	38.0 ± 1.0 ^a	*45.0 ± 1.6 ^a	31.2 ± 1.7 ^b	34.0 ± 2.1 ^b	9.6**	17.6***	8.3***
C _{tot}	90.1 ± 1.6 ^a	91.8 ± 0.7 ^a	92.5 ± 0.8 ^a	91.4 ± 0.6 ^a	91.2 ± 1.0 ^a	92.2 ± 1.9 ^a	<0.1	0.9	0.4
DOC	62.1 ± 5.7 ^a	73.3 ± 4.8 ^a	63.4 ± 5.1 ^a	56.4 ± 4.8 ^a	65.3 ± 2.9 ^b	59.2 ± 4.3 ^{ab}	1.5	4.7*	0.1
N _{tot}	7.8 ± 0.1 ^a	7.8 ± 0.0 ^a	7.9 ± 0.0 ^a	7.9 ± 0.1 ^a	7.8 ± 0.1 ^a	7.9 ± 0.1 ^a	<0.1	0.8	0.5
NH ₄ ⁺	*2.7 ± 0.7 ^a	*1.3 ± 0.1 ^b	*1.6 ± 0.2 ^a	*10.9 ± 1.7 ^a	*2.3 ± 0.3 ^b	*3.5 ± 0.5 ^b	88.0***	36.6***	8.7***
NO ₃ ⁻	*12.0 ± 2.0 ^a	*2.5 ± 0.5 ^b	*7.2 ± 1.6 ^a	*57.9 ± 4.4 ^a	*6.5 ± 1.2 ^b	*14.9 ± 2.9 ^c	59.9***	44.9***	7.7***
DON	3.9 ± 0.3 ^a	*5.5 ± 1.0 ^a	4.0 ± 0.3 ^a	2.9 ± 0.6 ^a	*3.5 ± 0.3 ^a	4.1 ± 0.2 ^a	5.2*	2.2	2.1
Mic _C	*1.9 ± 0.1 ^a	*1.9 ± 0.1 ^a	1.6 ± 0.1 ^b	*1.2 ± 0.1 ^a	*1.4 ± 0.1 ^b	1.7 ± 0.1 ^b	26.6***	2.1	11.8***
Mic _N	*0.29 ± 0.01 ^a	*0.30 ± 0.02 ^a	0.28 ± 0.01 ^a	*0.23 ± 0.01 ^a	*0.25 ± 0.01 ^{ab}	0.26 ± 0.01 ^b	18.5***	0.7	2.9 ^(*)

Values are mean ± SE. An asterisk * indicates a significant difference between warming treatments within the same competition treatment (all $P < 0.05$). Different letters indicate significant differences between competition treatments within the same warming treatment. SWC: soil water concentration (%), $n = 32$, residual $df = 183$), C_{tot}: total carbon (mg g^{-1} , 8, 42), DOC: dissolved organic C ($\mu\text{g g}^{-1}$, 32, 186), N_{tot}: total nitrogen (mg g^{-1} , 12, 66), NH₄⁺: ammonium ($\mu\text{g g}^{-1}$, 32, 181), NO₃⁻: nitrate ($\mu\text{g g}^{-1}$, 32, 181), DON: dissolved organic nitrogen ($\mu\text{g g}^{-1}$, 32, 181), Mic_C: microbial carbon (mg g^{-1} , 32, 167), Mic_N: microbial nitrogen (mg g^{-1} , 32, 167). Statistical analyses (F -values): Effects of temperature (T, $df = 1$), planting (P, $df = 2$) and their interactions (T × P, $df = 2$), levels of significances (***: $P < 0.001$, **: $P < 0.01$, *: $P < 0.05$, (*): $P < 0.1$)

$P < 0.001$) and NO₃⁻ ($F_{(2,181)} = 44.9$, $P < 0.001$) were lowest in soil planted with *A. odoratum*, followed by mixtures and *F. ovina* monocultures. Soil concentrations of NH₄⁺ ($F_{(1,181)} = 88.0$, $P < 0.001$) and NO₃⁻ ($F_{(1,181)} = 59.9$, $P < 0.001$) were greater in the warming than in the ambient treatment. Dissolved organic N (DON) was lower in the ambient treatment than the warming treatment ($F_{(1,181)} = 5.2$, $P = 0.023$). Pair-wise comparisons for DON were, however, only significant in *A. odoratum* monocultures ($P = 0.007$) and not in the other planting treatments (*F. ovina* monocultures: $P = 0.171$; mixtures: $P = 0.895$). Dissolved organic carbon (DOC) was changed by the planting treatment ($F_{(2,186)} = 4.7$, $P = 0.010$): DOC concentrations were, or tended to be greater in *A. odoratum* than in *F. ovina* monocultures (ambient: $P = 0.054$; warming: $P = 0.028$). There was, however, no warming effect on soil DOC, and neither total soil carbon (C_{tot}), nitrogen (N_{tot}) or pH were affected by the warming and planting treatments (Table 1). Soil water concentration at the end of the experiment was greatest in warmed *F. ovina* monoculture, whereas no differences between the other treatments were observed.

Both microbial biomass C ($F_{(1,167)} = 26.6$, $P < 0.001$) and N ($F_{(1,167)} = 4.1$, $P = 0.045$) were greater in the ambient than in the warming treatment (Table 1).

However, effects of warming on microbial biomass C (temperature × planting: $F_{(2,167)} = 11.8$, $P < 0.001$) and N (temperature × planting: $F_{(1,167)} = 5.5$, $P = 0.005$) varied with planting design: under ambient conditions, microbial biomass C was lowest in mixtures, whereas under warmed conditions it was smallest in *F. ovina* monocultures. Similarly, microbial biomass N under warming was lower in *F. ovina* monocultures than in mixtures, but not different from *A. odoratum* monocultures. We observed no differences in microbial N between the planting treatments under ambient conditions.

Root and shoot biomass

Elevated temperature on average doubled the shoot biomass of *A. odoratum*, whereas warming only marginally influenced shoot biomass of *F. ovina* (temperature × species: $F_{(1,247)} = 91.9$, $P < 0.001$, Fig. 1a). The planting treatment only affected shoot biomass of *A. odoratum* in the warming treatment (temperature × planting: $F_{(1,247)} = 4.5$, $P = 0.035$): *A. odoratum* shoot biomass per plant was 50 % higher in mixtures than in monocultures (competition ratio (CR) = 1.5 ± 0.1). However, total shoot biomass per pot (2 plants) in warmed *A. odoratum* monocultures did not differ from the total biomass of the two species in mixtures




Fig. 1 **a** Average root and shoot biomass per individual (g, \pm SE, $n = 32$), separately shown for each temperature and planting treatment. Total root length is shown in Fig. S1. **b** Average nitrogen (N) concentrations in root and shoot biomass (% , \pm SE, $n = 12$). Please note that the y-axis (shoot N) starts at 2 %. **a & b** Values of the two individuals in the monoculture treatment were

pooled prior analysis. Significant ($P < 0.05$) pair-wise comparisons are indicated by *: difference between monoculture and mixed treatments within the same species and temperature treatment; s: difference between species within the same temperature and competition treatment; w: difference between temperature treatments within the same species and competition treatment

($P = 0.745$). Planting treatment had no effect on the shoot biomass of *A. odoratum* under ambient temperature ($CR = 0.9 \pm 0.1$), or on shoot biomass of *F. ovina* under ambient ($CR = 1.0 \pm 0.1$) or warmed conditions ($CR = 1.2 \pm 0.1$).

Warming decreased root biomass of *F. ovina* in monoculture and mixtures, but it had no effect on root biomass of *A. odoratum* in monoculture, although it increased root biomass of this species in mixtures (temperature \times species: $F_{(1,247)} = 49.7$, $P < 0.001$, Fig. 1a). As a result, under elevated temperatures, root biomass of *A. odoratum* was more than four times greater than of *F. ovina*, whereas root biomass did not differ between the two species under ambient conditions (Fig. 1a). Warming decreased the root:shoot ratio of the test species ($F_{(1,247)} = 178.9$, $P < 0.001$): the effect of temperature on the root:shoot ratio of *F. ovina* was greater than on that of *A. odoratum* (temperature \times species: $F_{(1,247)} = 39.8$, $P < 0.001$), leading to a significantly higher root:shoot ratio of *A. odoratum* than of *F. ovina* in the warming treatment. There was no significant planting effect on root biomass or root:shoot ratio of either species ($F_{(1,247)} = 1.6$, $P = 0.201$, $F_{(1,247)} = 0.8$, $P = 0.383$). As with root biomass, root length of *F. ovina* was least in the warming than in the ambient treatment, whereas root length of *A. odoratum* grown in mixtures was greater under warming than ambient conditions (temperature \times planting: $F_{(1,143)} = 55.4$, $P < 0.001$, Fig. S1A). No warming effect on root length was observed in *A. odoratum* monocultures; hence, root length

of *A. odoratum* was greater than of *F. ovina*, but only under warmed conditions.

N concentrations in root and shoot

Temperature effects on shoot N differed between the two grass species (temperature \times species: $F_{(1,87)} = 37.1$, $P < 0.001$): shoot N in *A. odoratum* was greater under warming than under ambient temperature, whereas for *F. ovina* no effect of warming was detected (Fig. 1b). Planting design also influenced the two species differently (planting \times species: $F_{(1,87)} = 19.4$, $P < 0.001$). Although pair-wise comparisons were not significant, shoot N in *A. odoratum* tended to be greater in mixtures than in monocultures, whereas it was the other way around in *F. ovina*. Hence, shoot N concentrations under ambient and monoculture conditions were higher in *F. ovina* than in *A. odoratum* ($P = 0.004$), whereas N concentrations were lower in *F. ovina* than in *A. odoratum* in warmed mixture ($P = 0.001$). In general, root N concentrations were greater under elevated than under ambient temperature ($F_{(1,87)} = 9.1$, $P < 0.001$, Fig. 1b). However, pair-wise comparisons revealed that this response to warming was only significant in *A. odoratum* roots grown in mixtures ($P = 0.015$).

$^{13}\text{C}^{15}\text{N}$ excess values in root and shoot biomass

Enrichment of plant material, measured as absolute ^{15}N excess values, differed strongly between the two grass

species (roots: $F_{(1,71)} = 21.1$, $P < 0.001$; shoots: $F_{(1,71)} = 72.9$, $P < 0.001$). On average, ^{15}N excess values in roots and shoots of *A. odoratum* were higher than in those of *F. ovina*, which is indicative of greater uptake of all N forms (Fig. 2). Differences in ^{13}C excess values between the two species, however, were only weakly or not significant (roots: $F_{(1,47)} = 3.6$, $P = 0.064$; shoots: $F_{(1,71)} = 5.8$, $P = 0.020$), although there was a trend towards higher ^{13}C concentrations in *A. odoratum* than *F. ovina* (Table 3).

Plant uptake of N was affected by chemical N form (roots: $F_{(2,71)} = 18.1$, $P < 0.001$; shoots: $F_{(2,71)} = 22.2$, $P < 0.001$), planting design (roots: $F_{(1,71)} = 10.0$, $P = 0.002$; shoots: $F_{(1,71)} = 22.0$, $P < 0.001$) and warming treatment (roots: $F_{(2,71)} = 6.5$, $P = 0.013$; shoots: $F_{(2,71)} = 57.1$, $P < 0.001$). Most interestingly, planting treatment influenced the uptake of N forms by *A. odoratum* under ambient conditions: in monoculture, ^{15}N excess rates in *A. odoratum* roots and shoots were greater for ammonium than alanine (roots; shoots: $P < 0.001$; $P < 0.001$) or tri-alanine ($P = 0.066$; $P = 0.023$), whereas in mixture uptake of N supplied as tri-alanine was greater than as ammonium ($P = 0.049$; $P = 0.860$) or alanine ($P = 0.011$; $P = 0.005$). This shift in N forms taken up by *A. odoratum* can mainly be deduced from a smaller ammonium uptake in mixture

than in monoculture ($P < 0.001$; $P = 0.016$), whereas we observed no difference in uptake of N supplied as tri-alanine between the planting treatments. In *F. ovina* roots and shoots grown at ambient conditions, differences between N forms were less obvious than for *A. odoratum*. In monoculture, uptake of N supplied as alanine was less than for ammonium ($P < 0.001$; $P < 0.001$) and tri-alanine ($P = 0.020$; $P = 0.083$). In mixture, we observed no difference in uptake of different N forms on the basis of ^{15}N excess in *F. ovina* roots, but values in shoots were greater when plants were labelled with ammonium than with alanine ($P = 0.001$) or tri-alanine ($P = 0.032$). Recovery rates in roots of the applied ammonium and alanine were the same for the two species in mixture, whereas more tri-alanine was found in roots of *A. odoratum* than in roots of *F. ovina* ($P = 0.012$, Table 2). No differences in recovery rates in roots between the two species were found when they were grown in monocultures. Both in monocultures and mixtures, all labelling solutions were found to a higher extent in *A. odoratum* shoots than in *F. ovina* shoots. Over all, the conclusions gained from comparing the relative numbers presented as recovery rates are, however, the same as found when using excess values. Root uptake of ^{13}C was also affected by the chemical form ($F_{(1,47)} = 16.5$, $P < 0.001$); under ambient conditions ^{13}C

Fig. 2 Average ^{15}N excess rates in root (a) and shoot biomass (b) after a chasing period of 3 h ($\mu\text{mol g}^{-1}$, \pm SE, $n = 4$), separately shown for NH_4^+ , alanine and tri-alanine. Differences between the applied tracer solutions within a given treatment combination (column) are indicated by different letters (all $P < 0.05$). Average ^{13}C excess values are shown in Table 3. Please note the different scales between the two species




Table 2 Average recovery rates of the applied labelling solutions in roots and shoots after a chasing period of 3 h (%), SE, $n = 4$), separately shown for ammonium, alanine and tri-alanine

	<i>F. ovina</i>			<i>A. odoratum</i>		
	Ammonium	Alanine	Tri-alanine	Ammonium	Alanine	Tri-alanine
Roots						
Ambient monoculture	2.3 ± 1.0 ^a	0.6 ± 0.1 ^b	1.1 ± 0.2 ^{ab}	3.9 ± 0.8 ^a	1.1 ± 0.2 ^b	2.1 ± 0.5 ^{ab}
Ambient mixture	1.5 ± 0.2 ^a	0.9 ± 0.3 ^a	*0.8 ± 0.4 ^a	0.8 ± 0.4 ^a	0.6 ± 0.1 ^a	*2.8 ± 0.6 ^b
Warming monoculture	*0.1 ± 0.0 ^a	*0.1 ± 0.0 ^a	*0.4 ± 0.1 ^b	*1.9 ± 0.5 ^a	*1.9 ± 0.5 ^a	*1.9 ± 0.1 ^a
Warming mixture	*0.2 ± 0.1 ^a	*0.3 ± 0.2 ^a	*0.3 ± 0.1 ^a	*3.4 ± 1.1 ^a	*1.2 ± 0.5 ^b	*2.1 ± 1.2 ^{ab}
Shoots						
Ambient monoculture	*1.4 ± 0.5 ^a	*0.3 ± 0.0 ^b	*0.4 ± 0.1 ^{ab}	*6.2 ± 1.5 ^a	*1.1 ± 0.1 ^b	*2.1 ± 0.6 ^b
Ambient mixture	*0.6 ± 0.1 ^a	*0.2 ± 0.1 ^b	*0.3 ± 0.1 ^{ab}	*1.9 ± 0.6 ^a	*0.8 ± 0.3 ^b	*2.4 ± 0.6 ^a
Warming monoculture	*0.2 ± 0.0 ^{ab}	*0.2 ± 0.1 ^a	*0.5 ± 0.1 ^b	*2.2 ± .8 ^a	*1.8 ± 0.5 ^a	*2.1 ± 0.3 ^a
Warming mixture	*0.2 ± 0.0 ^{ab}	*0.1 ± 0.0 ^a	*0.2 ± 0.0 ^b	*5.9 ± 2.0 ^a	*1.5 ± 0.3 ^a	*2.6 ± 1.0 ^a

Differences between the applied tracer solutions within a given treatment combination are indicated by different letters. An asterisk * indicates a significant difference between species within the same labelling solution and competition treatment (all $P < 0.05$)

excess values were higher when plants were labelled with tri-alanine than with alanine (Table 3).

Warming changed the observed planting effects on ¹⁵N uptake under ambient conditions (temperature × form in roots: $F_{(2,71)} = 3.0$, $P = 0.056$; shoots: $F_{(2,71)} = 6.6$, $P = 0.002$); in general, we detected no differences in ¹⁵N and ¹³C excess values in both species between the applied N forms under warmed conditions (Fig. 2, Table 3). As an exception to this pattern, ¹⁵N excess values for *F. ovina* roots in monoculture were greater for tri-alanine than ammonium ($P = 0.033$) or alanine ($P = 0.023$), and for *A. odoratum* shoots, ¹⁵N excess values were greater for ammonium than alanine ($P = 0.033$). Due to the differences in biomass (Fig. 1),

higher recovery rates of the applied labelling solutions were found under elevated temperatures in roots and shoots of *A. odoratum* compared to *F. ovina* (Table 2).

We found significant correlations between ¹³C and ¹⁵N excess values in *A. odoratum* roots for alanine ($R^2 = 0.287$, $P = 0.027$) and tri-alanine ($R^2 = 0.401$, $P = 0.011$, Fig. 3). The slope of the alanine correlation line ($m = 0.9$) was slightly steeper than that of tri-alanine ($m = 0.5$). We estimate that about a third to a sixth of alanine and tri-alanine were taken up directly, respectively. This indicates that direct uptake of alanine was greater than for tri-alanine, that the proportion ¹³C lost in plant respiration was greater when acquired as tri-alanine than when acquired as alanine, or that the C

Table 3 Mean ¹³C excess values (nmol ¹³C excess g⁻¹) in roots and shoots, separately shown for the ¹³C labelling solutions alanine and tri-alanine. Values are mean ± SE, $n = 4$

	Ambient				Warming			
	<i>F. ovina</i> monoculture	<i>A. odoratum</i> monoculture	<i>F. ovina</i> mixture	<i>A. odoratum</i> mixture	<i>F. ovina</i> monoculture	<i>A. odoratum</i> monoculture	<i>F. ovina</i> mixture	<i>A. odoratum</i> mixture
Roots								
Alanine	279 ± 49 ^a	*235 ± 60 ^a	*129 ± 39 ^b	259 ± 50 ^a	439 ± 49 ^a	558 ± 113 ^a	348 ± 91 ^a	440 ± 109 ^a
Tri-alanine	474 ± 35 ^a	*475 ± 23 ^a	*268 ± 64 ^a	487 ± 67 ^a	567 ± 46 ^a	702 ± 83 ^a	397 ± 71 ^a	585 ± 103 ^a
Shoots								
Alanine	190 ± 82 ^a	269 ± 47 ^a	55 ± 111 ^a	310 ± 168 ^a	298 ± 41 ^a	428 ± 74 ^a	221 ± 149 ^a	324 ± 78 ^a
Tri-alanine	168 ± 50 ^a	292 ± 75 ^a	189 ± 134 ^a	212 ± 59 ^a	347 ± 46 ^a	475 ± 63 ^a	235 ± 58 ^a	347 ± 84 ^a

Different letters indicate significant differences between species and competition treatments within the same warming treatment and N form (all $P < 0.05$). An asterisk * indicates a significant difference between N forms within a given treatment




Fig. 3 Relationship between ^{13}C and ^{15}N excess values in roots of *A. odoratum* and *F. ovina*, separately shown for alanine (open circles) and tri-alanine (closed circles). Broken (alanine: $R^2 = 0.287$, $P = 0.027$) and solid lines (tri-alanine: $R^2 = 0.401$, $P = 0.011$) show significant regressions between the excess of both

isotopes in *A. odoratum* roots. The regressions in *F. ovina* roots were not significant (alanine: $R^2 = 0.022$, $P = 0.557$; tri-alanine: $R^2 < 0.001$, $P = 0.946$). The dotted lines show the molar ^{13}C : ^{15}N ratios for the nitrogen sources injected (3:1)

and N from the compounds partitioned differently between roots and shoots. But given that these relationships are only weakly significant, these numbers need to be interpreted with care. In *F. ovina* roots, we observed no correlations between ^{13}C and ^{15}N excess values (alanine: $R^2 = 0.022$, $P = 0.557$; tri-alanine: $R^2 < 0.001$, $P = 0.946$). The slopes of the correlation lines separately calculated for each planting and warming treatment did not differ from the patterns described above.

Discussion

The aim of this study was to test for the effects of warming on the uptake of different N forms and competitive interactions of two grass species of temperate grasslands with contrasting functional traits. Our first hypothesis was that the two grass species take up different forms when grown in monocultures, and this difference is greater in mixture to avoid competition for soil N. In contrast to this hypothesis, and to previous studies on inorganic and organic N uptake (Harrison et al. 2007; Harrison et al. 2008; Weigelt et al. 2005), we found that both *F. ovina* and *A. odoratum* took up more N supplied as ammonium than as alanine when grown in monocultures. When grown in mixture, however, *A. odoratum*, but not *F. ovina*, switched from taking up more N from ammonium than alanine, to greater uptake of N supplied as tri-alanine than as alanine or ammonium. The difference in N uptake between these species in mixture is reflected in their functional root traits: *Festuca* is known to place roots selectively in nutrient-rich hotspots, whereas *Anthoxanthum* spreads its roots more evenly

in soil, allowing uptake of a greater variety of N forms (Mommer et al. 2011). This suggests that when grown in mixture, *F. ovina* was more competitive than *A. odoratum* in taking up the same N form as in monoculture, thereby reducing *A. odoratum*'s ammonium uptake. Competition for N between plants and microbes was presumably strong in both monocultures and mixtures and, therefore, *A. odoratum* could not compensate the reduced ammonium uptake by acquiring more N supplied as alanine or tri-alanine. Moreover, we exclude that competition between microbes and plants explains the decreased ammonium uptake by *A. odoratum* as this would likewise have affected ammonium uptake by *F. ovina*. Our data, therefore, suggest that this shift in N uptake by *A. odoratum* was mainly induced by a lower competitiveness for ammonium in comparison with *F. ovina*, which lends support to the idea that acquisition of different N forms contributes to coexistence of competing grass species (Ashton et al. 2010; Kahmen et al. 2006; McKane et al. 2002).

As hypothesised, we found that warming changed N use by the two plant species, in that we detected no difference in uptake of the different N forms when they were grown in mixture compared to monocultures in this treatment. It is possible that increased soil inorganic N availability under warming compensated for the need for niche differentiation on the basis of N form, which was detected in mixtures under ambient conditions. Indeed, nitrate and ammonium concentrations were greater in the warming than ambient treatment, which is likely to be due to accelerated organic matter turnover in this organic-rich grassland soil (Bai et al. 2013; Prescott 2010; Rennenberg et al. 2009; Zhang et al. 2008). An alternative mechanism is that warming

influenced the competitiveness of the two grass species, which might have weakened in the requirement for niche differentiation; whereas under ambient conditions the biomass of the two species was similar, *A. odoratum* clearly outcompeted *F. ovina* in the warming treatment. In a study conducted by Schippers and Olff (2000), *A. odoratum* was still more vigorous than *F. ovina* at 15 °C, indicating that the optimum temperature of *F. ovina* is rather closer to 12 °C than to 22 °C. The lower root:shoot ratio and plant N concentrations of *F. ovina* compared to *A. odoratum* indicate that the differences in competitiveness between our test species can be related to a more effective nutrient uptake by *A. odoratum* compared to *F. ovina* in the warmed treatment (Mommer et al. 2011). Otherwise, the low root biomass of *F. ovina* under warming might have been a consequence of the high soil water availability in the *F. ovina* monoculture relative to *A. odoratum* and mixtures. This would mean that due to sufficient water availability in the topsoil there was no need for *F. ovina* to allocate resources to root growth and hence *F. ovina* was likely less competitive in taking up nutrients compared to *A. odoratum*. With its higher root density, *A. odoratum* is likely to be also more competitive under water-limiting conditions, as predicted to increase in frequency with climate change (IPCC 2013); this question, however, was not tested in our experiment and needs further investigation. It is possible that, in the long term, niche partitioning on the basis of uptake of different forms of N will occur in the real world under warming, especially due to acclimatisation of microbial activity and increased plant biomass production (Lu et al. 2013; Luo et al. 2001) or immigration of other species (Klanderud and Birks 2003; Parolo and Rossi 2008). We therefore conclude, in accordance with our second hypothesis, that warming reduces the need for niche differentiation on the basis of N form in grass species, at least in the short timescale of our study.

Even though our data show how competition and temperature influence the uptake of N forms by *F. ovina* and *A. odoratum*, the applied $^{13}\text{C}^{15}\text{N}$ labelling technique has some limitations. First, it is possible that N forms other than those we supplied to soil might have also been important for plant nutrition. Unlike in the field (Wilkinson et al. 2015), concentrations of nitrate were higher than ammonium or DON in soil of the present experiment. We found that soil nitrate concentrations were reduced under ambient conditions by the presence of *A. odoratum*, indicating that nitrate was a

significant part of nutrition for *A. odoratum*. Soil concentrations of nitrate in mixtures, however, suggest that *A. odoratum* did not increase its nitrate acquisition when grown alongside with *F. ovina*. Hence, even though *A. odoratum* may have taken up a significant amount of nitrate, our conclusions, gained from the reduced ammonium uptake in mixture compared to monoculture, would not be different. Second, correlations between ^{13}C and ^{15}N excess values in *A. odoratum* roots and $^{13}\text{C}^{15}\text{N}$ excess values in microbes may indicate that a higher fraction of tri-alanine, compared to alanine, was first mineralised then taken up as inorganic N, as similarly reported by Farrell et al. (2013). We presume, however, that direct uptake of tri-alanine was nevertheless an important source for plant nutrition: on the one hand we found higher plant ^{13}C excess values for tri-alanine than for alanine, indicating that direct uptake of the peptide was, in absolute numbers, higher than of the monomer; on the other hand, differences between ^{13}C and ^{15}N correlations might be explained by faster within-plant mineralisation of tri-alanine compared to alanine (Hill et al. 2011; Warren 2012). In other words, residual carbon, including ^{13}C , might have been respired to a higher extent when applied as peptide than as amino acid, resulting in a smaller $^{13}\text{C}^{15}\text{N}$ ratio. To reduce such uncertainties about direct uptake of labelled isotopes in future experiments, the application of other techniques might be helpful, such as compound-specific stable isotope measurements (Sauheitl et al. 2009a), position-specific labeling (Apostel et al. 2013) and the use of ^{14}C -labelled isotopes (Hill et al. 2013). However, all available techniques are subject to some caveats and assumptions. Third, pool dilution of applied labelling solutions has to be taken into account when interpreting ^{13}C and ^{15}N uptake in plant samples (Jones et al. 2005). We neither have any data on alanine or tri-alanine concentrations in our pots nor do we know the specificity of the transporters in the roots. However at the location where we collected field soil, concentrations of alanine, tri-alanine and other amino acids and peptides competing for the same root transporters were smaller than those of ammonium (Farrell et al. 2011a; Farrell et al. 2011b). Hence, the chance of a plant root to take up labelled ammonium was smaller in comparison with labelled N supplied in organic forms. Otherwise, considering the faster turnover rates of amino acids and peptides comparing to ammonium, plants have much more capacity to take up $^{15}\text{N}\text{-NH}_4^+$ during the labelling period. Taking pool dilution into account by multiplying

soil N pools in the field reported by Farrell et al. (2011b) by ^{15}N excess values recorded within root or shoot tissue in the present experiment, we estimate that 30–100× more $^{15}\text{N-NH}_4$ was recovered in plant material than the tested organic ^{15}N forms. Likely, differences in N uptake in ambient monocultures would be even more distinct when considering pool dilution, whereas shifts in N uptake by *A. odoratum* grown in mixture would be less obvious. However, as this correction would likewise apply for both monocultures and mixture, the relative difference in ammonium uptake would not be different and hence, our main conclusions from this experiment are still likely to be true. We recognize that these data do not necessarily reflect the N pools in our experiment, but they provide an approximation of the likely level of dilution commonly found in the field.

Conclusions

Our data show that grass species grown in mixture and under ambient conditions reduce competition by taking up different N forms. Thereby, N supplied in organic forms as amino acids and peptides can play a major role for plant nutrition. Hence, the possibilities for a plant species to create its own niche are manifold and may include intricacies such as acquiring different N forms. Increased availability of inorganic N due to warming deregulated the need for differential uptake of N forms. Hence, we conclude that uptake of different N forms is mainly important at nutrient-limiting conditions. Besides taking up different N forms, grass species have also been shown to coexist through spatiotemporal shifts in nutrient acquisition (McKane et al. 1990; Pomon et al. 2007). Whereas we exclude spatiotemporal shifts in N uptake as a source for niche differentiation in the present study, these other strategies might explain why in some field studies niche differentiation by taking up different N forms has been reported (Ashton et al. 2010; Kahmen et al. 2006), whereas in others it has not (Ashton et al. 2008; Harrison et al. 2007).

Acknowledgments We thank Helen Grant and Andy Stott from the NERC Life Sciences Mass Spectrometer Facility, Centre for Ecology and Hydrology, Lancaster, UK for conducting the stable isotope measurements. For funding these measurements, we acknowledge the National Environment Research Council NERC (Proposal no. CEH_L_069_05_2013). We thank Corina Del Fabbro, Deborah Ashworth, Hayley Craig, David Grantham, Richard Preziosi and Simon Turner for their help in performing

the experiment. We acknowledge Corina Del Fabbro for her support in data analysis. Financial support was provided by the Swiss National Science Foundation SNSF (Proposal no. 145973) and BBSRC (Grant Reference BB/I009000/2).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Andresen LC, Michelsen A, Jonasson S, Beier C, Ambus P (2009) Glycine uptake in heath plants and soil microbes responds to elevated temperature, CO_2 and drought. *Acta Oecol* 35:786–796
- Apostel C, Dippold M, Glaser B, Kuzyakov Y (2013) Biochemical pathways of amino acids in soil: assessment by position-specific labeling and ^{13}C -PLFA analysis. *Soil Biol Biochem* 67:31–40
- Ashton IW, Miller AE, Bowman WD, Suding KN (2008) Nitrogen preferences and plant-soil feedbacks as influenced by neighbors in the alpine tundra. *Oecologia* 156:625–636
- Ashton IW, Miller AE, Bowman WD, Suding KN (2010) Niche complementarity due to plasticity in resource use: plant partitioning of chemical N forms. *Ecology* 91:3252–3260
- Bai E, Li S, Xu W, Li W, Dai W, Jiang P (2013) A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics. *New Phytol* 199:441–451
- Bardgett RD, Jones AC, Jones DL, Kemmitt SJ, Cook R, Hobbs PJ (2001) Soil microbial community patterns related to the history and intensity of grazing in sub-montane ecosystems. *Soil Biol Biochem* 33:1653–1664
- Bardgett RD, Streeter TC, Bol R (2003) Soil microbes compete effectively with plants for organic-nitrogen inputs to temperate grasslands. *Ecology* 84:1277–1287
- Brookes PC, Landman A, Pruden G, Jenkinson D (1985) Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biol Biochem* 17:837–842
- Carter MR (2008) Soil sampling and methods of analysis. CRC Press, Boca Raton
- Christou M, Avramides EJ, Roberts JP, Jones DL (2005) Dissolved organic nitrogen in contrasting agricultural ecosystems. *Soil Biol Biochem* 37:1560–1563
- de Vries FT, Bloem J, Quirk H, Stevens CJ, Bol R, Bardgett RD (2012) Extensive management promotes plant and microbial nitrogen retention in temperate grassland. *PLoS One* 7
- Elberse WT, Berendse F (1993) A comparative study of the growth and morphology of eight grass species from habitats with different nutrient availabilities. *Funct Ecol* 7:223–229
- Falkengren-Grerup U, Mansson KF, Olsson MO (2000) Uptake capacity of amino acids by ten grasses and forbs in relation to soil acidity and nitrogen availability. *Environ Exp Bot* 44:207–219

- Farrell M, Hill PW, Farrar J, Bardgett RD, Jones DL (2011a) Seasonal variation in soluble soil carbon and nitrogen across a grassland productivity gradient. *Soil Biol Biochem* 43:835–844
- Farrell M, Hill PW, Wanniarachchi SD, Farrar J, Bardgett RD, Jones DL (2011b) Rapid peptide metabolism: A major component of soil nitrogen cycling? *Glob Biogeochem Cycles* 25
- Farrell M, Hill PW, Farrar J, DeLuca TH, Roberts P, Kielland K, Dahlgren R, Murphy DV, Hobbs PJ, Bardgett RD, Jones DL (2013) Oligopeptides represent a preferred source of organic N uptake: a global phenomenon? *Ecosystems* 16:133–145
- Grime JP, Hodgson JG, Hunt R (2007) Comparative plant ecology a functional approach to common British species. Castlepoint Press, Dalbeattie
- Harrison KA, Bol R, Bardgett RD (2007) Preferences for different nitrogen forms by coexisting plant species and soil microbes. *Ecology* 88:989–999
- Harrison KA, Bol R, Bardgett RD (2008) Do plant species with different growth strategies vary in their ability to compete with soil microbes for chemical forms of nitrogen? *Soil Biol Biochem* 40:228–237
- Hill PW, Quilliam RS, DeLuca TH, Farrar J, Farrell M, Roberts P, Newsham KK, Hopkins DW, Bardgett RD, Jones DL (2011) Acquisition and assimilation of nitrogen as peptide-bound and D-enantiomers of amino acids by wheat. *PLoS One* 6
- Hill PW, Marsden KA, Jones DL (2013) How significant to plant N nutrition is the direct consumption of soil microbes by roots? *New Phytol* 199:948–955
- IPCC (2013) Climate change 2013: the physical science basis. contribution of Working Group I to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge
- Jones DL, Healey JR, Willett VB, Farrar JF, Hodge A (2005) Dissolved organic nitrogen uptake by plants - an important N uptake pathway? *Soil Biol Biochem* 37:413–423
- Kahmen A, Renker C, Unsicker SB, Buchmann N (2006) Niche complementarity for nitrogen: An explanation for the biodiversity and ecosystem functioning relationship? *Ecology* 87: 1244–1255
- Klanderud K, Birks HJB (2003) Recent increases in species richness and shifts in altitudinal distributions of Norwegian mountain plants. *The Holocene* 13:1–6
- Lu M, Zhou X, Yang Q, Li H, Luo Y, Fang C, Chen J, Yang X, Li B (2013) Responses of ecosystem carbon cycle to experimental warming: a meta-analysis. *Ecology* 94:726–738
- Luo Y, Wan S, Hui D, Wallace LL (2001) Acclimatization of soil respiration to warming in a tall grass prairie. *Nature* 413:622–625
- McKane RB, Grigal DF, Russelle MP (1990) Spatiotemporal differences in N-15 uptake and the organization of an old-field plant community. *Ecology* 71:1126–1132
- McKane RB, Johnson LC, Shaver GR, Nadelhoffer KJ, Rastetter EB, Fry B, Giblin AE, Kielland K, Kwiatkowski BL, Laundre JA, Murray G (2002) Resource-based niches provide a basis for plant species diversity and dominance in arctic tundra. *Nature* 415:68–71
- Medina-Roldan E, Paz-Ferreiro J, Bardgett RD (2012) Grazing exclusion affects soil and plant communities, but has no impact on soil carbon storage in an upland grassland. *Agric Ecosyst Environ* 149:118–123
- Mommer L, Visser EJW, van Ruijven J, de Caluwe H, Pierik R, de Kroon H (2011) Contrasting root behaviour in two grass species: a test of functionality in dynamic heterogeneous conditions. *Plant Soil* 344:347–360
- Näsholm T, Persson J (2001) Plant acquisition of organic nitrogen in boreal forests. *Physiol Plant* 111:419–426
- Näsholm T, Ekblad A, Nordin A, Giesler R, Högborg M, Högborg P (1998) Boreal forest plants take up organic nitrogen. *Nature* 392:914–916
- Näsholm T, Kielland K, Ganeteg U (2009) Uptake of organic nitrogen by plants. *New Phytol* 182:31–48
- Parolo G, Rossi G (2008) Upward migration of vascular plants following a climate warming trend in the Alps. *Basic Appl Ecol* 9:100–107
- Pomon A, Escaravage N, Lamaze T (2007) Complementarity in mineral nitrogen use among dominant plant species in a subalpine community. *Am J Bot* 94:1778–1785
- Prescott CE (2010) Litter decomposition: what controls it and how can we alter it to sequester more carbon in forest soils? *Biogeochemistry* 101:133–149
- Rennenberg H, Dannenmann M, Gessler A, Kreuzwieser J, Simon J, Papen H (2009) Nitrogen balance in forest soils: nutritional limitation of plants under climate change stresses. *Plant Biol* 11:4–23
- Rodwell JS (1992) British plant communities. Volume 3. Grassland and montane communities. Cambridge University Press, Cambridge
- Ryser R, Wahl S (2001) Interspecific variation in RGR and the underlying traits among 24 grass species grown in full daylight. *Plant Biol* 3:426–436
- Sauheitl L, Glaser B, Weigelt A (2009a) Advantages of compound-specific stable isotope measurements over bulk measurements in studies on plant uptake of intact amino acids. *Rapid Commun Mass Spectrom* 23:3333–3342
- Sauheitl L, Glaser B, Weigelt A (2009b) Uptake of intact amino acids by plants depends on soil amino acid concentrations. *Environ Exp Bot* 66:145–152
- Saxe H, Cannell MGR, Johnsen B, Ryan MG, Vourlitis G (2001) Tree and forest functioning in response to global warming. *New Phytol* 149:369–399
- Schimel JP, Bennett J (2004) Nitrogen mineralization: challenges of a changing paradigm. *Ecology* 85:591–602
- Schippers P, Olff H (2000) Biomass partitioning, architecture and turnover of six herbaceous species from habitats with different nutrient supply. *Plant Ecol* 149:219–231
- Schippers P, Snoeiijing I, Kropff MJ (1999) Competition under high and low nutrient levels among three grassland species occupying different positions in a successional sequence. *New Phytol* 143:547–559
- Soper FM, Paungfoo-Lonhienne C, Brackin R, Rentsch D, Schmidt S, Robinson N (2011) *Arabidopsis* and *Lobelia anceps* access small peptides as a nitrogen source for growth. *Funct Plant Biol* 38:788–796
- Sorensen PL, Michelsen A, Jonasson S (2008) Nitrogen uptake during one year in subarctic pant functional groups and in microbes after long-term warming and fertilization. *Ecosystems* 11:1223–1233
- Vitousek PM, Howarth RW (1991) Nitrogen limitation on land and in the sea: how can it occur? *Biogeochemistry* 13:87–115

- Warren CR (2009) Why does temperature affect relative uptake rates of nitrate, ammonium and glycine: A test with *Eucalyptus pauciflora*. *Soil Biol Biochem* 41:778–784
- Warren CR (2012) Post-uptake metabolism affects quantification of amino acid uptake. *New Phytol* 193:522–531
- Weigelt A, Bol R, Bardgett RD (2005) Preferential uptake of soil nitrogen forms by grassland plant species. *Oecologia* 142: 627–635
- Wilkinson A, Hill PW, Vaieretti MV, Farrar JF, Jones DL, Bardgett RD (2015) Challenging the paradigm of nitrogen cycling: no evidence of in situ resource partitioning by coexisting plant species in grasslands of contrasting fertility. *Ecol Evol* 5:275–287
- Zhang DQ, Hui DF, Luo YQ, Zhou GY (2008) Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *J Plant Ecol* 1:85–93