

# Simple synthesis of $^{32}\text{P}$ -labelled inositol hexakisphosphates for study of phosphate transformations

Hayley Whitfield · Andrew M. Riley ·  
Soulla Diogenous · Himali Y. Godage ·  
Barry V. L. Potter · Charles A. Brearley

Received: 23 March 2017 / Accepted: 12 June 2017 / Published online: 27 June 2017

© The Author(s) 2017. This article is an open access publication

## Abstract

**Background and aims** In many soils inositol hexakisphosphate in its various forms is as abundant as inorganic phosphate. The organismal and geochemical processes that exchange phosphate between inositol hexakisphosphate and other pools of soil phosphate are poorly defined, as are the organisms and enzymes involved. We rationalized that simple enzymic synthesis of inositol hexakisphosphate labeled with  $^{32}\text{P}$  would greatly enable study of transformation of soil inositol phosphates when combined with robust HPLC separations of different inositol phosphates.

---

Responsible Editor: Daniel Menezes-Blackburn.

**Electronic supplementary material** The online version of this article (doi:10.1007/s11104-017-3315-9) contains supplementary material, which is available to authorized users.

---

H. Whitfield · C. A. Brearley (✉)  
School of Biological Sciences, University of Norwich, Norwich  
Research Park, Norwich NR4 7TJ, UK  
e-mail: c.brearley@uea.ac.uk

A. M. Riley · B. V. L. Potter  
Medicinal Chemistry & Drug Discovery, Department of  
Pharmacology, University of Oxford, Mansfield Rd, Oxford OX1  
3QT, UK

S. Diogenous · H. Y. Godage · B. V. L. Potter  
Wolfson Laboratory of Medicinal Chemistry, Department of  
Pharmacy and Pharmacology, University of Bath, Bath BA2 7AY,  
UK

**Methods** We employed the enzyme inositol pentakisphosphate 2-kinase, IP5 2-K, to transfer phosphate from  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  to axial hydroxyl(s) of *myo*-, *neo*- and 1*D-chiro*-inositol phosphate substrates.

**Results**  $^{32}\text{P}$ -labeled inositol phosphates were separated by anion exchange HPLC with phosphate eluents. Additional HPLC methods were developed to allow facile separation of *myo*-, *neo*-, 1*D-chiro*- and *scyllo*-inositol hexakisphosphate on acid gradients.

**Conclusions** We developed enzymic approaches that allow the synthesis of labeled *myo*-inositol 1, [ $^{32}\text{P}$ ]2,3,4,5,6-hexakisphosphate; *neo*-inositol 1, [ $^{32}\text{P}$ ]2,3,4, [ $^{32}\text{P}$ ]5,6-hexakisphosphate and 1*D-chiro*-inositol [ $^{32}\text{P}$ ]1,2,3,4,5, [ $^{32}\text{P}$ ]6-hexakisphosphate. Additionally, we describe HPLC separations of all inositol hexakisphosphates yet identified in soils, using a collection of soil inositol phosphates described in the seminal historic studies of Cosgrove, Tate and coworkers. Our study will enable others to perform radiotracer experiments to analyze fluxes of phosphate to/from inositol hexakisphosphates in different soils.

**Keywords** Phytate · *neo*- · 1*D-chiro*- · *myo*- · *scyllo*- inositol hexakisphosphate · Phosphate

## Abbreviations

HEPES 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid  
HPLC High pressure liquid chromatography  
i.d. Internal diameter

## Introduction

In consideration of the different forms of inositol hexakisphosphate identified in soils: 1D-*chiro*-, *myo*-, *neo*- and *scyllo*- ((Anderson 1964; Anderson and Malcolm 1974; Baker 1974, cited in Turner et al. 2002; Cosgrove 1962, 1963, 1966, 1969a, b; Cosgrove and Tate 1963; Halstead and Anderson 1970; L'Annunziata 1975; L'Annunziata and Fuller 1971; L'Annunziata et al. 1972; reviewed, Cosgrove 1980); Irving and Cosgrove 1982), it remains unclear what the biotic or abiotic origins of D-*chiro*-, *neo*- and *scyllo*-inositol phosphates are (L'Annunziata 2007; Turner and Richardson 2004; Turner et al. 2002). A limited number of studies have shown biotic contribution to the epimerization of unsubstituted inositols (Cosgrove 1969b; L'Annunziata 1975; L'Annunziata and Gonzalez 1977 (reviewed L'Annunziata 2007)), or have shown that chemical, and hence, geochemical, epimerization of *myo*-inositol pentakisphosphate is possible (Cosgrove 1972). It is clear from the foregoing that studies of soil phosphate transformations, particularly those arising from input of *myo*-inositol hexakisphosphate from plant sources, would be greatly enabled by the provision of  $^{32}\text{P}$  or  $^{33}\text{P}$ -labelled *myo*-inositol hexakisphosphate, and, indeed, of other inositol hexakisphosphates. Such materials would, with established extraction and separation techniques, allow facile determination of the exchange of phosphate between organic and inorganic pools and, with appropriate separation techniques, would allow study of transformations of different isomers of inositol hexakisphosphate. With labelled inositol hexakisphosphates and an increasing literature on the 'pathways' of *myo*-inositol hexakisphosphate degradation by phytases of different classes; cysteine phytase, histidine acid phytase, purple-acid phytase,  $\beta$ -propeller phytase (Konietzny and Greiner 2002), it would be possible to begin to describe 'pathways' of inositol hexakisphosphate turnover in soils and the contribution of different organisms to that turnover.

With these thoughts in mind, we have sought to synthesize  $^{32}\text{P}$ -labelled inositol hexakisphosphates by enzymic means. We have taken opportunity of a recombinant inositol pentakisphosphate 2-kinase (IP5 2-K, also known as IPK1) characterized (Banos-Sanz et al. 2012; Gonzalez et al. 2010; Gosein and Miller 2013; Sweetman et al. 2006). This enzyme transfers the

gamma-phosphate) from ATP to the sole axial hydroxyl on carbon 2 of *myo*-inositol phosphates, and is believed to be the enzyme responsible for synthesis of *myo*-inositol hexakisphosphate in all kingdoms that make this molecule. We reasoned that the enzyme might be capable of transferring phosphate from ATP to the axial hydroxyl(s) of other inositols bearing equatorial phosphates.

The structures of the different inositol 'parents' of the inositol phosphates used in this study are shown in Fig. 1. The rules for numbering of carbon atoms, and hence of inositol phosphates bearing phosphate substituents on particular carbons are given in the IUPAC-IUB rules (IUPAC-IUB 1973, 1977). For *myo*-inositol phosphates, only, a relaxation of the rules (NC-IUB 1989) allows numbering of carbons by the D- (1D) or L- (1L) nomenclature, e.g. to assist in delineating metabolic sequences. The 1D- and 1L- numbering of *myo*-inositol is shown in Fig. 1. Supplemental Fig. 1 shows symmetry aspects of the 'parent' inositols of the *myo*-, *neo*-, 1D-*chiro*- and *scyllo*-inositol phosphates discussed hereafter, while Supplemental Fig. 2 shows the structures of the substrates and products obtained therefrom.


For the assistance of researchers wishing to adopt these methods, we provide (Table 1) a summary of the methodologies used with an indication of their applicability and a reference to prior use of the method. We also list the figures in this manuscript in which the method is applied.

## Methods

### Inositol phosphate substrates

The synthesis of *neo*-inositol 1,3,4,6-tetrakisphosphate (*neo*-Ins(1,3,4,6)P<sub>4</sub>), 1D-*chiro*-inositol 1,3,4,6-tetrakisphosphate (1D-*chiro*-Ins(1,3,4,6)P<sub>4</sub>), 1D-*chiro*-inositol 2,3,4,5-tetrakisphosphate (1D-*chiro*-Ins(2,3,4,5)P<sub>4</sub>), *myo*-inositol 1,3,4,5,6-pentakisphosphate (*myo*-Ins(1,3,4,5,6)P<sub>5</sub>), *scyllo*-inositol pentakisphosphate and of *neo*- and 1D-*chiro* inositol hexakisphosphates was described (Diogenous 1999; Liu et al. 1999; Liu et al. 2001; Godage et al. 2013; Riley et al. 2006; Turner et al. 2012)). *Myo*-inositol hexakisphosphate was obtained from Merck Millipore (Product No. 407125).

**Fig. 1** The structures of *myo*-, *neo*-, *scyllo*- and 1*D*-*chiro*-inositols. The projections shown left to right are a Mills projection, a Haworth projection and a three-dimensional structure. Carbon atoms in the ring are numbered according to the IUPAC/IUPAC-IUB recommendations (1973, 1977). For *myo*-inositol, both 1*L*- (upper) and 1*D*- (lower) notation is shown



## Radioisotopes

ATP, [ $\gamma$ - $^{32}$ P]- 3000 Ci mmol $^{-1}$  was obtained from PerkinElmer.

## Cloning and expression of AtIPK1

*Arabidopsis thaliana* IPK1 (AGI number): At5G42810, was cloned from *Arabidopsis thaliana* Col-0 cDNA with forward primer AAGTTCGTTTTTCAGGGCCCG ATGGAGATGATTTTGGAGGAGAA and reverse primer ATGGTCTAGAAA GCTTTAGCTG TGGGAAGTTTTG (vector specific sequence in italics) using Phusion High Fidelity Polymerase (Thermo Scientific). Purified product (Wizard SV Gel and PCR Cleanup System, Promega) was inserted into pOPINF linearized with HindIII and KpnI (Berrow et al. 2007) by ligation independent cloning using In Fusion HD enzyme kit (Clontech). The vector adds a *N*-terminal hexahistidine tag to the recombinant protein. Recombinant plasmid was transformed into *E coli* Rosetta (DE3) (Novagen) and protein production induced from an overnight culture grown in LB containing 0.5%

(w/v) glucose and ampicillin by transfer to LB containing 0.5 mM IPTG and ampicillin with further growth for 7 h at 25 °C.

## Protein purification

AtIPK1 was purified according to (Banos-Sanz et al. 2012).

## Enzyme assays

For assays without radiolabel, inositol phosphate-dependent conversion of ATP to ADP was followed by reverse-phase ion pair HPLC and subsequent detection of nucleotides at 260 nm (Caddick et al. 2008). Briefly, 1  $\mu$ g enzyme was incubated with 500  $\mu$ M inositol phosphate and 50  $\mu$ M ATP in 20 mM HEPES, pH 7.3, 1 mM MgCl $_2$  for 2 h at 25 °C, the volume of the assay was 20  $\mu$ L. Reactions were terminated by the addition of 1  $\mu$ L of conc. HCl, followed after 5 min on ice by the addition of 50  $\mu$ L water. Aliquots (50  $\mu$ L) were analysed by HPLC.

**Table 1** Summary of methods of this study and their application to the study of inositol phosphate metabolism

Subject	Method	Comment	Description/Use	Reference
Stereochemistry of inositol phosphates		A comprehensive review of inositol (phosphate) chemistry and stereochemistry	Fig. 1	Thomas et al. 2016
Source of inositol phosphates	Complete synthesis or extracted from soils	The soil extracted inositol phosphates were the kind gift of Alan Richardson (CSIRO) from the personal stocks of the late Dennis Cosgrove, or were provided by Max Tate (The University of Adelaide)	Figs. 1, 2, 3, 4, 5, and 6; Suppl. Fig. 1	Diogenou 1999; Liu et al. 1999, 2001; Riley et al. 2006; Turner et al. 2012; Godage et al. 2013; Cosgrove, Tate and others (reviewed this manuscript)
Source of synthetic enzyme			Figs. 2, 3, 4, 5, and 6; Suppl. Fig. 1	Sweetman et al. 2006
Non-radioactive verification of enzyme specificity	Reverse-phase ion-pair chromatography with UV detection of nucleotides	Used to titrate enzyme concentration before use of radiolabel	Fig. 2	Caddick et al. 2008
Production of $^{32}\text{P}$ -labeled inositol phosphates	Anion-exchange chromatography on Partisphere SAX columns with on-line detection by Cerenkov counting	Used to separate/verify reaction products	Fig. 3	Breartley and Hanke 1996a, b; Nagy et al. 2009; Stephens 1990
Substrate specificity/product profiles of different phytases	Anion-exchange chromatography on Partisphere SAX columns with on-line detection by Cerenkov counting	Can be used to separate/purify particular isomers	Fig. 4	Stentz et al. 2014
Non radioactive detection of inositol phosphates on acid gradients with UV detection	Anion exchange chromatography on CarboPac PA200 columns with post-column addition of ferric nitrate	A particularly robust separation method for higher inositol phosphates using volatile (HCl) or non-volatile (methanesulfonic) acid eluents. Could be combined with Cerenkov counting for radiolabeled inositol phosphates	Fig. 5	Phillippy and Bland 1988; Blaabjerg et al. 2010
Non radioactive detection of pentakisphosphates that can be generated from soil inositol hexakisphosphates	Anion exchange chromatography on CarboPac PA200 columns with post-column addition of ferric nitrate	A particularly robust separation method for higher inositol phosphates using volatile (HCl) or non-volatile (methanesulfonic) acid eluents. Could be combined with Cerenkov counting for radiolabeled inositol phosphates	Fig. 6	This manuscript
The structures of substrates and products of <i>AtIP5</i> 2 K			Suppl. Fig. 1	This manuscript

For assays with radiolabel, ATP-dependent conversion of inositol phosphates to higher (more phosphorylated) species was followed by anion-exchange HPLC with on-line detection of  $^{32}\text{P}$ . Briefly, 1  $\mu\text{g}$  enzyme was incubated with 500  $\mu\text{M}$  inositol phosphate and 5  $\mu\text{M}$  ATP in the presence of 0.37 MBq  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  in 20 mM HEPES, pH 7.3, 1 mM  $\text{MgCl}_2$  for 1 h at 25  $^\circ\text{C}$ , the volume of the assay was 20  $\mu\text{L}$ . The reaction products were diluted with water and approximately 1–5% of the products were analysed by HPLC.

#### HPLC separation of inositol phosphates

Radiolabeled inositol phosphates were separated by anion exchange HPLC on Partisphere SAX columns eluted at a flow rate of 1  $\text{mL min}^{-1}$  with a gradient derived by mixing solvent from reservoirs containing (A) water and (B) 1.25 M  $(\text{NH}_4)_2\text{HPO}_4$ , adjusted to pH 3.8 with  $\text{H}_3\text{PO}_4$ , according to the following schedule: time (min), % B; 0, 0; 5, 0; 65, 100. Radioactivity was detected by Cerenkov counting in a Radiomatic A500 Series Flo Detector (Canberra Packard, Pangbourne, Bucks, UK) fitted with a 0.5 ml flow cell using an integration interval of 12 s (Hanke et al. 2012).

Non-labelled inositol phosphates were resolved by anion exchange HPLC on a 250  $\times$  3 mm i.d. CarboPac PA200 column (Dionex UK, Ltd) and guard column 50  $\times$  3 mm of the same material, eluted at a flow rate of 0.4  $\text{mL min}^{-1}$  with gradients of either HCl or methanesulfonic acid (Blaabjerg et al. 2010). Inositol phosphates were detected after post-column addition of 0.1% (w/v) ferric nitrate in 2%  $\text{HClO}_4$  (Phillippy and Bland 1988) delivered at a flow rate of 0.2  $\text{mL min}^{-1}$ . The gradient for both eluents was (A) water, (B) 0.6 M acid: time (min), % B; 0, 0; 25, 100; 38, 100.

#### HPLC separation of nucleotides

Aliquots of the products of enzyme assays were analysed according to (Caddick et al. 2008).

#### Nomenclature

For the purpose of this article, the term ‘Ins’ with prefix 1D-*chiro*-, *myo*-, *neo*- or *scyllo*- is used as an abbreviation of the described inositol phosphate. Hence, *neo*-inositol 1,3,4,6-tetrakisphosphate is abbreviated *neo*-Ins(1,3,4,6) $P_4$ . It should be noted, however, that the numbering of phosphate substituents (of the carbon


atoms to which they are attached) is not necessarily the same for different stereoisomers of inositol. The reader is referred to Shears and Turner (2007) for a concise description of terminology and to (Thomas et al. 2016) for a comprehensive review of inositol and inositol phosphate nomenclature and terminology.

## Results

*AtIP5* 2-K can be used to synthesize a range of inositol phosphate epimers, verifiable by non-radioactive assay of inositol phosphate production

We have previously described the use of *AtIP5* 2-K to synthesize *myo*-Ins(1, [ $^{32}\text{P}$ ]2, 3, 4, 5, 6) $P_6$  from  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$  and *myo*-Ins(1,3,4,5,6) $P_5$  (Nagy et al. 2009). Here, we have further examined the ability of *AtIP5*-2 K to phosphorylate *myo*-Ins(1,3,4,6) $P_4$ . We did so, not only because *myo*-Ins(1,3,4,6) $P_4$  possesses an axial 2-OH, but also because of the availability of the *neo*-inositol epimer, *neo*-Ins(1,3,4,6) $P_4$  (Diogenous 1999). This inositol phosphate shares the plane of symmetry that bisects the *myo*-Ins(1,3,4,6) $P_4$  molecule between C2 and C5, but also possesses a second axial hydroxyl on C5 which creates a  $C_2$ -axis of rotational symmetry that bisects the C1-C6 bond and the C3-C4 bond (Supplementary Figure 1). The consequence of this is that single phosphorylation of C2 generates the same product as phosphorylation of C5, while in contrast phosphorylation of C2 of *myo*-Ins(1,3,4,6) $P_4$  is not equivalent to phosphorylation of C5.

We incubated *AtIP5* 2-K with 500  $\mu\text{M}$  inositol phosphate and 50  $\mu\text{M}$  ATP. The products were resolved by reverse-phase ion-pair HPLC with detection of nucleotides at 260 nm (Fig. 2). Peak areas were integrated and the % of the nucleotide converted to ADP was calculated. A control incubation without inositol phosphate confirmed that *AtIP5* 2-K is not a phosphatase; the 1.3% of nucleotide recovered as ADP is typical of the level of contamination of commercial ATP with ADP (Fig. 2a). Inclusion of *myo*-Ins(1,3,4,6) $P_4$  increased ADP production, 12.1% of total nucleotide was recovered as ADP, without production of AMP (Fig. 2b), whereas for the physiological substrate *myo*-Ins(1,3,4,5,6) $P_5$ , included at 50  $\mu\text{M}$ , 26% of nucleotide was recovered as ADP (Fig. 2c). Clearly, *myo*-Ins(1,3,4,6) $P_4$  is a substrate, albeit a poorer one than *myo*-Ins(1,3,4,5,6) $P_5$ . In contrast, *neo*-Ins(1,3,4,6) $P_4$



**Fig. 2** Phosphotransferase reactions catalysed by AtIP5 2-K. The nucleotide products of reactions of enzyme, ATP and different inositol phosphates were separated by ion-pair RP-HPLC and detected by absorbance at 260 nm. **a** no inositol phosphate; **b** *myo*-Ins(1,3,4,6) $P_4$ ; **c** *myo*-Ins(1,3,4,5,6) $P_5$ ; **d** *neo*-Ins(1,3,4,6) $P_4$ ; **e** 1D-*chiro*-Ins(2,3,4,5) $P_4$ ; **f** 1D-*chiro*-Ins(1,3,4,6) $P_4$

was a strong substrate with ADP production at 90.8% (Fig. 2d).

Similarly, inclusion of 1D-*chiro*-Ins(2,3,4,5) $P_4$  at 500  $\mu$ M resulted in 92% conversion of ATP to ADP (Fig. 2e), while 1D-*chiro*-Ins(1,3,4,6) $P_4$  whose two hydroxyls, on C2 and C5 are equatorial (Supplementary

Figure 1) was not a substrate for AtIP5 2-K, with ADP representing 1.5% of the total nucleotide (Fig. 2f).


These results confirm the utility of *neo*-Ins(1,3,4,6) $P_4$  and 1D-*chiro*-Ins(2,3,4,5) $P_4$  as substrates from which higher *neo*- and 1D-*chiro*-inositol phosphates can be synthesized with AtIP5 2-K. They further confirm the exclusive phosphorylation of axial hydroxyls by this enzyme (Gonzalez et al. 2010; Sweetman et al. 2006).

We additionally tested racemic mixtures of 1D/L-*neo*-Ins(1,2,4) $P_3$  and 1D/L-*neo*-Ins(1,3,4) $P_3$  as substrates in extended (16 h) incubations at 500  $\mu$ M concentration with 50  $\mu$ M ATP. We did not observe production of ADP, thus these molecules are not substrates despite possessing one and two axial hydroxyls at C5, and C2 and C5 respectively (data not shown).

AtIP5 2-K can be used to synthesize a range of  $^{32}$ P-labelled inositol phosphates

Having established that *neo*-Ins(1,3,4,6) $P_4$  and 1D-*chiro*-Ins(2,3,4,5) $P_4$  are novel substrates of AtIP5 2 K, we performed enzyme assays to produce  $^{32}$ P-labelled inositol phosphates. The substrate concentrations used were 500  $\mu$ M inositol phosphate and 5  $\mu$ M ATP. Reactions were terminated and the products spiked with additional ATP to allow online tandem UV-radioactivity monitoring of the chromatography (Fig. 3). We included *myo*-Ins(1,3,4,5,6) $P_5$  as the canonical substrate and observed (Fig. 3a) in addition to a major peak of *myo*-Ins $P_6$  eluting at a retention time of 55 min, a small peak of unidentified material eluting at 38.8 min, a major peak of unreacted ATP at 25.2 min and a peak of inorganic phosphate at 14.6 min. We note that others have reported impurities in commercial [ $^{32}$ P]ATP that elute on Partisphere SAX columns with similar chromatographic mobility to *myo*-Ins $P_4$ s (Stephens 1990).

Provision of *neo*-Ins(1,3,4,6) $P_4$  as substrate yielded (Fig. 3b) a major peak of label with the chromatographic property of Ins $P_5$  at 46.4 min and a more polar (highly charged peak) with the chromatographic property of an Ins $P_6$ . This peak eluted at 56.4 min, slightly after *myo*-Ins $P_6$ . The trace also showed peaks of assumed impurity, ATP and inorganic phosphate. Considering the structure of *neo*-Ins(1,3,4,6) $P_4$  and its  $C_2$  axis of rotational symmetry (Supplemental Figs. 1, 2), the two axial hydroxyls are superposable so there is only one possible Ins $P_5$  product. IUPAC conventions recommend the naming of substituents by the lowest numbered locants,



**Fig. 3** Synthesis of  $^{32}\text{P}$ -labelled inositol phosphates with AtIP5 2-K. The inositol phosphate products of reactions of enzyme,  $^{32}\text{P}$ ATP and different inositol phosphates were separated by Partisphere Strong Anion Exchange HPLC and radioactivity estimated by on-line Cerenkov counting. **a** *myo*-Ins(1,3,4,5,6) $P_5$ ; **b** *neo*-Ins(1,3,4,6) $P_4$ ; **c** 1D-*chiro*-Ins(1,3,4,6) $P_4$ , **d** 1D-*chiro*-Ins(2,3,4,5) $P_4$ . The peaks labelled are the  $^{32}\text{P}$  labelled products or reactants (ATP)


hence single phosphorylation of one axial hydroxyl of *neo*-Ins(1,3,4,6) $P_4$  yields *neo*-Ins(1,2,3,4,6) $P_5$  = *neo*-

Ins(1,3,4,5,6) $P_5$  with preferred use of the former name. The addition of  $^{32}\text{P}$  to an unlabeled substrate therefore yields *neo*-Ins(1,[ $^{32}\text{P}$ ]2,3,4,6) $P_5$ . The elution of a second more polar peak, we assume to represent the double phosphorylation of *neo*-Ins(1,3,4,6) $P_4$  to give *neo*-Ins(1,[ $^{32}\text{P}$ ]2,3,4,[ $^{32}\text{P}$ ]5,6) $P_6$ .

We also tested 1D-*chiro*-Ins(1,3,4,6) $P_4$  and 1D-*chiro*-Ins(2,3,4,5) $P_4$  as co-substrates with  $^{32}\text{P}$ ATP (Fig. 3c, d). Significantly, the former, which bears equatorial hydroxyls on C2 and C5, and lacks axial hydroxyls (Fig. 1; Supplemental Figs. 1, 2), was not a substrate and yielded peaks of unreacted ATP, inorganic phosphate and the assumed contaminant with retention time 39 min (Fig. 3c). In contrast, 1D-*chiro*-Ins(2,3,4,5) $P_4$ , like *neo*-Ins(1,3,4,6) $P_4$ , yielded  $^{32}\text{P}$ -labeled peaks with the chromatographic mobility of *InsP\_5*, retention time 48 min; and *InsP\_6*, retention time 59.5 min (Fig. 3d). Comparison of the retention time of peaks of *InsP\_5* and *InsP\_6* products (Fig. 3a–d) reveals that the *InsP\_5* and *InsP\_6* products of different epimers of inositol (tetrakisphosphate) are chromatographically distinct. Again, 1D-*chiro*-Ins(2,3,4,5) $P_4$ , another substrate with two axial hydroxyls, yielded products which, relative to the substrate, were singly and doubly phosphorylated by AtIP5 2-K. Moreover, 1D-*chiro*-Ins(2,3,4,5) $P_4$  possesses a  $C_2$ -axis of symmetry, here bisecting the C1-C6 and C3-C4 bonds, which superposes C1 and C2 substituents. Consequently, a common *InsP\_5* product is generated from phosphorylation of either axial hydroxyl and the product is 1D-*chiro*-Ins(1,2,3,4,5) $P_5$ , here 1D-*chiro*-Ins( $^{32}\text{P}$ ]1,2,3,4,5) $P_5$ . The *InsP\_6* product is 1D-*chiro*-Ins( $^{32}\text{P}$ ]1,2,3,4,5,[ $^{32}\text{P}$ ]6) $P_6$ .

Different phytases yield characteristic product profiles from  $^{32}\text{P}$ -labeled *myo*-inositol hexakisphosphate

By way of illustration of how different phytases yield different product profiles from the same substrate, we show (Fig. 4) the products of progressive dephosphorylation of *myo*-Ins(1,[ $^{32}\text{P}$ ]2,3,4,5,6) $P_6$  by histidine acid phytases of fungal and bacterial origin. We chose *Aspergillus ficuum* phytase, a D-3 phytase (data of Fig. 4c), where D-3 signifies the position of attack on *myo*-Ins $P_6$  and HD is one of the canonical motifs, D (aspartate) being a proton donor that activates a water molecule that is responsible for the cleavage of scissile phosphate. The other phytase (data of Fig. 4b) is another histidine acid phytase, but is the archetype of a recently




**Fig. 4** Separations of  $^{32}\text{P}$  labelled inositol phosphate products of dephosphorylation of *myo*-Ins(1,[ $^{32}\text{P}$ ]2,3,4,5,6) $P_5$  by HAE (*Bacteroides thetaiotomicron* Minpp) and HD (*Aspergillus niger*) phytases. The products of reactions of **a** no enzyme control; **b** *Bt*Minpp and **c** *A ficuum* phytase were separated by Partisphere Strong Anion Exchange HPLC and radioactivity estimated by on-line Cerenkov counting. The positions of elution of representative classes of *myo*-inositol phosphate products and of contaminating ATP in the *myo*-[ $^{32}\text{P}$ ]Ins $P_6$  preparation are indicated. The difference in retention time of inositol hexakisphosphate between different panels reflects the use of different Partisphere SAX columns for the separations

described HAE subclass where E (glutamate) is the likely proton donor. This enzyme generates three resolvable Ins $P_5$  products from *myo*-Ins $P_6$  (Stentz et al. 2014). These data reveal the utility of inositol hexakisphosphate labeled on the axial 2-position for study of phytases in vitro. Clearly, they indicate the potential utility of such compounds and such chromatography for study of soil processes.

#### Separation of inositol hexakisphosphates identified in soils

Using the post-column complexation method (Phillippy and Bland 1988) and a CarboPac PA-200 column eluted with a gradient of HCl, we were able to resolve in order of increasing retention time, *neo*-Ins $P_6$ , *muco*-Ins $P_6$ , 1*D*-*chiro*-Ins $P_6$ , *myo*-Ins $P_6$  and *scyllo*-Ins $P_6$  (Fig. 5a), all samples obtained from the laboratory of the late Dennis Cosgrove. We assume that the *muco*-Ins $P_6$  was that made by chemical phosphorylation of the inositol (Cosgrove 1975). The gradient shows a strongly sloping baseline arising from the acid eluent and this UV



**Fig. 5** Separations of soil-representative inositol phosphates on a Dionex CarboPac PA200 column. The column was eluted with a gradient of HCl (**a**) or methanesulfonic acid (**b**) and inositol phosphates detected by post-column complexation with ferric nitrate in perchloric acid and subsequent detection at 290 nm. **a**, **b** Inositol phosphates were obtained from the laboratory of the late DJ Cosgrove. **a** the traces from individual injections (approximately 20 nmol) of different inositol hexakisphosphates are offset on the Y-scale to aid visualisation. **b** Samples of individual injections of synthetic *neo*-Ins $P_4$  with retention time 16–17 min, *neo*-Ins $P_5$  with retention time 21–22 min and *neo*-Ins $P_6$  with retention time 26.9 min are overlaid




absorbance can be negated when using methanesulfonic acid as the eluent (Blaabjerg et al. 2010); however, at least up to 0.6 M, methanesulfonic acid was not a strong enough eluent to elute *scyllo*-InsP<sub>6</sub> from this column.

### Separation of *neo*-inositol phosphates

Methanesulfonic acid was, however, suitable for separation of *neo*-InsPs: *neo*-InsP<sub>4s</sub>, *neo*-InsP<sub>5s</sub> and *neo*-InsP<sub>6</sub> (Fig. 5b), again, these compounds were likely produced by chemical phosphorylation of *neo*-inositol with sodium trimetaphosphate (Cosgrove 1969a). Irving (1980) reported that *neo*-InsP<sub>6</sub> was a good substrate of both the *Pseudomonas* (sp. unknown) bacterium (SB<sub>2</sub>) phytase of Cosgrove (Cosgrove et al. 1970) and *A. ficuum* phytase (Irving and Cosgrove 1974). We note that there are nine possible *neo*-InsP<sub>4s</sub>, comprised of three pairs of enantiomers and three *meso*-compounds and there are three possible *neo*-InsP<sub>5s</sub> (Thomas et al. 2016). Of the *neo*-InsP<sub>5s</sub>, two: 1L-*neo*-Ins(1,2,3,4,5)P<sub>5</sub> = 1D-*neo*-Ins(1,2,3,5,6)P<sub>5</sub> and 1D-*neo*-Ins(1,2,3,4,5)P<sub>5</sub> = 1L-*neo*-Ins(1,2,3,5,6)P<sub>5</sub> are a pair of enantiomers, the third is the *meso*-compound *neo*-Ins(1,2,3,4,6)P<sub>5</sub>. In the absence of chiral HPLC methods for separating enantiomers of inositol phosphates, the separation of two peaks of *neo*-InsP<sub>5</sub> from the Cosgrove samples (Fig. 5b) is all that is achievable, but nevertheless could be diagnostic in studies of *neo*-InsP<sub>6</sub> transformation in soils.

### Separation of inositol pentakisphosphates; the initial products of phytase action on inositol hexakisphosphate

Finally, we show (Fig. 6) the separation of a variety of InsP<sub>5s</sub> including the four separable peaks of *myo*-InsP<sub>5</sub>, identified in Fig. 6b by the position of the single hydroxyl; the two enantiomeric pairs 1D-1/3-OH and 1D-4/6-OH, indicated [1/3-OH] and [4/6-OH] on the figure, cannot be separated into individual enantiomers on non-chiral HPLC. Figure 6b also shows the single *scyllo*-InsP<sub>5</sub> and several InsP<sub>5s</sub> present in a 1L-*chiro*-InsP<sub>5</sub> sample. There are three possible 1L-*chiro*-InsP<sub>5s</sub> and, similarly, three possible 1D-*chiro*-InsP<sub>5s</sub>. Minor *chiro*-InsP<sub>5</sub> peaks with common retention times were observed in the 1D-*chiro*-InsP<sub>6</sub> sample (Fig. 6a) and the 1L-*chiro*-InsP<sub>5</sub> sample (Fig. 6b), with two of the three co-eluting precisely with peaks in a 1L-*chiro*-InsP<sub>5</sub> sample (Fig. 6a). Clearly, this column/eluent



**Fig. 6** HPLC separation of soil-representative inositol pentakisphosphates. A Dionex CarboPac PA200 column was eluted with a gradient of methanesulfonic acid. Inositol phosphates were obtained from the laboratory of the late DJ Cosgrove. **a** The traces from individual injections of different inositol phosphate samples: a 1D-*chiro*-InsP<sub>6</sub> sample with three 1D-*chiro*-InsP<sub>5s</sub>, a 1L-*chiro*-InsP<sub>5</sub> sample with two 1L-*chiro*-InsP<sub>5s</sub> and a *neo*-InsP<sub>5</sub> sample. **b** The traces from individual injections of different inositol phosphate samples: a 1L-*chiro*-InsP<sub>6</sub> sample with three 1L-*chiro*-InsP<sub>5s</sub>, a *scyllo*-InsP<sub>5</sub> sample and an acid hydrolysate of *myo*-InsP<sub>6</sub> with all four resolvable *myo*-InsP<sub>5s</sub>. **a, b** Traces are offset on the Y-scale (left or right) to aid visualisation

combination has great resolving power for all the InsP<sub>5s</sub> expected of soil samples.

## Discussion

Inositol phosphates are major components of total soil phosphate and commonly the dominant organic phosphates in these environments (Turner et al. 2002). Despite their abundance, the origins of *neo*-, 1D-*chiro*- and *scyllo*-inositol hexakisphosphates in soils are poorly defined. It is plausible that they derive from the considerable inputs to soil of plant-derived *myo*-inositol

hexakisphosphate, representing up to several percent of seed dry weight (Raboy 2003). It has been estimated that the sequestration of phosphorus in the *myo*-inositol hexakisphosphate, phytate, reserves of cropped organs of plants is equivalent to the per annum application of phosphorus as fertilizer to soils (Lott et al. 2000). It is remarkable therefore that we do not know, how plant-derived phytate is, likely, converted to other forms (epimers) of inositol phosphates, nor whether the processes are organismal or geochemical.

The epimerization of unsubstituted inositols by unsterilized soil is well described (L'Annunziata and Gonzalez 1977) and pathways by which *myo*-inositol and *scyllo*-inositol are converted have been engineered in *Bacillus subtilis* (Kang et al. 2017; Tanaka et al. 2013; Yamaoka et al. 2011). Similarly, *Geobacillus kaustophilus* HTA426, has three dehydrogenases that are capable of acting as epimerases to interconvert *myo*-, *scyllo*-, and 1D-*chiro*- inositols (Yoshida et al. 2012). The American cockroach, *Periplaneta americana*, was reported to possess epimerase activity producing *neo*-inositol. (Hipps et al. 1973). Plants have the ability to epimerize a number of methylated inositols, reviewed (Thomas et al. 2016). These studies seem to suggest that epimerase activity is restricted to inositols lacking phosphate substituents. It is intriguing therefore that *scyllo*-phosphoinositides were detected in barley (Narasimhan et al. 1997) and that highly phosphorylated inositol phosphates and diphosphoinositol phosphates containing *neo*- rather than *myo*-inositol are the major form of inositol phosphate in *Entamoeba histolytica* (Martin et al. 2000), despite earlier suggestion to the contrary (Martin et al. 1993).

Because the axial 2-phosphate is the last phosphate added in the synthesis of *myo*-inositol hexakisphosphate in plants (Brearley and Hanke 1996) and *Dictyostelium discoideum* (Stephens and Irvine 1990) it is possible to use the enzyme catalyzing this step to make *myo*-inositol 1,[<sup>32</sup>P]2,3,4,5,6-hexakisphosphate. Moreover, because most phytases of plant, bacterial or fungal origin do not remove this phosphate until very late in the sequence of *myo*-inositol hexakisphosphate degradation, if at all (Konietzny and Greiner 2002), the <sup>32</sup>P label will be retained in successive *myo*-InsP<sub>5</sub>, InsP<sub>4</sub>, InsP<sub>3</sub> and InsP<sub>2</sub> products of dephosphorylation. Consequently, addition of <sup>32</sup>P-labeled *myo*-InsP<sub>6</sub> to soils will, by simple chromatography using the methods elaborated here, allow researchers to study *myo*-inositol hexakisphosphate turnover in their soil of choice.

From a practical perspective, the use of a high energy β emitter such as <sup>32</sup>P allows facile detection of radioactivity in column eluates by on-line Cerenkov counting, obviating the requirement for the addition of scintillation fluid. The use of flow-detectors therefore allows for simple collection of radiolabelled fractions, albeit in high salt concentrations required to elute highly polar inositol phosphates. For phosphate eluents, it is a simple exercise to desalt the collected fractions on Dowex AG1 X8 resin with volatile ammonium formate / formic acid mixtures, subsequently removed by freeze-drying (Stephens 1990; Brearley et al. 1997), while for HCl eluents the HCl can be removed directly by freeze-drying.

From a diagnostic perspective, because different phytases produce different *myo*-InsP<sub>5</sub> products (after all, this is the explanation of their classification e.g. as D3, *Aspergillus*; D6, *E.coli* or D5, lily pollen alkaline phytases (Konietzny and Greiner 2002)), simple analysis of products at the level of *myo*-InsP<sub>5</sub> will allow identification of the likely class of enzyme predominantly responsible for initial degradation of *myo*-InsP<sub>6</sub> in different soils. Indeed, it will be fascinating to correlate *myo*-inositol hexakisphosphate degradation products with metagenomic characterization of phytase and microorganism abundance in different soils of the sort recently described (Neal et al. 2017).

In consideration of the routes of degradation of other inositol hexakisphosphates, the other labeled isomers that we describe will be of particular value. Perhaps unsurprisingly, we note the seminal work of Cosgrove (1969a, 1970) and Irving and Cosgrove (1971) in characterization of products of dephosphorylation of *myo*-inositol-, *scyllo*-inositol-, and 1D-*chiro*-inositol hexakisphosphate by a bacterial phytase, and similar studies of wheat phytase by Lim and Tate (1971, 1973) following the methods of Tomlinson and Ballou (1962). These works, reviewed (Irving 1980), show that the axial phosphates of *myo*- and 1D-*chiro*-inositol hexakisphosphates are the last to be removed by plant, bacterial and fungal phytases where tested. Similar conclusions can be drawn for fungal phytase action on 1L-*chiro*-inositol hexakisphosphate (Adelt et al. 2003). With these observations in mind, the use of the labeled InsP<sub>5</sub> and InsP<sub>6</sub> species and the powerful separation approaches identified in the foregoing should allow for facile assessment of the exchange of phosphate between different inositol phosphates in soil contexts.

**Acknowledgements** This work was supported with funding from the Biotechnology and Biological Sciences Research Council, UK Grant BB/N002024/1 to CAB and the Wellcome Trust. BVLP is a Wellcome Trust Senior Investigator (Grant 101010). We thank Max Tate (University of Adelaide) for the generous gift of soil-extracted inositol phosphates and thank Alan Richardson (CSIRO) for the gift of the same from the stocks of the late Dennis Cosgrove.

**Author contributions** HW, BVLP, AMR and CAB designed the study. All authors discussed the results, HW and CAB wrote the manuscript with input from other coauthors.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Adelt S, Podeschwa M, Dallmann G, Altenbach H-J, Vogel G (2003) Stereo- and regiospecificity of yeast phytases—chemical synthesis and enzymatic conversion of the substrate analogues *neo*- and *L-chiro*-inositol hexakisphosphate. *Bioorg Chem* 31:44–67
- Anderson G (1964) Investigations on the analysis of inositol hexaphosphate in soil. *Trans Int Congress Soil Sci* 4:563–571
- Anderson G, Malcolm RE (1974) The nature of alkali soluble soil organic phosphates. *J Soil Sci* 25:282–297
- Baker RT (1974) Chemical nature and distribution of soil organic phosphate in two chronosequences of soils. PhD thesis, Department of Soil Science, University of Canterbury, New Zealand
- Banos-Sanz JI, Sanz-Aparicio J, Whitfield H, Hamilton C, Brearley CA, Gonzalez B (2012) Conformational changes in inositol 1,3,4,5,6-pentakisphosphate 2-kinase upon substrate binding: role of *N*-terminal lobe and enantiomeric substrate preference. *J Biol Chem* 287:29237–29249
- Berrow NS, Alderton D, Sainsbury S, Nettleship J, Assenberg R, Rahman N, Stuart DI, Owens RJ (2007) A versatile ligation-independent cloning method suitable for high-throughput expression screening applications. *Nucleic Acids Res* 35:e45
- Blaabjerg K, Hansen-Møller J, Poulsen HD (2010) High-performance ion chromatography method for separation and quantification of inositol phosphates in diets and digesta. *J Chromatogr B Anal Technol Biomed Life Sci* 878:347–354
- Brearley CA, Hanke DE (1996) Metabolic evidence for the order of addition of individual phosphate esters in the *myo*-inositol moiety of inositol hexakisphosphate in the duckweed *Spirodela polyrrhiza* L. *Biochem J* 314:227–233
- Brearley CA, Parmar PN, Hanke DE (1997) Metabolic evidence for PtdIns(4,5)P-2-directed phospholipase C in permeabilized plant protoplasts. *Biochem J* 324:123–131
- Caddick SE, Harrison CJ, Stavridou I, Mitchell JL, Hemmings AM, Brearley CA (2008) A *Solanum tuberosum* inositol phosphate kinase (StITPK1) displaying inositol phosphate-inositol phosphate and inositol phosphate-ADP phosphotransferase activities. *FEBS Lett* 582:1731–1737
- Cosgrove DJ (1962) Forms of inositol hexaphosphate in soils. *Nature* 194:1265–1266
- Cosgrove DJ (1963) The chemical nature of soil organic phosphorus. I. Inositol phosphates. *Aust J Soil Res* 1:203–214
- Cosgrove DJ (1966) Detection of isomers of phytic acid in some Scottish and Californian soils. *Soil Sci* 102:42–43
- Cosgrove DJ (1969a) Ion-exchange chromatography of inositol polyphosphates. *Ann N Y Acad Sci* 165:677–686
- Cosgrove DJ (1969b) The chemical nature of soil organic phosphorus—II: characterization of the supposed *dl-chiro*-inositol hexaphosphate component of soil phytate as *d-chiro*-inositol hexaphosphate. *Soil Biol Biochem* 1:325–327
- Cosgrove DJ (1970) Inositol phosphate phosphatase of microbiological origin. Inositol pentaphosphate intermediates in the dephosphorylation of the hexaphosphates of *myo*-inositol, *scyllo*-inositol, and *D-chiro*-inositol, by a bacterial (*Pseudomonas* sp.) phytase. *Aust J Biol Sci* 23:1207–1220
- Cosgrove DJ (1972) The origin of inositol polyphosphates in soil. Some model experiments in aqueous systems involving the chemical phosphorylation of *myo*-inositol and the epimerization of *myo*-inositol pentaphosphates. *Soil Biol Biochem* 4:387–396
- Cosgrove DK (1975) The phosphorylation of epi-inositol and muco-inositol with polyphosphoric acid. *Carbohydr Chem* 40:380–384
- Cosgrove DJ (1980) Inositol phosphates—their chemistry, biochemistry and physiology. Elsevier Scientific, Amsterdam
- Cosgrove DJ, Tate ME (1963) Occurrence of *neo*-inositol hexaphosphate in soils. *Nature* 200:568–569
- Cosgrove DJ, Irving GC, Bromfield SM (1970) Inositol phosphate phosphatase of microbial origin. The isolation of soil bacteria having inositol phosphate phosphatase activity. *Aust J Biol Sci* 23:339–343
- Diogenous S (1999) Synthesis of novel ligands for intervention in the polyphosphoinositide pathway of cell signalling. Ph.D. Thesis University of Bath
- Godage HY, Riley AM, Woodman TJ, Thomas MP, Mahon MF, Potter BVL (2013) Regioselective opening of *myo*-inositol orthoesters: mechanism and synthetic utility. *J Org Chem* 78:2275–2288
- Gonzalez B, Banos-Sanz JI, Villate M, Brearley CA, Sanz-Aparicio J (2010) Inositol 1,3,4,5,6-pentakisphosphate 2-kinase is a distant IPK member with a singular inositide binding site for axial 2-OH recognition. *Proc Natl Acad Sci U S A* 107:9608–9613
- Gosein V, Miller GJ (2013) Roles of phosphate recognition in inositol 1,3,4,5,6-pentakisphosphate 2-kinase (IPK1) substrate binding and activation. *J Biol Chem* 288:26908–26913
- Halstead RL, Anderson G (1970) Chromatographic fractionation of organic phosphates from alkali, acid, and aqueous acetylacetone extracts of soils. *Can J Soil Sci* 50:111–119
- Hanke DE, Parmar PN, Caddick SE, Green P, Brearley CA (2012) Synthesis of inositol phosphate ligands of plant hormone-receptor complexes: pathways of inositol hexakisphosphate turnover. *Biochem J* 444:601–609

- Hipps PP, Sehgal RK, Holland WH, Sherman WR (1973) Identification and partial characterization of inositol: NAD<sup>+</sup> epimerase and inosose: NAD(P)H reductase from the fat body of the American cockroach, *Periplaneta americana* L. *Biochemistry* 12:4507–4512
- Irving G CJ (1980) Intermediates in the dephosphorylation of P6-inositols by phytase enzymes. In: Cosgrove DJ (ed) *Inositol phosphates: their chemistry biochemistry and physiology*. Studies in organic chemistry 4. Elsevier Scientific Publishing Company, Amsterdam, pp 99–117
- Irving G CJ, Cosgrove DJ (1971) Inositol phosphate phosphatases of microbiological origin. Some properties of a partially purified bacterial (*Pseudomonas* sp.) phytase. *Aust J Biol Sci* 24:547–557
- Irving G CJ, Cosgrove DJ (1974) Inositol phosphate phosphatases of microbiological origin. Some properties of the partially purified phosphatases of *Aspergillus ficuum* NRRL 3135. *Aust J Biol Sci* 27:361–368
- Irving G CJ, Cosgrove DJ (1982) The use of gas-liquid chromatography to determine the proportions of inositol isomers present as pentakis- and hexakisphosphates in alkaline extracts of soils. *Commun Soil Sci Plant Anal* 13: 957–967
- IUB Nomenclature Committee of the International Union of Biochemistry (1989) Numbering of atoms in *myo*-inositol. *Biochem J* 258:1–2
- IUPAC–IUB Commission on Biochemical Nomenclature (CBN) (1973) Nomenclature of cyclitols. Recommendations 1973. *Biochem J* 153:23–31
- IUPAC–IUB Commission on Biochemical Nomenclature (CBN) (1977) Nomenclature of phosphorus-containing compounds of biochemical importance. Recommendations 1976. *Proc Natl Acad Sci U S A* 74:2222–2230
- Kang DM, Tanaka K, Takenaka S, Ishikawa S, Yoshida KI (2017) *Bacillus subtilis* ioU encodes an additional NADP<sup>+</sup>-dependent *scyllo*-inositol dehydrogenase. *Biosci Biotechnol Biochem* 1–7
- Konietzny U, Greiner R (2002) Molecular and catalytic properties of phytate-degrading enzymes (phytases). *Int J Food Sci Technol* 37:791–812
- L'Annunziata MF (1975) The origin and transformations of the soil inositol phosphate isomers. *Soil Sci Soc Am Proc* 39: 377–379
- L'Annunziata MF, Fuller WH (1971) Soil and plant relationships of inositol phosphate stereoisomers; the identification of *D-chiro*- and *muco*-inositol phosphates in a desert soil and plant system. *Soil Sci Soc Am Proc* 35(587–595):239–253
- L'Annunziata MF, Gonzalez J (1977) Soil metabolic transformations of carbon-14-*myo*-inositol, carbon-14- phytic acid and carbon-14-iron(III) phytate. In: *Soil organic matter studies*, vol. 1. International Atomic Energy Agency, Publication No. IAEA-SM-211/66, Vienna, Austria, pp
- L'Annunziata MF, Fuller WH, Brantley DS (1972) *D-chiro*-Inositol phosphate in a forest soil. *Soil Sci Soc Am Proc* 36:183–184
- L'Annunziata MF (2007) Seed phosphorus and the development of low-phytate crops. In: Turner BL, Richardson AE, Mullaney EJ (eds) *Inositol phosphates: linking agriculture to the environment*. CAB International, Wallingford, pp 41–59
- Lim PE, Tate ME (1971) The phytases I. Lysolecithin-activated phytase from wheat bran. *Biochim Biophys Acta* 250:155–164
- Lim PE, Tate ME (1973) The phytases II. *Biochim Biophys Acta* 302:316–328
- Liu C, Davis RJ, Nahorski SR, Ballereau S, Spiess B, Potter BVL (1999) Synthesis, calcium mobilizing, and physicochemical properties of *D-chiro*-inositol 1,3,4,6-tetrakisphosphate, a novel and potent ligand at the *D-myo*-inositol 1,4,5-trisphosphate receptor. *J Med Chem* 42:1991–1998
- Liu C, Riley AM, Yang X, Shears SB, Potter BVL (2001) Synthesis and biological activity of *D*- and *L-chiro*-inositol 2,3,4,5-tetrakisphosphate: design of a novel and potent inhibitor of Ins(3,4,5,6)P<sub>4</sub> 1-kinase/Ins(1,3,4)P<sub>3</sub> 5/6-kinase. *J Med Chem* 44:2984–2989
- Lott JNA, Ockenden I, Raboy V, Batten G (2000) Phytic acid and phosphorus in crop seeds and fruits: a global estimate. *Seed Sci Res* 10:11–33
- Martin JB, Bakker-Grunwald T, Klein G (1993) <sup>31</sup>P-NMR analysis of *Entamoeba histolytica*. Occurrence of high amounts of two inositol phosphates. *Eur J Biochem* 214:711–718
- Martin JB, Laussmann T, Bakker-Grunwald T, Vogel G, Klein G (2000) neo-inositol polyphosphates in the amoeba *Entamoeba histolytica*. *J Biol Chem* 275:10134–10140
- Nagy R, Grob H, Weder B, Green P, Klein M, Frelet-Barrand A, Schjoerring JK, Brearley C, Martinoia E (2009) The *Arabidopsis* ATP-binding cassette protein AtMRP5/AtABCC5 is a high affinity inositol hexakisphosphate transporter involved in guard cell signaling and phytate storage. *J Biol Chem* 284:33614–33622
- Narasimhan B, Pliska-Matyshak G, Kinnard R, Carstensen S, Ritter MA, Von Weymar L, Murthy P (1997) Novel phosphoinositides in barley aleurone cells (additional evidence for the presence of phosphatidyl-scylo-inositol). *Plant Physiol* 113:1385–1393
- Neal AL, Rossmann M, Brearley C, Akkari E, Guyomar C, Clark IM, Allen E, Hirsch PR (2017) Land-use influences phosphatase gene microdiversity in soils. *Environ Microbiol*. doi:10.1111/1462-2920.13778
- Phillippy BQ, Bland JM (1988) Gradient ion chromatography of inositol phosphates. *Anal Biochem* 175:162–166
- Raboy V (2003) *myo*-Inositol-1,2,3,4,5,6-hexakisphosphate. *Phytochemistry* 64:1033–1043
- Riley AM, Trusselle M, Kuad P, Borkovec M, Cho J, Choi JH, Qian X, Shears SB, Spiess B, Potter BVL (2006) *scyllo*-inositol pentakisphosphate as an analogue of *myo*-inositol 1,3,4,5,6-pentakisphosphate: chemical synthesis, physicochemistry and biological applications. *Chembiochem* 7:1114–1122
- Shears SB, Turner BL (2007) Nomenclature and terminology of inositol phosphates: clarification and a glossary of terms. In: Turner BL, Richardson AE, Mullaney EJ (eds) *Inositol phosphates: linking agriculture to the environment*. CAB International, Wallingford, pp 1–6
- Stentz R, Osborne S, Horn N, Li AW, Hautefort I, Bongaerts R, Rouyer M, Bailey P, Shears SB, Hemmings AM, Brearley CA, Carding SR (2014) A bacterial homolog of a eukaryotic inositol phosphate signaling enzyme mediates cross-kingdom dialog in the mammalian gut. *Cell Rep* 6:646–656
- Stephens LR (1990) Preparation and separation of inositol tetrakisphosphates and inositol pentakisphosphates and the

- establishment of enantiomeric configurations by the use of L-idoitol dehydrogenase. In: Irvine RF (ed) Methods in inositide research. Raven Press, New York, NY, pp 9–30
- Stephens LR, Irvine RF (1990) Stepwise phosphorylation of myo-inositol leading to myo-inositol hexakisphosphate in *Dictyostelium*. Nature 346:580–583
- Sweetman D, Johnson S, Caddick SE, Hanke DE, Brearley CA (2006) Characterization of an *Arabidopsis* inositol 1,3,4,5,6-pentakisphosphate 2-kinase (AtIPK1). Biochem J 394:95–103
- Tanaka K, Tajima S, Takenaka S, Yoshida K (2013) An improved *Bacillus subtilis* cell factory for producing *scyllo*-inositol, a promising therapeutic agent for Alzheimer's disease. Microb Cell Factories 12:124
- Thomas MP, Mills SJ, Potter BVL (2016) The "other" inositols and their phosphates: synthesis, biology, and medicine (with recent advances in myo-inositol chemistry). Angew Chem Int Ed Engl 55:1614–1650
- Tomlinson RV, Ballou CE (1962) Myo-inositol polyphosphates intermediates in the dephosphorylation of phytic acid by phytase. Biochemistry 1:166–171
- Turner BL, Richardson AE (2004) Identification of *scyllo*-inositol phosphates in soil by solution phosphorus-31 nuclear magnetic resonance spectroscopy. Soil Sci Soc Am J 68:802–808
- Turner BL, Paphazy MJ, Haygarth PM, McKelvie ID (2002) Inositol phosphates in the environment. Philos Trans R Soc Lond Ser B Biol Sci 357:449–469
- Turner BL, Cheesman AW, Godage HY, Riley AM, Potter BVL (2012) Determination of *neo*- and *D-chiro*-inositol hexakisphosphate in soils by solution 31P NMR spectroscopy. Environ Sci Technol 46:4994–5002
- Yamaoka M, Osawa S, Morinaga T, Takenaka S, Yoshida K (2011) A cell factory of *Bacillus subtilis* engineered for the simple bioconversion of *myo*-inositol to *scyllo*-inositol, a potential therapeutic agent for Alzheimer's disease. Microb Cell Factories 10:69
- Yoshida K, Sanbongi A, Murakami A, Suzuki H, Takenaka S, Takami H (2012) Three inositol dehydrogenases involved in utilization and interconversion of inositol stereoisomers in a thermophile, *Geobacillus kaustophilus* HTA426. Microbiology 158:1942–1952