



Siegel modular forms of degree two and level five

Haowu Wang¹ · Brandon Williams² 

Received: 2 January 2022 / Accepted: 11 June 2022 / Published online: 4 August 2022
© The Author(s) 2022

Abstract

We construct a ring of meromorphic Siegel modular forms of degree 2 and level 5, with singularities supported on an arrangement of Humbert surfaces, which is generated by four singular theta lifts of weights 1, 1, 2, 2 and their Jacobian. We use this to prove that the ring of holomorphic Siegel modular forms of degree 2 and level $\Gamma_0(5)$ is minimally generated by eighteen modular forms of weights 2, 4, 4, 4, 4, 4, 6, 6, 6, 6, 10, 11, 11, 11, 13, 13, 13, 15.

Keywords Siegel modular forms · Borcherds lifts · Rings of modular forms

Mathematics Subject Classification 11F46 · 11F27

1 Introduction

It is an interesting problem to determine the structure of rings of Siegel modular forms with respect to congruence subgroups. A famous theorem of Igusa [8] shows that every Siegel modular form of degree two and even weight for the full modular group $\mathrm{Sp}_4(\mathbb{Z})$ can be written uniquely as a polynomial in forms $\phi_4, \phi_6, \phi_{10}, \phi_{12}$ of weights 4, 6, 10, 12, and that odd-weight Siegel modular forms are precisely the products of even weight Siegel modular forms with a distinguished cusp form ψ_{35} of weight 35. It was proved by Aoki and Ibukiyama [1] that the rings of modular forms for the congruence subgroups:

H. Wang was supported by the Institute for Basic Science (IBS-R003-D1).

✉ Brandon Williams
brandon.williams@matha.rwth-aachen.de
Haowu Wang
haowu.wangmath@gmail.com

¹ Center for Geometry and Physics, Institute for Basic Science (IBS), Pohang 37673, Korea

² Lehrstuhl A für Mathematik, RWTH Aachen 52056, Aachen, Germany

$$\Gamma_{0,1}^{(2)}(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sp}_4(\mathbb{Z}) : c \equiv 0(N), \det(a) \equiv \det(d) \equiv 1(N) \right\}, \quad N = 2, 3, 4$$

have an analogous structure: they are generated by four algebraic independent modular forms together with their Jacobian (or first Rankin–Cohen–Ibukiyama bracket). The rings $M_*(\Gamma_0^{(2)}(N))$ where

$$\Gamma_0^{(2)}(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{Sp}_4(\mathbb{Z}) : c \equiv 0(N) \right\},$$

therefore, have a simple structure as well.

The goal in this paper is to extend the methods of Aoki and Ibukiyama to level $N = 5$. This is not quite straightforward, as the natural underlying ring is no longer $M_*(\Gamma_{0,1}^{(2)}(5))$ but rather a ring $M_*^!(\Gamma_{0,1}^{(2)}(5))$ of meromorphic Siegel modular forms with singularities on Humbert surfaces. We will define a hyperplane arrangement \mathcal{H} as the $\Gamma_0^{(2)}(5)$ -orbit of the Humbert surface

$$\{Z = \begin{pmatrix} \tau & z \\ z & w \end{pmatrix} \in \mathbb{H}_2 : \det(Z) = 1 - 5z\},$$

which, if one views points in \mathbb{H}_2 as parameterizing abelian surfaces, is a locus of principally polarized abelian surfaces with real multiplication that respects a $\Gamma_0^{(2)}(5)$ level structure. We then investigate the ring $M_*^!(\Gamma_{0,1}^{(2)}(5))$ of meromorphic Siegel modular forms on $\Gamma_{0,1}^{(2)}(5)$ with singularities supported on \mathcal{H} . Using a generalization of the modular Jacobian approach of [12], we prove in Theorem 3.6 that $M_*^!(\Gamma_{0,1}^{(2)}(5))$ is generated by four algebraically independent singular additive lifts f_1, f_2, g_1, g_2 of weights 1, 1, 2, 2 and by their Jacobian; in particular, the associated threefold $X_{0,1}^{(2)}(5)$ is rational. The local isomorphism from Sp_4 to $\mathrm{SO}(3, 2)$ and Borcherds’ theory of orthogonal modular forms with singularities are essential. $\mathrm{Proj}(M_*^!(\Gamma_{0,1}^{(2)}(5)))$ is the Looijenga compactification [9] of the complement $(\mathbb{H}_2 \setminus \mathcal{H})/\Gamma_{0,1}^{(2)}(5)$, which plays a similar role to the Satake–Baily–Borel compactification of $Y_{0,1}^{(2)}(5)$.

It follows from the above that every Siegel modular form of level $\Gamma_0^{(2)}(5)$ can be expressed uniquely in terms of the basic forms f_1, f_2, g_1, g_2 . It is not clear to the authors how to compute the ring $M_*(\Gamma_0^{(2)}(5))$ of (holomorphic) Siegel modular forms from this information alone; however, allowing a formula of Hashimoto [6] for the dimensions of cusp forms (itself an application of the Selberg trace formula), the ring structure becomes a straightforward Gröbner basis computation. We will prove that $M_*(\Gamma_0^{(2)}(5))$ is minimally generated by eighteen modular forms of weights 2, 4, 4, 4, 4, 4, 6, 6, 6, 6, 10, 11, 11, 11, 13, 13, 13, 15 in Theorem 4.2.

This paper is organized as follows. In Sect. 2, we review the realization of Siegel modular groups as orthogonal groups and the theory of Borcherds lifts. In Sect. 3, we determine two rings of meromorphic Siegel modular forms. In Sect. 4, we use this to determine the ring of holomorphic Siegel modular forms for $\Gamma_0^{(2)}(5)$.

2 Theta lifts to Siegel modular forms of degree two

2.1 $\Gamma_0^{(2)}(N)$ as an orthogonal group

Recall that the Pfaffian of an antisymmetric (4×4) -matrix M is

$$\text{pf}(M) = \text{pf} \begin{pmatrix} 0 & a & b & c \\ -a & 0 & d & e \\ -b & -d & 0 & f \\ -c & -e & -f & 0 \end{pmatrix} = af - be + cd.$$

We view pf as a quadratic form and define the associated bilinear form:

$$\langle x, y \rangle = \text{pf}(x + y) - \text{pf}(x) - \text{pf}(y).$$

The Pfaffian is invariant under conjugation $M \mapsto A^T M A$ by $A \in \text{SL}_4(\mathbb{R})$, and this action identifies $\text{SL}_4(\mathbb{R})$ with the Spin group $\text{Spin}(\text{pf})$. The symplectic group $\text{Sp}_4(\mathbb{R})$, by definition, preserves

$$\mathcal{J} = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

under conjugation, so it also preserves the orthogonal complement \mathcal{J}^\perp , and indeed, it is exactly the Spin group of pf restricted to \mathcal{J}^\perp . If the entries of M are labelled as above, then $M \in \mathcal{J}^\perp$ if and only if $b + e = 0$.

For any $N \in \mathbb{N}$, the group $\Gamma_0^{(2)}(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Sp}_4(\mathbb{Z}) : c \equiv 0(N) \right\}$ stabilizes the lattice

$$L = \left\{ M = \begin{pmatrix} 0 & a & b & c \\ -a & 0 & d & -b \\ -b & -d & 0 & f \\ -c & b & -f & 0 \end{pmatrix} : a, b, c, d, f \in \mathbb{Z}, a \equiv 0(N) \right\},$$

which is of type $U \oplus U(N) \oplus A_1$. By [7, Sect. 2] the special discriminant kernel $\widetilde{\text{SO}}(L)$ of L is exactly the projective modular group $\Gamma_{0,1}^{(2)}(N)/\{\pm I\}$ under this identification, where

$$\begin{aligned} \widetilde{\text{SO}}(L) &= \{g \in \text{SO}(L) : g(v) - v \in L \text{ for all } v \in L'\}, \\ \Gamma_{0,1}^{(2)}(N) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0^{(2)}(N) : \det(a) \equiv 1(N) \right\}. \end{aligned}$$

It follows that

$$\Gamma_0^{(2)}(N)/\{\pm I\} = \langle \widetilde{\text{SO}}(L), \varepsilon_u : u \in (\mathbb{Z}/N\mathbb{Z})^\times \rangle,$$

where ε_u is the matrix

$$\varepsilon_u = \begin{pmatrix} u & 0 & b & 0 \\ 0 & 1 & 0 & 0 \\ N & 0 & u^* & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in \Gamma_0^{(2)}(N)$$

for any integer solutions u^*, b to $uu^* - Nb = 1$ (the choice does not matter). The map induced by ε_u on $L'/L \cong A'_1/A_1 \oplus U(N)'/U(N)$ acts trivially on A'_1/A_1 and acts on $U(N)'/U(N) \cong \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}/N\mathbb{Z}$ as the map

$$\varepsilon_u : \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}/N\mathbb{Z} \rightarrow \mathbb{Z}/N\mathbb{Z} \oplus \mathbb{Z}/N\mathbb{Z}, (x, y) \mapsto (ux, u^{-1}y).$$

The symplectic group $\text{Sp}_4(\mathbb{R})$ acts on the Siegel upper half-space \mathbb{H}_2 by Möbius transformations:

$$M \cdot Z = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot Z = (aZ + b)(cZ + d)^{-1}.$$

Let $j(M; Z) = \det(cZ + d)$ be the usual automorphy factor. We embed the Siegel upper half-space into $L \otimes \mathbb{C}$ as follows:

$$Z = \begin{pmatrix} \tau & z \\ z & w \end{pmatrix} \mapsto \mathcal{Z} := \phi(Z) := \begin{pmatrix} 0 & 1 & z & w \\ -1 & 0 & -\tau & -z \\ -z & \tau & 0 & \tau w - z^2 \\ -w & z & z^2 - \tau w & 0 \end{pmatrix}.$$

Then one has the relation

$$M^T \mathcal{Z} M = j(M; Z)\phi(M \cdot Z), \quad M \in \text{Sp}_4(\mathbb{R})$$

as one can check on any system of generators.

For any $\lambda \in L'$ of positive norm $D = Q(\lambda)$, the space

$$\{Z \in \mathbb{H}_2 : \mathcal{Z} \in \lambda^\perp\}$$

is known as a **Humbert surface** $H(D, \lambda)$ of discriminant D . If λ is written in the form $\begin{pmatrix} 0 & a & b & c \\ -a & 0 & d & -b \\ -b & -d & 0 & f \\ -c & b & -f & 0 \end{pmatrix}$ then

$$H(D, \lambda) = \{Z = \begin{pmatrix} \tau & z \\ z & w \end{pmatrix} \in \mathbb{H}_2 : a\det(Z) - c\tau + 2bz + dw + f = 0\}.$$

If γ instead is a coset of L'/L , then we define

$$H(D, \gamma) = \sum_{\substack{\lambda \in \gamma \\ \lambda \text{ primitive in } L' \\ Q(\lambda)=D}} H(D, \lambda).$$

These unions are locally finite and, therefore, descend to well-defined divisors on $\tilde{O}(L) \setminus \mathbb{H}_2$. We will use the notation $H(D, \pm\gamma)$ because $\lambda^\perp = (-\lambda)^\perp$ implies $H(D, \gamma) = H(D, -\gamma)$. Note that many references omit the condition that λ is primitive in L' , so $H(D, \pm\gamma)$ satisfy inclusions; our divisors $H(D, \pm\gamma)$ do not.

2.2 Theta lifts

Let L be the lattice in the space of (4×4) antisymmetric matrices from the previous subsection. The **weight k theta kernel** is

$$\Theta_k(\tau; Z) = \frac{\pi^k}{\det(V)^k \Gamma(k)} \sum_{\lambda \in L'} \langle \lambda, Z \rangle^k e^{-\frac{\pi y}{\det(V)} |\langle \lambda, Z \rangle|^2} e^{2\pi i \bar{\tau} \text{pf}(\lambda)} \mathbf{e}_\lambda,$$

where $\tau = x + iy \in \mathbb{H}$ and $Z = U + iV \in \mathbb{H}_2$; and \mathcal{Z} is the image of Z in $L \otimes \mathbb{C}$. By applying a theorem of Vignéras on indefinite theta series [11] one can deduce the behaviour of Θ_k under the action of $\text{SL}_2(\mathbb{Z})$ on τ : it transforms like a modular form of weight $\kappa := k - 1/2$ with respect to the Weil representation ρ_L (see Definition 2.1). On the other hand, for any $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0^{(2)}(N)$,

$$\begin{aligned} \Theta_k(\tau; M \cdot Z) &= \frac{\pi^k}{\det \text{im}(M \cdot Z)^k \Gamma(k)} \sum_{\lambda \in L'} \det(cZ + d)^{-k} \langle \lambda, M^T \mathcal{Z} M \rangle \\ &\quad \times e^{-\frac{\pi y}{\det(V)} |\langle \lambda, M^T \mathcal{Z} M \rangle|^2} e^{2\pi i \bar{\tau} \text{pf}(\lambda)} \mathbf{e}_\lambda \\ &= \frac{\pi^k}{\det(V)^k \Gamma(k)} \overline{\det(cZ + d)^k} \sum_{\lambda \in L'} \langle M^{-T} \lambda M^{-1}, \mathcal{Z} \rangle^k \\ &\quad \times e^{-\frac{\pi y}{\det(V)} |\langle M^{-T} \lambda M^{-1}, \mathcal{Z} \rangle|^2} e^{2\pi i \bar{\tau} \text{pf}(M^{-T} \lambda M^{-1})} \mathbf{e}_\lambda \\ &= \overline{\det(cZ + d)^k} \sigma(M) \Theta_k(\tau; Z), \end{aligned}$$

where σ is the map

$$\sigma : \Gamma_0^{(2)}(N) \longrightarrow \text{Aut } \mathbb{C}[L'/L], \quad \sigma(M) \mathbf{e}_\lambda := \mathbf{e}_{M^T \lambda M}.$$

Following Borcherds [3], one defines the theta lift of a vector-valued modular form F with a pole at ∞ as the regularized integral of F against the kernel Θ_k :

Definition 2.1 (i) The *Weil representation* ρ_L associated to an even lattice (L, Q) is the representation $\rho : \text{Mp}_2(\mathbb{Z}) \rightarrow \text{GL } \mathbb{C}[L'/L]$ defined by

$$\begin{aligned} \rho \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, 1 \right) \epsilon_\gamma &= e^{-2\pi i Q(\gamma)} \epsilon_\gamma; \\ \rho \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \sqrt{\tau} \right) \epsilon_\gamma &= e^{\pi i \text{sig}(L)/4} |L'/L|^{-1/2} \sum_{\beta \in L'/L} e^{2\pi i \langle \gamma, \beta \rangle} \epsilon_\beta, \end{aligned}$$

where $\text{Mp}_2(\mathbb{Z})$ is the metaplectic group of pairs (M, ϕ) where $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$ and ϕ is a square root of $c\tau + d$, and $\epsilon_\gamma, \gamma \in L'/L$ is the standard basis of the group ring $\mathbb{C}[L'/L]$.

(ii) A *weakly holomorphic vector-valued modular form* for ρ_L is a holomorphic function $F : \mathbb{H} \rightarrow \mathbb{C}[L'/L]$ which satisfies

$$F((M, \phi) \cdot \tau) = \phi(\tau)^{2k} \rho_L(M) F(\tau), \quad (M, \phi) \in \text{Mp}_2(\mathbb{Z})$$

and which is meromorphic at the cusp ∞ , i.e. its Fourier series has only finitely many negative exponents.

(iii) Let $k \geq 1$ and let $F \in M_k^1(\rho_L)$ be a weakly holomorphic modular form. The **(singular) theta lift** of F is

$$\Phi_F(Z) = \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}^{\text{reg}} \langle F(\tau), \Theta_k(\tau; Z) \rangle y^\kappa \frac{dx \, dy}{y^2}.$$

Here, the regularization means one takes the limit as $w \rightarrow \infty$ of the integral over $\mathcal{F}_w = \{\tau = x + iy \in \mathbb{H} : x^2 + y^2 \geq 1, |x| \leq 1/2, y \leq w\}$; in effect, it means one integrates first with respect to x , which mollifies the contribution of the principal part of F to the integral; and then secondly with respect to y . The behaviour of the theta lift under Möbius transformations is

$$\begin{aligned} \Phi_F(M \cdot Z) &= \int^{\text{reg}} \langle F(\tau), \Theta_k(\tau; M \cdot Z) \rangle y^{\kappa-2} \, dx \, dy \\ &= \det(cZ + d)^k \int^{\text{reg}} \sum_{\gamma \in L'/L} F_\gamma(\tau) \overline{\Theta_{k; M^{-T}\gamma M^{-1}}(\tau; Z)} y^{\kappa-2} \, dx \, dy \\ &= \det(cZ + d)^k \Phi_{\sigma(M)^{-1}F}(Z). \end{aligned}$$

Therefore, the singular theta lift Φ_F transforms like a Siegel modular form of weight k on the subgroup of $\Gamma_0^{(2)}(N)$ that fixes F . Borcherds’ results [3, Theorem 14.3] show that Φ_F is meromorphic, with singularities of multiplicity k along Humbert surfaces associated to the principal part of the input F . This is the so-called Borcherds additive lift. Since the Borcherds additive lift is a generalization of the Gritsenko lift [5], we will also call it the singular Gritsenko lift. When the input F has weight $\kappa = -\frac{1}{2}$ (i.e. $k = 0$), the modified exponential of Φ_F defines a remarkable modular form which has an infinite product expansion (cf. [3, Theorem 13.3]), called a *Borcherds product*, or more specifically the Borcherds lift of F . In this paper we will need both types of singular theta lifts.

Remark 2.2 It is often useful to consider the *pullback* or restriction of a Siegel modular form to a Humbert surface. The result is traditionally interpreted as a Hilbert modular

form attached to a real-quadratic field. From the point of view of orthogonal modular forms, this is very simple: to restrict a form $\Phi(Z)$ to the sublattice v^\perp (with $v \in L'$), one simply restricts to Z satisfying $\langle Z, v \rangle = 0$.

The pullback of a theta lift Φ_F as above is again a theta lift, $\Phi_{\vartheta F}$, where $\vartheta F \in M_{k+1/2}(\rho_{v^\perp})$ is the *theta contraction*; this is the vector-valued modular form associated to the Weil representation of $L_v = L \cap v^\perp$ characterized by

$$\langle \vartheta F, \Theta_{k;L_v}(\tau; Z) \rangle = \langle F, \Theta_{k;L}(\tau; Z) \rangle, \quad Z \in v^\perp,$$

where $\Theta_{k;L_v}$ and $\Theta_{k;L}$ are the weight k theta kernels attached to L_v and L , respectively. More explicitly one can define ϑF as the zero-value of a vector-valued Jacobi form for the Weil representation attached to L_v whose associated theta decomposition is F itself [14]. The important point is that one can check rigorously whether a theta lift Φ_F vanishes identically on a Heegner divisor, with the computations taking place only on the level of vector-valued modular forms.

3 The ring of meromorphic Siegel modular forms of level 5

We consider the ring $M_*^!(\Gamma_0^{(2)}(5))$ of meromorphic Siegel modular forms of level $\Gamma_0^{(2)}(5)$ whose poles may lie only on the orbit \mathcal{H} of the Humbert surface:

$$\{Z = \begin{pmatrix} \tau & z \\ z & w \end{pmatrix} \in \mathbb{H}_2 : \det(Z) = 1 - 5z\},$$

which is a locus of principally polarized RM abelian surfaces with $\Gamma_0^{(2)}(5)$ level structure. In view of the discussion of Sect. 2.1, \mathcal{H} splits as the union of two irreducible $\Gamma_{0,1}^{(2)}(5)$ -orbits of Humbert surfaces:

$$\mathcal{H} = H(1/20, \pm\gamma_1) + H(1/20, \pm\gamma_2),$$

each invariant under the discriminant kernel of $L = U \oplus U(5) \oplus A_1$, where we have fixed any coset $\gamma_1 \in L'/L$ of norm $1/20 + \mathbb{Z}$ and define $\gamma_2 = \varepsilon_2(\gamma_1)$. The Humbert surface $H_{1/5}$ of discriminant $1/5$, the orbit of $\{(\begin{smallmatrix} \tau & z \\ z & w \end{smallmatrix}) \in \mathbb{H}_2 : \tau = 2z\}$ under $\Gamma_0^{(2)}(5)$, also splits into two $\Gamma_{0,1}^{(2)}(5)$ -invariant divisors:

$$H_{1/5} = H(1/5, \pm\delta_1) + H(1/5, \pm\delta_2),$$

where $\delta_n = 2\gamma_n \in L'/L$.

For a finite-index subgroup $\Gamma \leq \Gamma_0^{(2)}(5)$ or $\Gamma \leq O(L)$, we define $M_*^!(\Gamma, \chi)$ to be the ring of meromorphic forms, holomorphic away from \mathcal{H} , which are modular under Γ with character χ .

We first prove a form of Koecher’s principle for meromorphic modular forms with poles supported on \mathcal{H} .

Lemma 3.1 *Let $f \in M_k^!(\Gamma_{0,1}^{(2)}(5), \chi)$. If k is negative, then f is identically zero. If $k = 0$, then f is constant.*

Proof We prove the lemma in the context of $O(3, 2)$. Let v and $u \neq \pm v$ be primitive vectors of norm $1/20$ in L' , such that $v^\perp, u^\perp \in \mathcal{H}$. Suppose that f is not identically zero and has poles of multiplicity c_v along v^\perp . We denote the intersection of v^\perp and the symmetric domain \mathbb{H}_2 (resp. the lattice L) by $v^\perp \cap \mathbb{H}_2$ (resp. L_v). Then L_v is a lattice of signature $(2, 2)$ and discriminant 5 , equivalent to the lattice $U + \mathbb{Z}[(1 + \sqrt{5})/2]$ where the quadratic form is the field norm. It is easy to see that the space $L_v \otimes \mathbb{Q}$ contains no isotropic planes, so the Koecher principle holds for modular forms on $\tilde{O}(L_v)$. We find that the projection of u in L_v has non-positive norm, which implies that the intersection of u^\perp and $v^\perp \cap \mathbb{H}_2 \cong \mathbb{H} \times \mathbb{H}$ is empty. Thus, the quasi-pullback of f to $v^\perp \cap \mathbb{H}_2$, i.e. the leading term in the power series expansion about that hyperplane, is a nonzero holomorphic modular form of weight $k - c_v$. By Koecher's principle, we conclude $k - c_v \geq 0$ and, therefore, $k \geq 0$, and when $k = 0$, we must have $c_v = 0$, and thus, f is holomorphic and must be constant (by Koecher's principle on $\tilde{O}(L)$).

We now construct some basic modular forms using Borcherds additive lifts (singular Gritsenko lifts) and Borcherds products.

Lemma 3.2 *There are singular Gritsenko lifts f_1, f_2 of weight one on $\tilde{O}(L)$ whose divisors are exactly*

$$\text{div}(f_1) = -H(1/20, \pm\gamma_1) + 4H(1/20, \pm\gamma_2) + H(1/5, \pm\delta_1)$$

and

$$\text{div}(f_2) = 4H(1/20, \pm\gamma_1) - H(1/20, \pm\gamma_2) + H(1/5, \pm\delta_2).$$

Proof Using the algorithm of [13], we find a weakly holomorphic modular form of weight $1/2$ for the Weil representation associated to L for which the Fourier expansion takes the form:

$$2q^{-1/20}(\mathbf{e}_{\gamma_1} - \mathbf{e}_{-\gamma_1}) + O(q^{1/20}),$$

which is mapped under the Gritsenko lift to a meromorphic form f_1 with simple poles only on $H(1/20, \pm\gamma_1)$ and $H(1/20, \pm\gamma_4)$. Applying the automorphism ε_2 on L'/L to the input into f_1 yields the input into f_2 .

On the other hand, we found a weakly holomorphic modular form of weight $-1/2$ for which the principal part at ∞ is

$$2\mathbf{e}_0 - 2q^{-1/20}(\mathbf{e}_{\gamma_1} + \mathbf{e}_{-\gamma_1}) + 4q^{-1/20}(\mathbf{e}_{\gamma_2} + \mathbf{e}_{-\gamma_2}) + q^{-1/5}(\mathbf{e}_{\delta_1} + \mathbf{e}_{-\delta_1}),$$

which is mapped under the Borcherds lift to a meromorphic modular form F_1 (possibly with character) of weight one and the claimed divisor. By taking theta contractions of the input form, one finds that f_1 vanishes on $H(1/5, \pm\delta_1)$. Then the quotient f_1/F_1 lies in $M_0^1(\tilde{O}(L), \chi)$ so it is constant by Lemma 3.1.

Remark 3.3 The Fourier expansions of f_1 and f_2 begin

$$\begin{aligned}
 f_1\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= 1 + 3q + 3s + 4q^2 + (2r^{-1} + 6 + 2r)qs + 4s^2 + O(q, s)^3; \\
 f_2\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= q - s - 2q^2 + 2s^2 + 4q^3 + (4r^2 + 2 + 4r)qs(q - s) \\
 &\quad - 4s^3 + O(q, s)^4,
 \end{aligned}$$

where as usual $q = e^{2\pi i\tau}$, $r = e^{2\pi iz}$, $s = e^{2\pi iw}$. For more coefficients, see Fig. 1. Setting $s = 0$, one obtains the (holomorphic) modular forms:

$$\Phi(f_1) = 1 + 3q + 4q^2 \pm \dots, \quad \Phi(f_2) = q - 2q^2 + 4q^3 \pm \dots$$

of weight one and level $\Gamma_1(5)$ which freely generate the ring $M_*(\Gamma_1(5))$.

There are nine Heegner divisors of discriminant $1/4$. One is the mirror of the reflective vector $r = 1/2 \in A'_1$, represented by the diagonal in \mathbb{H}_2 , and the other eight are of the form $H(1/4, r + \gamma)$ where γ are the isotropic cosets of $U(5)' / U(5)$. It will be convenient to fix concrete representatives. We take the Gram matrix $\mathbf{S} = \begin{pmatrix} 0 & 0 & 5 \\ 0 & 2 & 0 \\ 5 & 0 & 0 \end{pmatrix}$ for $U(5) \oplus A_1$, such that $L'/L \cong \mathbf{S}^{-1}\mathbb{Z}^3/\mathbb{Z}^3$ and fix the cosets

$$\begin{aligned}
 \gamma_1 &= (1/5, 1/2, 4/5) + L, & \gamma_2 &= (2/5, 1/2, 2/5) + L, \\
 \gamma_3 &= (3/5, 1/2, 3/5) + L, & \gamma_4 &= (4/5, 1/2, 1/5) + L,
 \end{aligned}$$

of norm $1/20 + \mathbb{Z}$. The norm $1/4$ cosets other than r are labelled

$$\alpha_n = (n/5, 1/2, 0) + L, \quad \beta_n = (0, 1/2, n/5) + L, \quad n \in \{1, 2, 3, 4\}.$$

Lemma 3.4 *There are singular Gritsenko lifts g_1, g_2, h_1, h_2 of weight two on $\tilde{\mathcal{O}}(L)$ whose divisors are exactly*

$$\begin{aligned}
 \text{div } g_1 &= 3H(1/20, \pm\gamma_1) - 2H(1/20, \pm\gamma_2) + H(1/4, \pm\alpha_2); \\
 \text{div } g_2 &= -2H(1/20, \pm\gamma_1) + 3H(1/20, \pm\gamma_2) + H(1/4, \pm\alpha_1); \\
 \text{div } h_1 &= 3H(1/20, \pm\gamma_1) - 2H(1/20, \pm\gamma_2) + H(1/4, \pm\beta_2); \\
 \text{div } h_2 &= -2H(1/20, \pm\gamma_1) + 3H(1/20, \pm\gamma_2) + H(1/4, \pm\beta_1).
 \end{aligned}$$

Proof The proof is essentially the same argument as Lemma 3.2. Using the pullback trick, one constructs weight two Gritsenko lifts which vanish on the claimed discriminant $1/4$ Heegner divisors. Then one constructs Borcherds products of weight two with the claimed divisors. The respective quotients lie in $M_0^1(\tilde{\mathcal{O}}(L), \chi)$ and are, therefore, constant by Lemma 3.1. To determine the precise (weakly holomorphic) vector-valued modular forms which lift to g_1, g_2, h_1, h_2 , one only needs to compute the four-dimensional space of weakly holomorphic forms of weight $3/2$ for ρ_L with a pole of order at most $1/20$ at ∞ and identify the unique (up to scalar) forms whose pullback to α_n^\perp or β_n^\perp is respectively zero. The input forms G_1, G_2, H_1, H_2 can be chosen such that their Fourier expansions begin as follows:

$$\begin{aligned}
 G_1 &: q^{-1/20}(\mathbf{e}_{\gamma_2} + \mathbf{e}_{\gamma_3}) + (\mathbf{e}_{(0,0,1/5)} + \mathbf{e}_{(0,0,4/5)} - \mathbf{e}_{(0,0,2/5)} + \mathbf{e}_{(0,0,3/5)}) + O(q^{1/20}), \\
 G_2 &: q^{-1/20}(\mathbf{e}_{\gamma_1} + \mathbf{e}_{\gamma_4}) + (\mathbf{e}_{(0,0,2/5)} + \mathbf{e}_{(0,0,3/5)} - \mathbf{e}_{(0,0,1/5)} + \mathbf{e}_{(0,0,4/5)}) + O(q^{1/20}), \\
 H_1 &: q^{-1/20}(\mathbf{e}_{\gamma_2} + \mathbf{e}_{\gamma_3}) + (\mathbf{e}_{(1/5,0,0)} + \mathbf{e}_{(4/5,0,0)} - \mathbf{e}_{(2/5,0,0)} + \mathbf{e}_{(3/5,0,0)}) + O(q^{1/20}), \\
 H_2 &: q^{-1/20}(\mathbf{e}_{\gamma_1} + \mathbf{e}_{\gamma_4}) + (\mathbf{e}_{(2/5,0,0)} + \mathbf{e}_{(3/5,0,0)} - \mathbf{e}_{(1/5,0,0)} + \mathbf{e}_{(4/5,0,0)}) + O(q^{1/20}).
 \end{aligned}$$

These expansions determine G_1, G_2, H_1, H_2 uniquely because there are no vector-valued cusp forms of weight $3/2$ for ρ_L .

Remark 3.5 The Fourier expansions of g_1, g_2, h_1, h_2 begin as follows:

$$\begin{aligned}
 g_1\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= q + q^2 - 5qs + 2q^3 + (-3r^{-1} + 1 - 3r)q^2s \\
 &\quad + (-r^{-1} + 7 - r)qs^2 + O(q, s)^4; \\
 g_2\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= -q - q^2 + (r^{-1} + 3 + r)qs - 2q^3 + (-r^{-1} + 7 - r)q^2s \\
 &\quad + (-3r^{-1} + 1 - 3r)qs^2 + O(q, s)^4; \\
 h_1\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= s - 5qs + s^2 + (-r^{-1} + 7 - r)q^2s \\
 &\quad + (-3r^{-1} + 1 - 3r)qs^2 + 2s^3 + O(q, s)^4; \\
 h_2\left(\begin{pmatrix} \tau & z \\ z & w \end{pmatrix}\right) &= -s + (r^{-1} + 3 + r)qs - s^3 + (-3r^{-1} + 1 - 3r)q^2s \\
 &\quad + (-r^{-1} + 7 - r)qs^2 - 2s^3 + O(q, s)^4.
 \end{aligned}$$

We can now determine the structure of $M_*^1(\Gamma_{0,1}^{(2)}(5))$. Recall that $\Gamma_{0,1}^{(2)}(5)/\{\pm I\} \cong \widetilde{\text{SO}}(L)$. The decomposition

$$M_k^1(\widetilde{\text{SO}}(L)) = M_k^1(\widetilde{\text{O}}(L)) \oplus M_k^1(\widetilde{\text{O}}(L), \det)$$

suggests that we first consider the ring of modular forms for the discriminant kernel $\widetilde{\text{O}}(L)$. We will show that $M_*^1(\widetilde{\text{O}}(L))$ is freely generated using a generalization of the modular Jacobian approach of [12, Theorem 5.1]. We briefly introduce the main objects of this approach. For any four $\psi_i \in M_{k_i}^1(\widetilde{\text{O}}(L))$ with $1 \leq i \leq 4$, their Jacobian (see [12, Theorem 2.5] and [1, Proposition 2.1])

$$J(\psi_1, \psi_2, \psi_3, \psi_4) = \begin{vmatrix} k_1\psi_1 & k_2\psi_2 & k_3\psi_3 & k_4\psi_4 \\ \frac{\partial\psi_1}{\partial\tau} & \frac{\partial\psi_2}{\partial\tau} & \frac{\partial\psi_3}{\partial\tau} & \frac{\partial\psi_4}{\partial\tau} \\ \frac{\partial\psi_1}{\partial z} & \frac{\partial\psi_2}{\partial z} & \frac{\partial\psi_3}{\partial z} & \frac{\partial\psi_4}{\partial z} \\ \frac{\partial\psi_1}{\partial w} & \frac{\partial\psi_2}{\partial w} & \frac{\partial\psi_3}{\partial w} & \frac{\partial\psi_4}{\partial w} \end{vmatrix}$$

lies in $M_{k_1+k_2+k_3+k_4+3}^1(\widetilde{\text{O}}(L), \det)$. The Jacobian $J(\psi_1, \psi_2, \psi_3, \psi_4)$ is not identically zero if and only if the four forms ψ_i are algebraically independent over \mathbb{C} .

The discriminant kernel $\widetilde{\text{O}}(L)$ contains reflections associated to vectors of norm 1 in L (the so-called 2-reflections)

$$\sigma_v : x \mapsto x - (x, v)v.$$

The hyperplane v^\perp is called the mirror of the reflection σ_v . Since $\det(\sigma_v) = -1$, the chain rule implies that the above Jacobian vanishes on all mirrors of 2-reflections. Conversely, the main theorem of [12], and its generalization to meromorphic modular forms with constrained poles, implies that

Theorem 3.6 *The ring $M_*^1(\tilde{\mathcal{O}}(L))$ is a free algebra:*

$$M_*^1(\tilde{\mathcal{O}}(L)) = \mathbb{C}[f_1, f_2, g_1, g_2].$$

Define $J := J(f_1, f_2, g_1, g_2)$. Then

$$M_*^1(\Gamma_{0,1}^{(2)}(5)) = \mathbb{C}[f_1, f_2, g_1, g_2, J].$$

Proof The Jacobian J of f_1, f_2, g_1, g_2 has weight 9 and vanishes on the mirrors of 2-reflections, which form a union of Heegner divisors of discriminants 1/4 and 1 denoted Δ . Using the Fourier expansions of the forms it is easy to check that J is not identically zero. Using the algorithm of [13], we find a Borcherds product J_0 with divisor

$$\text{div } J_0 = \Delta + 6H(1/20, \pm\gamma_1) + 6H(1/20, \pm\gamma_2).$$

The quotient J/J_0 lies in $M_{\chi}^1(\Gamma_{0,1}^{(2)}(5))$ and is, therefore, a constant denoted c by Lemma 3.1. We will now prove the claim by an argument which appeared essentially in [12]. Suppose that $M_*^1(\tilde{\mathcal{O}}(L))$ was not generated by $h_1 := f_1, h_2 := f_2, h_3 := g_1$ and $h_4 := g_2$, and let $h_5 \in M_{k_5}^1(\tilde{\mathcal{O}}(L))$ be a modular form of minimal weight which is not contained in $\mathbb{C}[f_1, f_2, g_1, g_2]$. Set $k_1 = k_2 = 1$ and $k_3 = k_4 = 2$, such that k_i is the weight of h_i . For $1 \leq j \leq 5$, we define J_j as the Jacobian of the four modular forms h_i omitting h_j , such that $cJ_0 = J = J_5$. It is clear that $g_j := J_j/J$ is a modular form on $\tilde{\mathcal{O}}(L)$ with poles supported on \mathcal{H} . We compute the determinant and find the identity:

$$0 = \det \begin{pmatrix} k_1 h_1 & \dots & k_4 h_4 & k_5 h_5 \\ k_1 h_1 & \dots & k_4 h_4 & k_5 h_5 \\ \nabla h_1 & \dots & \nabla h_4 & \nabla h_5 \end{pmatrix} = \sum_{i=1}^5 (-1)^{i+1} k_i h_i J_i.$$

Since $J_i = J g_i$ and $g_5 = 1$, we have

$$\sum_{i=1}^5 (-1)^{i+1} k_i h_i g_i = 0, \quad \text{i.e.} \quad k_5 h_5 = \sum_{i=1}^4 (-1)^i h_i g_i.$$

Since h_5 was chosen to have minimal weight, $g_i \in \mathbb{C}[h_1, h_2, h_3, h_4]$ for all i , and thus, $h_5 \in \mathbb{C}[h_1, h_2, h_3, h_4]$, which is a contradiction.

Now any $h \in M_k^1(\tilde{\mathcal{O}}(L))$ vanishes on all mirrors of 2-reflections, which implies that $h/J \in M_{k-9}^1(\tilde{\mathcal{O}}(L))$. Therefore,

$$M_*^1(\Gamma_{0,1}^{(2)}(5)) = M_*^1(\tilde{\mathcal{S}}\tilde{\mathcal{O}}(L)) = M_*^1(\tilde{\mathcal{O}}(L)) \oplus M_*^1(\tilde{\mathcal{O}}(L), \det)$$

is generated by f_1, f_2, g_1, g_2 , and J .

Remark 3.7 The weight two singular Gritsenko lifts satisfy the relations:

$$g_1 - h_1 = h_2 - g_2 = f_1 f_2.$$

The product $f_1 f_2$ is holomorphic and in fact itself a Gritsenko lift, but it has a quadratic character under $\Gamma_0^{(2)}(5)$. There is a unique normalized Siegel modular form e_2 of weight two for $\Gamma_0^{(2)}(5)$, which can be constructed as the Gritsenko lift of the unique vector-valued modular form of weight $3/2$ for ρ_L invariant under all automorphisms of the discriminant form. (The uniqueness follows from Corollary 3.8.) In terms of the generators of $M_*^!(\Gamma_{0,1}^{(2)}(5))$, a computation shows

$$e_2 = f_1^2 + f_2^2 - 4(g_1 + g_2).$$

Corollary 3.8 *The ring $M_*^!(\Gamma_0^{(2)}(5))$ is minimally generated in weights 2, 2, 4, 4, 4, 4, 4, 11, 11, 11 by the ten forms*

$$\begin{aligned} f_1^2 + f_2^2, \quad e_2, \quad f_1^2 g_1 + f_2^2 g_2, \quad f_1 f_2 (g_1 - g_2), \quad f_1 f_2 (f_1 - f_2)(f_1 + f_2), \\ f_1^2 f_2^2, \quad g_1 g_2, \quad J f_1 f_2, \quad J(f_1^2 - f_2^2), \quad J(g_1 - g_2). \end{aligned}$$

Proof The group $\Gamma_0^{(2)}(5)$ is generated by the special discriminant kernel of L and by the order four automorphism ε_2 which acts on the generators of $M_*^!(\Gamma_{0,1}^{(2)}(5))$ by

$$\varepsilon_2 : f_1 \mapsto f_2, \quad f_2 \mapsto -f_1, \quad g_1 \mapsto g_2, \quad g_2 \mapsto g_1, \quad J \mapsto -J$$

as one can see on the input functions into the Gritsenko lifts. We conclude the action of ε_2 on J (as a Jacobian) from the actions of ε_2 on f_1, f_2, g_1 , and g_2 . The expressions in f_1, f_2, g_1, g_2, J in the claim generate the ring of invariants under this action.

Remark 3.9 The same argument shows that the kernel of $\varepsilon_2^2 = \varepsilon_4$ is generated by J and by the weight two forms:

$$f_1^2, f_2^2, f_1 f_2, g_1, g_2.$$

This corresponds to the quadratic Nebentypus $\chi \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = \left(\frac{5}{\det d} \right)$ on $\Gamma_0^{(2)}(5)$. Note that $f_1 f_2$ is the Siegel Eisenstein series of weight two for the character χ , and that the Jacobian J is the unique cusp form of weight nine for χ up to scalars.

Remark 3.10 There is a seven-dimensional space of modular forms of weight $7/2$ for ρ_L , and a four-dimensional subspace on which ε_2 acts trivially, so the weight four Maass space for $\Gamma_0^{(2)}(5)$ is four dimensional. Using the structure theorem above, we can identify it by comparing only a few Fourier coefficients:

$$\text{Maass}_4 = \text{Span}(\phi_1, \phi_2, \phi_3, \phi_4),$$

Table 1 $\dim M_k(\Gamma_0^{(2)}(5))$

k	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
dim	0	1	0	6	0	10	0	22	0	34	3	57	6	79	16	117	25	153	45

where

$$\begin{aligned} \phi_1 &= e_2^2 + f_1^2 f_2^2; \\ \phi_2 &= f_1^2 g_1 + f_2^2 g_2 - 2g_1 g_2; \\ \phi_3 &= f_1 f_2 (f_1^2 - 2f_1 f_2 - f_2^2 + 2g_1 - 2g_2); \\ \phi_4 &= 2g_1 g_2 + f_1 f_2 (g_1 - g_2). \end{aligned}$$

The form ϕ_4 is a cusp form and indeed spans $S_4(\Gamma_0^{(2)}(5))$, which was shown to be one dimensional by Poor and Yuen [10].

4 The ring of holomorphic Siegel modular forms of level 5

In this section, we investigate the ring $M_*(\Gamma_0^{(2)}(5))$ of holomorphic Siegel modular forms for $\Gamma_0^{(2)}(5)$. We will need the Hilbert–Poincaré series for this ring, which can be derived from dimension formulas available in the literature.

Theorem 4.1 *The Hilbert–Poincaré series of dimensions of modular forms for $\Gamma_0^{(2)}(5)$ is*

$$\sum_{k=0}^{\infty} \dim M_k(\Gamma_0^{(2)}(5))t^k = \frac{(1-t)^2(1+t^7)P(t)}{(1-t^2)^2(1-t^3)(1-t^4)^2(1-t^5)},$$

where $P(t)$ is the irreducible palindromic polynomial

$$\begin{aligned} P(t) &= 1 + 2t + 2t^2 + t^3 + 3t^4 + 5t^5 + 8t^6 + 8t^7 + 8t^8 + 5t^9 \\ &\quad + 3t^{10} + t^{11} + 2t^{12} + 2t^{13} + t^{14}. \end{aligned}$$

The first values of $\dim M_k(\Gamma_0^{(2)}(5))$ are given in Table 1.

Proof The dimensions of the spaces of cusp forms of weight $k \geq 5$ have been computed in closed form by Hashimoto by means of the Selberg trace formula and in lower weights by Poor and Yuen [10]: we have $\dim S_4(\Gamma_0^{(2)}(5)) = 1$ and $\dim S_k(\Gamma_0^{(2)}(5)) = 0$ for $k \leq 3$. All odd-weight modular forms are cusp forms, and by a more general theorem of Böcherer–Ibukiyama [2], for even $k > 2$,

$$\dim M_k(\Gamma_0^{(2)}(5)) = \dim S_k(\Gamma_0^{(2)}(5)) + 2 \cdot \dim S_k(\Gamma_0(5)) + 3.$$

We can now determine the generators of $M_*(\Gamma_0^{(2)}(5))$ using Corollary 3.8 together with the above generating series.

Theorem 4.2 *The ring of Siegel modular forms of level $\Gamma_0^{(2)}(5)$ is minimally generated by the weight two form*

$$e_2 = f_1^2 + f_2^2 - 4g_1 - 4g_2,$$

five weight four forms

$$f_1^2 g_1 + f_2^2 g_2, \quad f_1 f_2 (g_1 - g_2), \quad f_1 f_2 (f_1^2 - f_2^2), \quad f_1^2 f_2^2, \quad g_1 g_2,$$

four weight six forms

$$f_1^2 f_2^2 (g_1 + g_2), \quad f_1^3 f_2 g_1 - f_2^3 f_1 g_2, \quad f_1^2 g_1^2 + f_2^2 g_2^2, \quad g_1 g_2 (f_1^2 + f_2^2),$$

the weight ten form

$$f_1^2 f_2^2 (f_1^2 + f_2^2)^3,$$

three weight eleven forms

$$f_1 f_2 J, \quad (f_1^2 - f_2^2) J, \quad (g_1 - g_2) J,$$

three weight thirteen forms

$$(f_1^2 + f_2^2) f_1 f_2 J, \quad (f_1^4 - f_2^4) J, \quad (f_1^2 + f_2^2) (g_1 - g_2) J,$$

and the weight fifteen form

$$(f_1^2 - f_2^2)^3 J.$$

Proof From the divisors of f_1 , f_2 , g_1 , g_2 , and J , it is easy to see that all of the forms above (except for e_2 , which was discussed in the previous section) are holomorphic and ε_2 -invariant. The Hilbert series of this ring was computed in Macaulay2 [4] and coincides exactly with the series predicted by Theorem 4.1, so we can conclude that these forms are sufficient to generate all holomorphic Siegel modular forms.

Funding Open Access funding enabled and organized by Projekt DEAL.

Declarations

Conflict of interest Data sharing is not applicable to this article as no datasets were generated or analysed during the current study. The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix

In Figs. 1 and 2, we list more Fourier coefficients of the basic meromorphic forms f_1 , f_2 , g_1 , g_2 , as well as the unique expression for J^2 as a polynomial in these forms.

Note that the polynomial representing J^2 must split into two irreducible factors, corresponding to the two classes of reflections whose mirrors lie in the divisor of J . One of these factors is $g_1 + g_2$.

<i>a</i>	<i>b</i>	<i>c</i>	<i>f</i> ₁	<i>f</i> ₂	<i>g</i> ₁	<i>g</i> ₂	<i>a</i>	<i>b</i>	<i>c</i>	<i>f</i> ₁	<i>f</i> ₂	<i>g</i> ₁	<i>g</i> ₂
0	0	0	1	0	0	0	4	1	2	-8	8	12	4
1	0	0	3	1	1	-1	4	2	2	6	-6	-4	-8
0	0	1	3	-1	0	0	3	-3	3	2	0	0	3
2	0	0	4	-2	1	-1	3	-2	3	6	0	-5	-11
1	-1	1	2	0	0	1	3	-1	3	-6	0	40	1
1	0	1	6	0	-5	3	3	0	3	16	0	-50	-6
1	1	1	2	0	0	1	3	1	3	-6	0	40	1
0	0	2	4	2	0	0	3	2	3	6	0	-5	-11
3	0	0	2	4	2	-2	3	3	3	2	0	0	3
2	-1	1	4	-4	-3	-1	2	-2	4	6	6	-8	-4
2	0	1	2	-2	1	7	2	-1	4	-8	-8	4	12
2	1	1	4	-4	-3	-1	2	0	4	14	14	-7	-1
1	-1	2	4	4	-1	-3	2	1	4	-8	-8	4	12
1	0	2	2	2	7	1	2	2	4	6	6	-8	-4
1	1	2	4	4	-1	-3	1	-2	5	3	1	1	-1
0	0	3	2	-4	0	0	1	-1	5	-2	-4	-9	4
4	0	0	1	-3	3	-3	1	0	5	8	6	11	-1
3	-1	1	6	6	-2	4	1	1	5	-2	-4	-9	4
3	0	1	-2	-2	-6	2	1	2	5	3	1	1	-1
3	1	1	6	6	-2	4	0	0	6	6	-2	0	0
2	-2	2	2	0	0	2	7	0	0	4	-2	6	-6
2	-1	2	8	0	10	-2	6	-2	1	6	0	-5	3
2	0	2	0	0	-15	-5	6	-1	1	0	0	10	-10
2	1	2	8	0	10	-2	6	0	1	8	0	-20	24
2	2	2	2	0	0	2	6	1	1	0	0	10	-10
1	-1	3	6	-6	4	-2	6	2	1	6	0	-5	3
1	0	3	-2	2	2	-6	5	-2	2	6	8	5	5
1	1	3	6	-6	4	-2	5	-1	2	-4	-12	15	-35
0	0	4	1	3	0	0	5	0	2	6	18	-15	35
5	0	0	3	1	5	-5	5	1	2	-4	-12	15	-35
4	-1	1	0	-8	-2	10	5	2	2	6	8	5	5
4	0	1	0	6	-11	-5	4	-3	3	-4	-4	-1	-3
4	1	1	0	-8	-2	10	4	-2	3	6	6	24	12
3	-2	2	0	6	3	-5	4	-1	3	-12	-12	-43	-49
3	-1	2	0	0	-10	10	4	0	3	10	10	70	50
3	0	2	0	8	24	-20	4	1	3	-12	-12	-43	-49
3	1	2	0	0	-10	10	4	2	3	6	6	24	12
3	2	2	0	6	3	-5	4	3	3	-4	-4	-1	-3
2	-2	3	0	-6	3	-5	3	-3	4	-4	4	-3	-1
2	-1	3	0	0	-10	10	3	-2	4	6	-6	12	24
2	0	3	0	-8	24	-20	3	-1	4	-12	12	-49	-43
2	1	3	0	0	-10	10	3	0	4	10	-10	50	70
2	2	3	0	-6	3	-5	3	1	4	-12	12	-49	-43
1	-1	4	0	8	-2	10	3	2	4	6	-6	12	24
1	0	4	0	-6	-11	-5	3	3	4	-4	4	-3	-1
1	1	4	0	8	-2	10	2	-2	5	6	-8	-1	11
0	0	5	3	-1	0	0	2	-1	5	-4	12	9	-29
6	0	0	6	2	2	-2	2	0	5	6	-18	-21	41
5	-2	1	3	-1	0	0	2	1	5	-4	12	9	-29
5	-1	1	-2	4	-15	10	2	2	5	6	-8	-1	11
5	0	1	8	-6	5	5	1	-2	6	6	0	-5	3
5	1	1	-2	4	-15	10	1	-1	6	0	0	10	-10
5	2	1	3	-1	0	0	1	0	6	8	0	-20	24
4	-2	2	6	-6	-4	-8	10	1	1	6	0	10	-10
4	-1	2	-8	8	12	4	1	2	6	6	0	-5	3
4	0	2	14	-14	-1	-7	0	0	7	4	2	0	0

Fig. 1 Fourier coefficients of $\left(\begin{smallmatrix} a & b/2 \\ b/2 & c \end{smallmatrix}\right)$ in the basic forms $f_1, f_2, g_1, g_2, a + c \leq 7$

$$\begin{aligned}
 J^2 = & f_1^8 f_2^4 g_1^4 + 22f_1^7 f_2^3 g_1^4 + 119f_1^6 f_2^4 g_1^4 - 22f_1^5 f_2^5 g_1^4 + f_1^4 f_2^6 g_1^4 + 2f_1^8 f_2^2 g_1^3 g_2 + 46f_1^7 f_2^3 g_1^3 g_2 + 282f_1^6 f_2^4 g_1^3 g_2 \\
 & + 194f_1^5 f_2^5 g_1^3 g_2 - 42f_1^4 f_2^6 g_1^3 g_2 + 2f_1^3 f_2^7 g_1^3 g_2 + f_1^8 f_2^2 g_1^2 g_2^2 + 22f_1^7 f_2^3 g_1^2 g_2^2 + 120f_1^6 f_2^4 g_1^2 g_2^2 + 120f_1^4 f_2^6 g_1^2 g_2^2 \\
 & - 22f_1^3 f_2^7 g_1^2 g_2^2 + f_1^2 f_2^8 g_1^2 g_2^2 - 2f_1^7 f_2^3 g_1 g_2^3 - 42f_1^6 f_2^4 g_1 g_2^3 - 194f_1^5 f_2^5 g_1 g_2^3 + 282f_1^4 f_2^6 g_1 g_2^3 - 46f_1^3 f_2^7 g_1 g_2^3 \\
 & + 2f_1^2 f_2^8 g_1 g_2^3 + f_1^4 f_2^4 g_2^4 + 22f_1^5 f_2^5 g_2^4 + 119f_1^6 f_2^6 g_2^4 - 22f_1^3 f_2^7 g_2^4 + f_1^2 f_2^8 g_2^4 - 2f_1^7 f_2^3 g_2^5 - 110f_1^6 f_2^4 g_2^5 \\
 & - 166f_1^5 f_2^5 g_2^5 - 12f_1^4 f_2^6 g_2^5 - 4f_1^7 f_2^3 g_2^4 g_2 - 242f_1^6 f_2^4 g_2^4 g_2 - 450f_1^5 f_2^5 g_2^4 g_2 - 438f_1^4 f_2^6 g_2^4 g_2 - 24f_1^3 f_2^7 g_2^4 g_2 \\
 & - 2f_1^2 f_2^8 g_2^4 g_2 - 142f_1^6 f_2^4 g_1^3 g_2^2 + 62f_1^5 f_2^5 g_1^3 g_2^2 + 110f_1^4 f_2^6 g_1^3 g_2^2 - 370f_1^3 f_2^7 g_1^3 g_2^2 - 10f_1^2 f_2^8 g_1^3 g_2^2 - 10f_1^6 f_2^4 g_1^2 g_2^3 \\
 & + 370f_1^5 f_2^5 g_1^2 g_2^3 + 110f_1^4 f_2^6 g_1^2 g_2^3 - 62f_1^3 f_2^7 g_1^2 g_2^3 - 142f_1^2 f_2^8 g_1^2 g_2^3 + 2f_1^7 f_2^3 g_1^2 g_2^3 + 24f_1^6 f_2^4 g_1^2 g_2^3 - 438f_1^4 f_2^6 g_1^2 g_2^3 \\
 & + 450f_1^3 f_2^7 g_1^2 g_2^3 - 242f_1^2 f_2^8 g_1^2 g_2^3 + 4f_1^7 f_2^3 g_1 g_2^4 - 12f_1^6 f_2^4 g_1 g_2^4 + 166f_1^5 f_2^5 g_1 g_2^4 - 110f_1^4 f_2^6 g_1 g_2^4 + 2f_1^7 f_2^3 g_1 g_2^4 + f_1^6 g_1^6 \\
 & + 152f_1^5 f_2^5 g_1 g_2^4 + 48f_1^4 f_2^6 g_1 g_2^4 + 2f_1^6 g_1^5 g_2 + 340f_1^5 f_2^5 g_1^5 g_2 + 300f_1^4 f_2^6 g_1^5 g_2 + 96f_1^3 f_2^7 g_1^5 g_2 + f_1^6 g_1^4 g_2^2 + 212f_1^5 f_2^4 g_1^4 g_2^2 \\
 & - 312f_1^4 f_2^5 g_1^4 g_2^2 - 394f_1^3 f_2^6 g_1^4 g_2^2 + 24f_1^2 f_2^7 g_1^4 g_2^2 + 24f_1^5 f_2^3 g_1^3 g_2^3 - 540f_1^4 f_2^4 g_1^3 g_2^3 - 540f_1^3 f_2^5 g_1^3 g_2^3 - 24f_1^2 f_2^6 g_1^3 g_2^3 \\
 & + 24f_1^4 f_2^2 g_1^2 g_2^4 + 394f_1^3 f_2^3 g_1^2 g_2^4 - 312f_1^2 f_2^4 g_1^2 g_2^4 - 212f_1 f_2^5 g_1^2 g_2^4 + f_2^2 g_1^2 g_2^4 - 96f_1^3 f_2^3 g_1 g_2^5 + 300f_1^2 f_2^4 g_1 g_2^5 \\
 & - 340f_1 f_2^5 g_1 g_2^5 + 2f_2^6 g_1 g_2^5 + 48f_1^2 f_2^4 g_2^6 - 152f_1 f_2^5 g_2^6 + f_2^6 g_2^6 - 64f_1^4 g_1^7 - 144f_1^4 g_1^6 g_2 - 128f_1^3 f_2^3 g_1^2 g_2^2 \\
 & - 92f_1^4 g_1^5 g_2^2 + 264f_1^3 f_2^3 g_1^5 g_2^2 + 32f_1^2 f_2^4 g_1^5 g_2^2 - 12f_1^4 g_1^4 g_2^3 + 296f_1^3 f_2^3 g_1^4 g_2^3 + 4f_1^2 f_2^4 g_1^4 g_2^3 + 96f_1 f_2^5 g_1^4 g_2^3 \\
 & - 96f_1^3 f_2^3 g_1^3 g_2^4 + 4f_1^2 f_2^4 g_1^3 g_2^4 - 296f_1 f_2^5 g_1^3 g_2^4 - 12f_2^4 g_1^2 g_2^4 + 32f_1^2 f_2^3 g_1^2 g_2^5 - 264f_1 f_2^4 g_1^2 g_2^5 - 92f_2^3 g_1^2 g_2^5 \\
 & + 128f_1 f_2^5 g_1 g_2^6 - 144f_2^4 g_1 g_2^6 - 64f_2^3 g_2^7 - 128f_2^2 g_1^6 g_2^2 - 80f_2^2 g_1^5 g_2^2 - 128f_1 f_2^5 g_1^5 g_2^2 + 48f_1^2 g_1^4 g_2^4 + 48f_2^4 g_1^4 g_2^4 \\
 & + 128f_1 f_2^3 g_1^3 g_2^5 - 80f_2^3 g_1^3 g_2^5 - 128f_2^2 g_1^2 g_2^5 - 64g_1^5 g_2^4 - 64g_1^4 g_2^5.
 \end{aligned}$$

Fig. 2 Expression for J^2 in terms of the basic forms f_1, f_2, g_1, g_2

References

1. Aoki, H., Ibukiyama, T.: Simple graded rings of Siegel modular forms, differential operators and Borcherds products. *Int. J. Math.* **16**(3), 249–279 (2005)
2. Böcherer, S., Ibukiyama, T.: Surjectivity of Siegel Φ -operator for square free level and small weight. *Ann. Inst. Fourier (Grenoble)* **62**(1), 121–144 (2012)
3. Borcherds, R.: Automorphic forms with singularities on Grassmannians. *Invent. Math.* **132**(3), 491–562 (1998)
4. Grayson, D.R., Stillman, M.E.: Macaulay2, a software system for research in algebraic geometry. <http://www.math.uiuc.edu/Macaulay2/>
5. Gritsenko, V. Fourier–Jacobi functions in n variables. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **168** (Anal. Teor. Chisel i Teor. Funktsii. 9), 32–44, 187–188 (1988)
6. Hashimoto, K.: The dimension of the spaces of cusp forms on Siegel upper half-plane of degree two. *I. J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **30**(2), 403–488 (1983)
7. Hauffe-Waschbüsch, A., Krieg, A.: Congruence subgroups and orthogonal groups. *Linear Algebra Appl.* **618**, 22–36 (2021)
8. Igusa, J.: On Siegel modular forms of genus two. *Am. J. Math.* **84**, 175–200 (1962)
9. Looijenga, E.: Compactifications defined by arrangements. II. Locally symmetric varieties of type IV. *Duke Math. J.* **119**(3), 527–588 (2003)
10. Poor, C., Yuen, D.: Dimensions of cusp forms for $\Gamma_0(p)$ in degree two and small weights. *Abh. Math. Sem. Univ. Hamburg* **77**, 59–80 (2007)
11. Vignéras, M.-F.: Séries θ eta des formes quadratiques indéfinies. In: *Séminaire Delange–Pisot–Poitou*, 17e année (1975/1976). *Théorie des nombres: Fasc. 1, Exp. No. 20, 3.* (1977)
12. Wang, H.: The classification of free algebras of orthogonal modular forms. *Compos. Math.* **157**(9), 2026–2045 (2021)
13. Williams, B.: Poincaré square series for the Weil representation. *Ramanujan J.* **47**(3), 605–650 (2018)
14. Williams, B.: Remarks on the theta decomposition of vector-valued Jacobi forms. *J. Number Theory* **197**, 250–267. Erratum. *J. Number Theory* **201**(473–474), 2019 (2019)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.