

Erratum to: Clinical characteristics and outcomes of HIV-seropositive men treated with surgery for prostate cancer


Sudeh Izadmehr^{1,2} · Michael Leapman^{1,2} · Adele R. Hobbs^{1,2,3,4} · Maria Katsigeorgis⁴ · Fatima Nabizada-Pace^{1,2} · Seyed Behzad Jazayeri⁴ · David B. Samadi⁴

Published online: 25 August 2016
© Springer Science+Business Media Dordrecht 2016

Erratum to: Int Urol Nephrol DOI 10.1007/s11255-016-1338-4

In the original published version, Figure 2A was incorrectly published. The correct Fig. 2A is given here.

Fig. 2 Changes in Gleason score. Percentage of RALP patients that were upgraded or downgraded between biopsy and tumor removed by RALP (a). Change in Gleason score of HIV-positive prostate cancer before and after RALP (b)



The online version of the original article can be found under doi:[10.1007/s11255-016-1338-4](https://doi.org/10.1007/s11255-016-1338-4).

✉ David B. Samadi
robotmd@yahoo.com

¹ Department of Urology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

² University of California San Francisco School of Medicine, San Francisco, CA, USA

³ Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA, USA

⁴ Department of Urology, Hofstra Northwell School of Medicine, Lenox Hill Hospital, 485 Madison Avenue, 21 Floor, New York, NY 10022, USA