

ERRATUM

Erratum to: Gaseous Elemental Mercury Level and Distribution in a Heavily Contaminated Site: the Ex-chlor Alkali Plant in Torviscosa (Northern Italy)

**Alessandro Acquavita · Stefano Biasiol · Daniel Lizzi ·
Giorgio Mattassi · Mariangela Pasquon ·
Nicola Skert · Luca Marchiol**

Published online: 29 March 2017
© Springer International Publishing Switzerland 2017

Erratum to: Water Air Soil Pollut
DOI 10.1007/s11270-016-3234-z

The original publication contained the following errors.
(1) Table 3 should have been presented as shown here.
(2) In page 9, section “3.2 Biomonitoring Data”, 3rd paragraph's last sentence should be changed to: It should be emphasized that bioaccumulation recorded in the CAP area is among the highest ever recorded in autochthonous material in the Friuli Venezia Giulia region (Nimis et al. 1999; Tretiach and Baruffo 2001; Tretiach and Pittao 2008; Tretiach et al. 2011; Acquavita et al. 2015a, b).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11270-016-3234-z>

A. Acquavita (✉) · S. Biasiol · G. Mattassi · M. Pasquon ·
N. Skert
ARPA FVG, Via Cairoli 14, 33057 Palmanova, Italy
e-mail: alessandro.acquavita@arpa.fvg.it

D. Lizzi · L. Marchiol
Dipartimento di Scienze AgroAlimentari, Ambientali e Animali,
Università di Udine, Via delle Scienze 91, 33100 Udine, Italy

Table 3 Total Hg levels ($\mu\text{g g}^{-1}$) observed in biomonitoring with autochthonous lichens conducted in Italy and at Flix (NE Spain) (* data by tab. 12 in Supplementary material 10661_2010_1553_MOESM1_ESM.doc, Tretiach et al. 2011)

Site	Lichen species	Hg concentration (maximum level)	References
* Mt. Amiata	<i>P. sulcata</i> Taylor	3.93 ± 1.26	Bargagli et al. 1987
* Abbadia S. Salvatore, Mt. Amiata	<i>Parmelia saxatilis</i> (L.) Ach.	2.83 ± 0.65	Bargagli et al. 1987
* Abbadia S. Salvatore, Mt. Amiata	<i>Flavoparmelia caperata</i> (L.) Hale	2.43 ± 0.09	Bargagli et al. 1987
* Abbadia S. Salvatore, Mt. Amiata	<i>P. sulcata</i> Taylor	2.26 ± 1.71 (7.62)	Bargagli et al. 1989
* Bagnore, Mt. Amiata	<i>P. sulcata</i> Taylor	2.54 ± 1.27 (5.68)	Bargagli and Barghigiani 1991
* Bagnore, Mt. Amiata	<i>Xanthoria parietina</i> (L.) Th. Fr.	2.87 ± 0.66 (3.21)	Bargagli and Barghigiani 1991
* Isola di Vulcano	<i>Xanthoria parietina</i> (L.) Th. Fr.	0.23 ± 0.05 (0.27)	Bargagli and Barghigiani 1991
* Mt. Etna	<i>P. sulcata</i> Taylor	0.19 ± 0.04 (0.25)	Bargagli and Barghigiani 1991
* Isola di Vulcano	<i>Parmelia pastillifera</i> (Harm.) Hale	$0.22 \ddagger$	Barghigiani et al. 1990
* Mt. Etna	<i>P. sulcata</i> Taylor	0.18 ± 0.02	Bargagli et al. 1989
* Mt. Etna	<i>Pleurosticta acetabulum</i> (Neck.) Elix & Lumbsch	0.18 ± 0.03 (0.21)	Bargagli and Barghigiani 1991
* Isola di Vulcano	<i>Flavoparmelia caperata</i> (L.) Hale	$0.21 \pm 0.01 \ddagger$	Barghigiani et al. 1990
* Isola di Vulcano	<i>P. sulcata</i> Taylor	$0.21 \pm 0.01 \ddagger$	Barghigiani et al. 1990
* Mt. Etna	<i>Parmelia pastillifera</i> (Harm.) Hale	0.17 ± 0.02 (0.20)	Bargagli and Barghigiani 1991
* Mt. Etna	<i>Ramalina farinacea</i> (L.) Ach.	0.17 ± 0.02 (0.20)	Bargagli and Barghigiani 1991
* Lardereello	<i>Xanthoria parietina</i> (L.) Th. Fr.	2.16 ± 0.19 (2.41)	Bargagli and Barghigiani 1991
* Mt. Amiata	<i>Xanthoria parietina</i> (L.) Th. Fr.	1.18 ± 0.70 (1.96)	Loppi et al. 2006
* Acquapassante, Mt. Amiata	<i>Hypogymnia physodes</i> (L.) Nyl.	0.75 ± 0.25 (1.40)	Tretiach and Bargagli unpubl.
* Piancastagnato, Mt. Amiata	<i>P. sulcata</i> Taylor	0.49 ± 0.25 (0.93)	Loppi et al. 1999
* Travale-Radicondoli, Siena	<i>Flavoparmelia caperata</i> (L.) Hale	0.30 ± 0.16	Loppi et al. 1997
* Travale-Radicondoli, Siena	<i>Flavoparmelia caperata</i> (L.) Hale	0.20 ± 0.11 (0.56)	Loppi and Bargagli 1996
* Rosignano Solvay, Prov. Livorno	<i>Xanthoria parietina</i> (L.) Th. Fr.	0.74 ± 0.46 (1.84)	Bargagli et al. 1985
* Poggibonsi	<i>Flavoparmelia caperata</i> (L.) Hale	0.13 ± 0.03 (0.18)	Loppi et al. 2000
* Spilimbergo	<i>Xanthoria parietina</i> (L.) Th. Fr.	0.17 ± 0.08 (0.38)	Tretiach and Pittao 2008
* Lestans	<i>Punctelia borrieri</i> (Sm.) Krong	0.19 ± 0.07 (0.36)	Tretiach and Baruffo unpubl.
* Maniago	<i>Punctelia borrieri</i> (Sm.) Krong	0.06 ± 0.03 (0.13)	Tretiach and Baruffo unpubl.
* 20 km SO di Siena	<i>Flavoparmelia caperata</i> (L.) Hale	0.51 ± 0.23 (1.00)	Loppi et al. 1997
* N Italia	<i>Pseudevernia furfuracea</i> (L.)	$0.44 \pm 0.11 \ddagger$	Bergamaschi et al. 2007
* N Italia	<i>P. sulcata</i> Taylor	$0.20 \pm 0.05 \ddagger$	Bergamaschi et al. 2007

Table 3 (continued)

	Site	Lichen species	Hg concentration (maximum level)	References
*	N Italia	<i>Usnea gr. hirta</i>	0.28 ± 0.06 ‡	Bergamaschi et al. 2007
*	C Toscana	<i>Flavoparmelia caperata (L.) Hale</i>	0.17 ± 0.11	Loppi et al. 1997b
*	N Italia	<i>Hypogymnia physodes (L.) Nyl.</i>	0.10 ± 0.02 ‡	Bergamaschi et al. 2007
	Flix, NE Spain	<i>Xanthoria parietina (L.) Th. Fr.</i>	1.61 (0.39–3.79)	Esbrí et al. 2015
	Basso Isonzo, Laguna Marano e Grado	<i>Xanthoria parietina (L.) Th. Fr.</i>	0.12 ± 0.09 (0.39)	RAFG 2013
	Basovizza	<i>Xanthoria parietina (L.) Th. Fr.</i>	0.10 †	This study
	Torviscosa	<i>Xanthoria parietina (L.) Th. Fr.</i>	0.20 ± 0.08 (0.38)	This study