## **CORRECTION**



## Correction to: Hg and Se composition in demersal deep-sea fish from the North-East Atlantic

Guilherme Teixeira <sup>1</sup> · Joana Raimundo <sup>2,3</sup> · Joana Goulart <sup>4,5</sup> · Valentina Costa <sup>4,5</sup> · Gui M. Menezes <sup>5</sup> · Miguel Caetano <sup>2</sup> · Mário Pacheco <sup>1</sup> · Inês Martins <sup>4,5</sup>

Published online: 6 July 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

## Correction to: Environmental Science and Pollution Research https://doi.org/10.1007/s11356-020-08970-3

The correct Table 2 is presented in this paper. The original article is corrected.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Table 2** Arithmetic means  $\pm$  SD, total length, total Hg and Se concentrations ( $\mu$ gg<sup>-1</sup> dw and ww), Spearman correlations coefficient ( $r_s$ ) for total Hg and Se concentrations with length, Se:Hg molar ratios,

for the two tissues analysed. Statistically significant differences between the different species in muscle are indicated by \* and in liver by # (p < 0.05)

Species (Common name)	N	Tlength (range)	Tissue	THg dw	(μgg <sup>-1</sup> ) ww	THg: length (r <sub>s</sub> )	TSe dw	(μgg <sup>-1</sup> ) ww	TSe: length (r <sub>s</sub> )	Se:Hg molar ratio (range)	HBV- Se
Mora moro	30	55.05±3.36	M	$4.2 \pm 0.4$	$0.9 \pm 0.4$	0.30	$2.4 \pm 0.2$	$0.5 \pm 0.1$	0.1	$2.0 \pm 1.4$ (0.05-5.2)	-2.95
(Common mora)		(31-64)	L	$4.2^{\#} \pm 0.9$	$2.4\pm2.7$	0.14	$5.7^* \pm 0.9$	$3.1 \pm 2.8$	0.22	$10.5 \pm 13.6$ $(0.4-48.9)$	-2.96
Deania calcea	21	89.71±3.52	M	$8.8^* \pm 1.4$	$1.8 \pm 0.7$	0.21	$1.2 \pm 0.1$	$0.3 \pm 0.1$	-0.04	$0.4 \pm 0.15$ (0.2-0.7)	-14.56
(Birdbeak dogfish)		(82-106)	L	$2.7 \pm 1.5$	$1.5 \pm 1.5$	0.02	$2.7 \pm 0.6$	$1.5\pm0.8$	0.06	$6.5 \pm 7.9$ (0.6-30.1)	-2.38
Etmopterus pusillus	10	31.95 ±2.11	M	$3.2^* \pm 0.6$	$0.7\pm0.3$	0.48	$4.1 \pm 0.7$	$0.8 \pm 0.3$	0.82*	$3.5 \pm 1$ (1.6-4.7)	0.27
(Smooth lanternshark)		(26-40)	L	$0.32^{\#} \pm 0.1$	$0.2 \pm 0.1$	0.55	$4.1^* \pm 1.2$	$2.3 \pm 1.5$	0.12	$33.8 \pm 23.3$ (10.2-80.4)	2.25

The online version of the original article can be found at https://doi.org/10.1007/s11356-020-08970-3

- ☐ Inês Martins ines.ca.martins@uac.pt
- Department of Biology and CESAM, University of Aveiro, 3810-193 Aveiro, Portugal
- <sup>2</sup> IPMA- Portuguese Institute for Sea and Atmosphere, Av. Brasília, 1449-006 Lisbon, Portugal
- <sup>3</sup> Interdisciplinary Centre of Marine and Environmental Research (CIIMAR/CIMAR), University of Porto (U.Porto), Matosinhos, Portugal
- <sup>4</sup> IMAR- Department of Oceanography and Fisheries, University of Azores, 9901-862 Horta, Portugal
- OKEANOS- Research Unit- Faculty of Science and Technology, University of the Azores, 9901-862 Horta, Portugal

