



RETRACTION NOTE

Retraction Note to: Honokiol suppresses the development of post-ischemic glucose intolerance and neuronal damage in mice

Shinichi Harada¹ · Maya Kishimoto¹ · Mana Kobayashi¹ · Kazuo Nakamoto¹ · Wakako Fujita-Hamabe¹ · Hwei-Hsien Chen² · Ming-Huan Chan² · Shogo Tokuyama¹

Published online: 3 March 2020
© The Japanese Society of Pharmacognosy 2020

Retraction to: *J Nat Med* (2012) 66:591–599 <https://doi.org/10.1007/s11418-011-0623-x>

This article [1] has been retracted at the request of the corresponding author because an Investigation Committee established by Kobe Gakuin University (Kobe, Japan) has found numerous discrepancies between the raw data and the data presented in Figs. 6b, d. Statistical analysis of the raw data showed no significant difference between conditions. Authors S. Harada, K. Nakamoto, W. Fujita-Hamabe, H.-H. Chen, M.-H. Chan, and S. Tokuyama agree with this retraction. Authors M. Kishimoto and M. Kobayashi could not be reached for comment about this retraction.

Reference

1. Harada S, Kishimoto M, Kobayashi M et al (2012) Honokiol suppresses the development of post-ischemic glucose intolerance and neuronal damage in mice. *J Nat Med* 66:591. <https://doi.org/10.1007/s11418-011-0623-x>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s11418-011-0623-x>.

✉ Shogo Tokuyama
stoku@pharm.kobegakuin.ac.jp

¹ Department of Clinical Pharmacy, School of Pharmaceutical Sciences, Kobe Gakuin University, Hyogo, Japan

² Institute of Pharmacology and Toxicology, Tzu Chi University, Hualien, Taiwan