

MR-diffusion imaging in assessing chronic liver diseases: does a clinical role exist?

Imaging in diffusione RM nella valutazione delle epatopatie croniche: esiste un ruolo clinico?

**F. Pasquinelli¹ • G. Belli² • L.N. Mazzoni³ • F. Regini¹ • C. Nardi¹ • L. Grazioli⁴ • A. Linda Zignego⁵
S. Colagrande¹**

¹Careggi, Viale Morgagni 85, 50134 Florence, Italy

²Section of Medical Physics, Azienda Ospedaliero-Universitaria Careggi, Florence, Italy

³CIRM, University of Florence, Azienda Ospedaliero-Universitaria Careggi, Florence, Italy

⁴Department of Diagnostic for Imaging, 1st Radiology, Spedali Civili of Brescia, Italy

⁵Center for Systemic Manifestations of Hepatitis Viruses (MaSVE), Department of Internal Medicine, University of Florence, Florence, Italy

Correspondence to: S. Colagrande, Tel.: +39-055-7947189, Fax: +39-055-431970, e-mail: stefano.colagrande@unifi .it

Received: 23 February 2011 / Accepted: 23 March 2011 / Published online: 21 October 2011

© Springer-Verlag 2011

**Erratum to: La radiol med
DOI 10.1007/s11547-011-0730-5**

The original version of this article, unfortunately, contained a mistake. The name of the seventh author should have been A.L. Zignego instead of A. Linda Zignego

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s11547-011-0730-5>.

Published online: 16 December 2011