CORRECTION



Correction to: Computed tomography surveillance helps tracking COVID-19 outbreak

Akihiro Machitori¹ · Tomoyuki Noguchi^{2,3,4} · Yusuke Kawata¹ · Nobuhiko Horioka⁵ · Akihiro Nishie⁶ · Daisuke Kakihara⁶ · Kousei Ishigami⁶ · Shigeki Aoki⁷ · Yutaka Imai⁸

Published online: 27 August 2020 © Japan Radiological Society 2020

Correction to: Japanese Journal of Radiology https://doi.org/10.1007/s11604-020-01026-z

The authors would like to correct Abstract and text of original publication as follows:

Correction in Abstract Original publication:

Their PCR test results were positive (n = 62.5-398%), negative (n = 8.9-57%), unknown (n = 26.2-167%), and other disease (n = 2.4-15%).

The original article can be found online at https://doi.org/10.1007/ $s11604\hbox{-}020\hbox{-}01026\hbox{-}z.$

☐ Tomoyuki Noguchi tnogucci@radiol.med.kyushu-u.ac.jp

Akihiro Machitori dmachitori@hospk.ncgm.go.jp

Yusuke Kawata dkawata@hospk.ncgm.go.jp

Nobuhiko Horioka horioka-nobuhiko@mhlw.go.jp

Akihiro Nishie anishie@radiol.med.kyushu-u.ac.jp

Daisuke Kakihara kakky@radiol.med.kyushu-u.ac.jp

Kousei Ishigami ishigami@radiol.med.kyushu-u.ac.jp

Shigeki Aoki saoki@juntendo.ac.jp

Yutaka Imai imaiy@is.icc.u-tokai.ac.jp

Corrected sentence:

Their PCR test results were positive (n = 398, 62.5%), negative (n = 57, 8.9%), unknown (n = 167, 26.2%), and other disease (n = 15, 2.4%).

Correction in text (in Results section) Original publication:

The most common CT findings judged by radiologists were bilateral lung (male: female = 78.2-82.4%) and distributed GGOs (male: female = 80.8-83.0%). The crazy-paving pattern (male: female = 41.0-43.1%) and consolidation

- Department of Radiology, National Center for Global Health and Medicine, Kohnodai Hospital, 1-7-1 Kohnodai, Ichikawa City, Chiba Province 272-8516, Japan
- Department of Clinical Research, Center for Clinical Sciences, National Center for Global Health and Medicine, 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan
- Department of Radiology, National Hospital Organization Kyushu Medical Center, 1-8-1 Jigyohama, Chuo-ku, Fukuoka City, Fukuoka Province, Japan
- Department of Clinical Research, National Hospital Organization Kyushu Medical Center, 1-8-1 Jigyohama, Chuo-ku, Fukuoka City, Fukuoka Province 810-8563, Japan
- General Affairs Division, Health Policy Bureau, Ministry of Health, Labour and Welfare, 1-2-2 Kasumigaseki, Chiyoda-ku, Tokyo 100-8916, Japan
- Department of Clinical Radiology, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka City, Fukuoka Province 812-8582, Japan
- Department of Radiology, Juntendo University, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan
- Department of Radiology, Tokai University Hachioji Hospital, 1838 Ishikawa-cho, Hachioji City, Tokyo 192-0032, Japan



(male: female = 33.2–35.0%) were observed at about one-half the frequency of GGOs (Fig. 4).

Corrected sentence:

The most common CT findings judged by radiologists were bilateral lung (male: female = 78.2% : 82.4%) and distributed GGOs (male: female = 80.8% : 83.0%). The crazy-paving pattern (male: female = 41.0% : 43.1%) and consolidation (male: female = 33.2% : 35.0%) were observed at about one-half the frequency of GGOs (Fig. 4).

Correction in figure legend of Fig. 5 Original publication:

Significant differences between the PCR-positive and -negative patients in CT-surveillance were observed in the bilateral lung involvement (70.2–84.2%) and the

consolidation (33.9–50.9%). In the meantime, GGOs (83.5–83.9%) and the crazy-paving pattern (36.8–44.5%) were commonly observed

Corrected sentence:

Significant differences between the PCR-positive and -negative patients in CT-surveillance were observed in the bilateral lung involvement (70.2%, 84.2%) and the consolidation (33.9%, 50.9%). In the meantime, GGOs (83.5%, 83.9%) and the crazy-paving pattern (36.8%, 44.5%) were commonly observed

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

