

Recovery, Dependence or Death After Discharge

Deborah E. Barnes, PhD, MPH^{1,3}, W. John Boscardin, PhD^{1,3}, and C. Seth Landefeld, MD²

¹University of California, San Francisco, CA, USA; ²University of Alabama, Birmingham, AL, USA; ³San Francisco Veterans Affairs Medical Center, San Francisco, CA, USA.

J Gen Intern Med 28(3):343
DOI: 10.1007/s11606-012-2310-3
© Society of General Internal Medicine 2013

Authors' Reply—Thank you for your comments¹ related to our article on prediction of recovery, dependence or death in elders who become disabled during hospitalization.² We agree that validation of prognostic models such as the one we developed is critical. There are two key reasons for validation.³ The first is to account for the ‘optimism’ that occurs when a model is initially developed, since it is ‘optimized’ to the data. This can be thought of as ‘internal’ validation, and strategies such as dividing the sample into development and validation cohorts or bootstrapping techniques are utilized to estimate the extent to which c statistics may be inflated due to optimism.⁴ We utilized bootstrapping techniques, and the optimism-adjusted c statistics are reported at the end of the third paragraph of the results section (recovery: 0.76; dependence: 0.66; death: 0.73). The second reason for validation is to determine whether the model developed is predictive in other study populations and can be thought of

as ‘external’ validation.³ We acknowledged lack of external validation of our tool as a limitation. Since our article was published, we have been contacted by several other investigators with data that may allow us to take this important next step.

Corresponding Author: Deborah E. Barnes, PhD, MPH; University of California, San Francisco, CA, USA (e-mail: Deborah.barnes@ucsf.edu).

REFERENCES

1. **Sabour S.** Recovery, dependence or death after discharge. J Gen Intern Med. 2013; doi:10.1007/s11606-012-2311-2.
2. **Barnes DE, Mehta KM, Boscardin WJ, Fortinsky RH, Palmer RM, Kirby KA, Landefeld CS.** Prediction of recovery, dependence or death in elders who become disabled during hospitalization. J Gen Intern Med. 2012; doi:10.1007/s11606-012-2226-y.
3. **Justice AC, Covinsky KE, Berlin JA.** Assessing the generalizability of prognostic information. Ann Intern Med. 1999;130(6):515–24.
4. **Harrell FE Jr, Lee KL, Mark DB.** Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. Stat Med. 1996;15(4):361–87.