

Correction to: Comparative Effectiveness of Combining MTX with Biologic Drug Therapy Versus Either MTX or Biologics Alone for Early Rheumatoid Arthritis in Adults: a Systematic Review and Network Meta-analysis



*Katrina E. Donahue, M.D., M.P.H.^{1,2}, Elizabeth R. Schulman, M.D.³,
Gerald Gartlehner, M.D., M.P.H.^{4,5}, Beth L. Jonas, M.D., F.A.C.R.⁶,
Emmanuel Coker-Schwimmer, M.P.H.², Sheila V. Patel, B.S.P.H.⁴,
Rachel Palmieri Weber, Ph.D.², Carla M. Bann, Ph.D.⁴, and Meera Viswanathan, Ph.D.⁴*

¹Department of Family Medicine, University of North Carolina, Chapel Hill, NC, USA; ²Cecil G. Sheps Center for Health Services Research, Chapel Hill, NC, USA; ³Hospital for Special Surgery, New York, NY, USA; ⁴RTI International, Research Triangle Park, Durham, NC, USA; ⁵Department for Evidence-based Medicine and Clinical Epidemiology, Danube University, Krems, Austria; ⁶Department of Medicine, Division of Rheumatology, Allergy, and Immunology, University of North Carolina, Chapel Hill, NC, USA.

J Gen Intern Med
DOI: 10.1007/s11606-020-05839-6
© Society of General Internal Medicine 2020

CORRECTION TO: J GEN INTERN MED
<https://doi.org/10.1007/s11606-019-05230-0>

This article has been amended to include open access. The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s11606-019-05230-0>
