LETTER TO THE EDITOR



Long-Term Results of Laparoscopic Sleeve Gastrectomy for Morbid Obesity: 5 to 8-Year Results

Martine Uittenbogaart book Arijan A. P. M. Luijten François M. H. van Dielen Wouter K. G. Leclercq

Published online: 29 March 2017

© Springer Science+Business Media New York 2017

With great interest we read the study of Gadiot and colleagues regarding the long-term results of laparoscopic sleeve gastrectomy (LSG) in the January 2017 edition of *Obesity Surgery* [1]. First, we would like to express our appreciation of this very nice report portraying long-term data on a sizeable cohort of patients with LSG. Second, we would like to ask a few questions concerning the study.

In Table 3, the authors report on the failure rates at respectively 5 to 8 years postoperatively. Regarding inadequate weight loss following bariatric surgery, it is favourable to divide between weight loss failure (WLF)—inability to achieve 50% excess weight loss—and weight regain (WR)—significant

Martine Uittenbogaart m.uittenbogaart@mmc.nl

Arijan A. P. M. Luijten a.luijten@mmc.nl

François M. H. van Dielen f.vandielen@mmc.nl

Wouter K. G. Leclercq w.leclercq@mmc.nl

Obesity Centre Máxima, Máxima Medical Centre, Veldhoven, the Netherlands weight gain after initial weight loss success. Is it possible to provide us with the WLF and WR percentages?

Furthermore, could the authors elaborate on the mentioned percentage of subsequent revision to laparoscopic Roux-en-Y gastric bypass (LRYGB), specifically whether these patients managed to return to adequate weight loss after revision?

Last, did the results in this study have specific implications on your patient selection nowadays, with regard to eligibility for either LSG or LRYGB?

Compliance with Ethical Standards

Ethical Approval This article does not contain any studies with human participants or animals performed by any of the authors.

Informed Consent Does not apply.

Funding No funding applicable.

Conflict of Interest The authors declare that they have no conflict of interest.

References

 Gadiot RP, Biter LU, van Mil S, et al. Long-term results of laparoscopic sleeve gastrectomy for morbid obesity: 5 to 8-year results. Obes Surg. 2017;27(1):59–63. doi:10.1007/s11695-016-2235-8.

