

Erratum to: Usefulness of combining admission brain natriuretic peptide (BNP) plus hospital discharge bioelectrical impedance vector analysis (BIVA) in predicting 90 days cardiovascular mortality in patients with acute heart failure

Simona Santarelli¹ · Veronica Russo¹ · Irene Lalle¹ · Benedetta De Berardinis¹ · Silvia Navarin¹ · Laura Magrini¹ · Antonio Piccoli² · Marta Codognotto² · Luigi Maria Castello³ · Gian Carlo Avanzi³ · Humberto Villacorta^{4,5} · Bernardo Luiz Campanário Precht⁴ · Pilar Barreto de Araújo Porto⁴ · Aline Sterque Villacorta⁴ · Salvatore Di Somma¹ · On Behalf of Great Network

Published online: 8 February 2017
© SIMI 2017

Erratum to: Intern Emerg Med DOI 10.1007/s11739-016-1581-9

In the original publication, the hydration status between survivors and non-survivors has been incorrectly mentioned in the abstract section.

The correct text should read as below:

At discharge, BIVA shows a statistically significant difference in hydration status between non survivors and survivors [respectively, hydration index (HI) 85 vs 74, $p < 0.001$; reactance (X_c) 26.7 vs 37, $p < 0.001$; resistance (R) 445 vs 503, $p < 0.01$].

The online version of the original article can be found under doi:[10.1007/s11739-016-1581-9](https://doi.org/10.1007/s11739-016-1581-9).

✉ Salvatore Di Somma
salvatore.disomma@uniroma1.it

¹ Department of Medical-Surgery Sciences and Translational Medicine, Emergency Medicine, Postgraduate School of Emergency Medicine, University La Sapienza, Via di Grottarossa 1035/1039, 00189 Rome, Italy

² Department of Medicine, University of Padova, Padua, Italy

³ Department of Translational Medicine, University of Piemonte Orientale, Novara, Italy

⁴ Department of Cardiology, Fluminense Federal University, Niteroi, RJ, Brazil

⁵ Hospital Unimed Rio, Rio de Janeiro, RJ, Brazil