

## Erratum to: Robotic-assisted TKA Reduces Postoperative Alignment Outliers and Improves Gap Balance Compared to Conventional TKA

Eun-Kyoo Song MD, PhD, Jong-Keun Seon MD, PhD,  
Ji-Hyeon Yim MD, PhD, Nathan A. Netravali PhD,  
William L. Bargar MD

Published online: 14 July 2012  
© The Association of Bone and Joint Surgeons® 2012

**Erratum to: Clin Orthop Relat Res**  
**DOI 10.1007/s11999-012-2407-3**

In the third sentence of the Patients and Methods section it states, “We considered patients with primary osteoarthritis of the knee and a mechanical axis between 20° and 5°

valgus”. However, the sentence should state, “We considered patients with primary osteoarthritis of the knee and a mechanical axis between 20° varus and 5° valgus”.

In addition, Dr. Yim’s degree is MD rather than MD, PhD. He currently is studying for his PhD degree.

---

The online version of the original article can be found under  
doi:[10.1007/s11999-012-2407-3](https://doi.org/10.1007/s11999-012-2407-3).

---

E.-K. Song, J.-K. Seon, J.-H. Yim  
Department of Orthopedic Surgery, Chonnam National  
University Hwasun Hospital, Chonnam, Korea

N. A. Netravali  
Curexo Technology Corp, Fremont, CA, USA

W. L. Bargar (✉)  
Department of Orthopaedics, University of California  
at Davis School of Medicine, Sutter General Hospital,  
1020 29th Street, #450, Sacramento, CA 95816, USA  
e-mail: [wbargar@jointsurgeons.com](mailto:wbargar@jointsurgeons.com)