

Erratum to: How Much Clinical and Functional Impairment do Children Treated With Knee Rotationplasty Experience in Adulthood?

Maria Grazia Benedetti MD, Yusuke Okita PT, PhD, Elena Recubini MD,
Elisabetta Mariani MD, Alberto Leardini PhD, Marco Manfrini MD

Published online: 2 February 2016
© The Association of Bone and Joint Surgeons® 2016

Erratum to: Clin Orthop Relat Res
DOI 10.1007/s11999-016-4691-9

In the study, “How Much Clinical and Functional Impairment do Children Treated With Knee Rotationplasty Experience in Adulthood?” the year ranges on the X axis are incorrect as published. The year ranges should be “1986–1990”, “1991–1995”, “1996–2000”, “2001–2005”, and “2006–2009”. Figure 1 has been corrected to reflect this change.

The authors apologize for the error.

The online version of the original article can be found under
doi:[10.1007/s11999-016-4691-9](https://doi.org/10.1007/s11999-016-4691-9).

M. G. Benedetti, Y. Okita (✉), E. Recubini, E. Mariani
Physical Medicine and Rehabilitation Unit, Rizzoli Orthopedic
Institute, Via Pupilli 1, 40136 Bologna, Italy
e-mail: okita.yusuke@gmail.com

A. Leardini
Movement Analysis Laboratory, Rizzoli Orthopedic Institute,
Bologna, Italy

M. Manfrini
Orthopaedic and Traumatologic Clinic for Musculoskeletal
Tumors, Rizzoli Orthopedic Institute, Bologna, Italy

Fig. 1 The trend in surgical indications at our institution for high-grade sarcomas of the distal femur in children younger than 14 years (1986–2009) is shown.

