

Short-Term Effects of Sibutramine on Mineral Status and Selected Biochemical Parameters in Obese Women

Joanna Suliburska · Paweł Bogdański ·
Monika Szulińska · Danuta Pupek-Musialik

Received: 10 February 2012 / Accepted: 11 April 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract The aim of this study was to assess the effect of sibutramine on mineral status and selected biochemical parameters in obese women. The study was conducted on 24 patients who received 15 mg daily doses of sibutramine for 12 weeks, and on 20 patients who received placebo. At the baseline, after the sixth and twelfth weeks of treatment, body weight and blood pressure were measured, the BMI was calculated, and samples of blood and of first morning urine were collected. Serum lipid profiles, glucose levels, and nitric oxide levels were determined. The iron (Fe), copper (Cu), zinc (Zn), calcium (Ca), and magnesium (Mg) present in the serum and urine samples were assessed. The erythrocyte hemolysate of the patients was used to assay the activity of glutathione peroxidase (GSH-Px) and superoxide dismutase (SOD). No changes were observed in BMI, blood pressure, or nitric oxide during the study. After 12 weeks of treatment, a decrease was observed in total cholesterol, LDL cholesterol, triglyceride, glucose, and ferritin levels. GSH-Px and SOD activity increased after 12 weeks of sibutramine treatment. The Mg and Cu increases were observed in serum after the sixth and twelfth weeks of treatment. It was found that the Zn level decreased in serum after the twelfth week. The elimination of Ca, Mg,

Fe, Zn, and Cu in urine also declined in the twelfth week. No differences were found in the women taking the placebo. In conclusion, we found that sibutramine had a positive effect on lipid and glucose status in obese women. However, the drug disturbed the balance of minerals, especially Zn and Mg, in the subjects.

Keywords Sibutramine · Minerals · Lipid profile · Glucose · Obesity · Women

Introduction

The prevalence of obesity and its comorbidities has been increasing all over the world. Abdominal or visceral obesity is closely related to disturbances in lipid and glucose status, and also to a lower antioxidative status in the body [1]. Weight loss reduces the cardiovascular risks associated with obesity [2]. The treatment of obesity should be individually tailored and should be maintained for a long term. The first line of strategy for weight loss is a combination of diet, physical activity, and behavior modification. Anti-obesity drugs may be used in adult patients when dietary and lifestyle modifications have been unsuccessful.

For a long time, sibutramine was used as an effective anti-obesity drug in Europe, but because of the cardiovascular risk associated with it, European regulators suspended its market authorization, and the US Food and Drug Administration restricted its license. This study was conducted a few months before the withdrawal of sibutramine's license. Over the years, the effectiveness and side effects of sibutramine have been assessed in many studies. Subsequent studies have shown that the drug has a significant effect on weight loss, due to its satietogenic and calorogenic effects [3]. It also improved glycemic and lipid profile and

J. Suliburska (✉)
Department of Human Nutrition and Hygiene,
Poznan University of Life Sciences,
Wojska Polskiego 31 Str.,
60-624 Poznan, Poland
e-mail: jsulibur@up.poznan.pl

P. Bogdański · M. Szulińska · D. Pupek-Musialik
Department of Internal Medicine, Metabolic Disorders
and Hypertension, Poznan University of Medical Sciences,
Szamarzewskiego 84 Str.,
60-569 Poznan, Poland

uric acid, and for this reason, it was often recommended in diet and lifestyle advice [4]. Svacina et al. [5] documented the positive effect of a 3-month sibutramine treatment on dyslipidemia in the obese, even in those already being treated with statins. In studies with hypertensive subjects whose blood pressure was adequately controlled by antihypertensive drugs, a slight to significant decrease in systolic and diastolic blood pressure was observed with sibutramine use [6]. Moreover, Derosa et al. [7] found that sibutramine plus L-carnitine gave a greater improvement in body weight loss and glycemic and lipid profile compared to sibutramine alone. Sibutramine therapy has been associated with significant C-reactive protein reduction as compared with routine treatment in patients with coronary artery disease. Thus, short-term (4 months) therapy with sibutramine, together with diet and lifestyle modifications, was associated with improved endothelial function [8] but did not affect TNF- α [9]. In our previous unpublished research, we observed a considerable decrease in ferritin and Zn concentration in the serum of patient undergoing treatment with sibutramine. A review of the literature on the topic showed that there was limited data on the effects of sibutramine on mineral balance in obese women. The aim of this study was thus to assess the effect of sibutramine on mineral status and on selected biochemical parameters in obese women.

Material and Methods

Study Design

The study was conducted at the Department of Internal Diseases, Metabolic Disorders, and Arterial Hypertension, Poznan University of Medical Sciences (Poland), and at the Department of Human Nutrition and Hygiene, Poznan University of Life Sciences (Poland). The study protocol was approved by Bioethics Commission at Poznan University of Medical Sciences (approval no. 86/09).

Patients

In the Clinic of the Department of Internal Diseases, Metabolic Disorders, and Hypertension in Poznan, we treated obese patients with sibutramine. During the biochemical studies carried out in women taking sibutramine, we observed a considerable decrease in ferritin and Zn concentration in serum. In addition, these women often complained about the brittleness of their nails and hair and of taste disorders, which may have been due to low concentrations of Zn in their bodies. This imbalance of mineral status, apparently due to the use of sibutramine, was disturbing, especially when it is known that an appropriate level of minerals, particularly of Ca and Fe, is very important for

the health of women of childbearing age. We thus decided to conduct a study accessing the effect of sibutramine on mineral status in obese women.

The study was conducted on 44 obese women (body mass index [BMI] ≥ 30 kg/m²) aged 40.6 \pm 12.1 years. Individuals suffering from chronic disease or using any medications (including OTC products, with the exception of five women who took oral contraceptives) were excluded. For at least 3 months before the study, all women have remained on a controlled-energy diet (with nearly 500 kcal daily deficit) that took 58 % of its calories from carbohydrates, 27 % from fat, and 15 % from proteins, with a maximum cholesterol content of 300 mg/day and 35 g/day of fiber. Standard dietary advice was given by a qualified dietitian. Subjects did not change their diet during the study and were asked to restrain from smoking for the time of the study.

Women who were pregnant, breastfeeding, or capable of becoming pregnant were excluded. All subjects were informed about the study's purpose, protocol, and risks. All individuals provided informed written consent.

Treatment and Assessment

Twenty-four patients aged 40.6 \pm 12.1 were assigned to receive sibutramine 15 mg once daily for 12 weeks, while 20 patients aged 40.7 \pm 12.2 received placebos containing pure microcrystalline cellulose. Before the beginning of the study, all patients underwent a full physical examination. Anthropometric measurements of the subjects wearing light clothing and no shoes were conducted. Weight was measured to the nearest 0.1 kg, and height to the nearest 0.1 cm. The BMI was calculated by dividing the weight (kg) by the height squared (m²). Obesity was defined as BMI ≥ 30 kg/m². Blood pressure was measured according to the guidelines of the European Society of Hypertension using a digital electronic tensiometer (model 705IT, Omron Corporation, Kyoto, Japan). Regular or large adult cuffs were used, depending on the patient's arm circumference. The measurements were taken while fasting in the morning hours in a sitting position, with the legs uncrossed, the back and arms supported. Hypertension was defined by measurement of arterial blood pressure as the average of three measurements obtained after 10 min of physical resting (3 times at 2 different visits).

All participants had blood collected from a forearm vein in serum-separated tubes (without using an anticoagulant). The coagulated blood was left to clot at room temperature for 30 min, and then centrifuged for 15 min at 2,000 rpm at 4°C. Then the supernatant fluid was separated. Serum samples were stored at -20°C for no longer than 2–3 days. Blood samples were collected after an overnight fast, and after 30 min in the supine position.

Measurements of total cholesterol (TC), high density lipoproteins (HDL-C), low density lipoproteins (LDL-C), triglycerides (TG), and glucose were performed on each blood sample. Prior to the Fe, Cu, and Zn serum concentration assays, the serum was diluted with triton X-100 (Merck), and before the Ca and Mg concentration assays, the serum was diluted with triton and LaCl₃ (Merck).

Urine was collected in the morning, following an overnight fast. Morning urine samples were collected in sterilized vessels and stored for material analysis.

The samples of urine were mineralized in a mixture of concentrated nitric (65 %) and perchloric (60 %) acids (suprapure, Merck; v/v, 1:1) in a microwave oven (Milestone), and dissolved (v/v, 1:1) in Triton X-100 (Merck; all minerals) and LaCl₃ (Merck; Ca and Mg).

After the sixth and twelfth weeks of treatment, weight and blood pressure were measured, BMI was calculated, and the blood samples and first morning urine were collected from all patients. All serum and urine parameters were determined after a 12-h overnight fast.

Biochemical Assays

The content of Fe, Cu, Zn, Ca, and Mg in the serum and urine samples was determined by the flame atomic absorption spectrometry (AAS-3 spectrometer, Carl Zeiss, Germany with deuterium background correction). In order to obtain concentrations of the serum bioelements, the samples were diluted (v/v, 1:1) as follows: for Fe, Zn, and Cu analyses 0.01 % Triton X-100 (Merck) was used, while for the Mg and Ca analysis, aqueous solutions consisting of 0.01 % Triton X-100 (Merck) and 0.05 % lanthanum chloride (Merck) were used. The content of Fe, Cu, Zn, Ca, and Mg in serum and urine samples was determined at the following wavelengths: 248.3 nm (Fe), 324.8 nm (Cu), 213.9 nm (Zn), 422.7 nm (Ca), and 285.2 nm (Mg).

The accuracy of the method was verified using certified reference materials (HUM ASY CONTROL 2 and URN ASY CONTROL 2, Randox), and reached 95 %, 99 %, 94 %, 99 %, and 102 % for Ca, Mg, Fe, Zn, and Cu, respectively [10].

Plasma TC, LDL-C, HDL-C, TG, and glucose levels were measured using commercial kits. The accuracy and precision of the techniques used to assay the lipids and glucose were validated. Reproducibility was checked with a human serum control (Randox). Accuracy was assessed by means of the recovery value and its range between 95 % and 109 %, and the variability coefficient (an indicator of the method's precision) did not exceed 10 %.

The activity of glutathione peroxidase (GSH-Px) was assayed in the erythrocyte hemolysate using a method of Paglia and Valentine [11]. GSH-Px decomposes hydrogen

peroxide, causing oxidation of the reduced glutathione. Oxidized glutathione is then reduced in a reaction that is catalyzed by glutathione reductase. The coenzyme of the reaction is reduced nicotinamide adenine dinucleotide phosphate, which is transformed into oxidized form, and causes a change in light absorbance at 340 nm (spectrophotometer Sunrise, Tecan). Superoxide dismutase (SOD) activity was determined in the erythrocyte hemolysate using a modification of epinephrine-adenochrome detection system [12]. The enzyme inhibits the reaction of adrenaline autooxidation to adenochrome in an alkaline environment. The unit of SOD activity under the given conditions is defined as the amount of enzyme that inhibits 50 % of the reaction at the maximal increase of absorbance rectilinear segment of adenochrome formation at wavelength 450 nm (spectrophotometer Sunrise, Tecan). Erythrocyte GSH-Px and SOD activity was determined using a commercial kit (Randox), and expressed as units per gram of hemoglobin. The hemoglobin concentration in milligrams per milliliters was determined using the cyanmethemoglobin method [13].

Erythrocyte lysate was prepared from blood collected with EDTA. Blood was centrifuged at 700–1,000×g at 4°C. Plasma was then removed. The red cells were washed three times with cold isotonic saline solution. Erythrocytes were lysed with a ninefold volume of ice-cold HPLC-grade water, and then centrifuged at 10,000×g at 4°C. Supernatant was used for assaying.

The concentration of ferritin in the serum was determined by a two-site sandwich immunoassay using direct chemiluminometric technology (ADVIA Centaur XP analyzer), which uses constant amounts of two anti-ferritin antibodies [14].

The concentration of nitric oxide (NO) was determined using the spectrophotometric method on the serum, with the testing set by Oxis. The test determines nitric oxide concentration based on the enzymatic conversion of nitrate to nitrite by nitrate reductase. The reaction is followed by colorimetric detection of nitrate as an azo dye product of the Griess Reaction. The Griess reaction transforms nitrites to NO, which then reacts with sulfanilic acid, and the indirect product of the reaction interacts with N-(1-naphthyl) ethylenediamine, leading to the formation of a red-violet colored product having maximum absorbance at $\lambda = 540$ nm. The solution color change, measured spectrophotometrically (with Hyperion Micro Reader), is proportional to the nitric oxide concentration [15].

Statistical Analysis

The experimental results were given as means±standard deviation. The statistical analysis was carried out using STATISTICA 7.0 software and the ANOVA test. A *p* value less than 0.05 was considered statistically significant.

Results

Body Mass Index and Blood Pressure

The results of BMI and blood pressure measurements are presented in Table 1. No significant change was observed in BMI or in the systolic or diastolic blood pressure of patients during the study. However, the body mass index slightly decreased after 6 weeks (38.43 vs. 37.93, $p=0.345$) and 12 weeks (38.43 vs. 36.83, $p=0.204$). Both systolic and diastolic blood pressure did not change at all in comparison with the baseline.

Biochemical Parameters

After 6 weeks of sibutramine treatment, a significant decrease was found only in TC ($p=0.031$; Table 2). No change in LDL-C, TG, glucose, or ferritin levels was observed at 6 weeks. After 12 weeks, a significant decrease was observed for TC ($p=0.013$), LDL-C ($p=0.034$), TG ($p=0.044$), glucose ($p=0.046$), and ferritin ($p=0.030$). GSH-PX and SOD activity markedly increased ($p=0.036$ and $p=0.021$, respectively) after 12 weeks of sibutramine treatment. Serum levels of NO did not change during the study. In women taking the placebo, no significant difference was observed.

Minerals

A significant increase in Mg and Cu concentration was observed in the serum of women with sibutramine after 6 ($p=0.037$ and $p=0.029$, respectively) and 12 weeks ($p=0.030$ and $p=0.021$, respectively; Table 3). It was also found that Zn levels markedly decreased ($p=0.018$) in serum after 12 weeks, and a slight decrease was also observed after

6 weeks of treatment. In the case of Fe serum concentration, a slight decrease was found during the study.

The elimination of Ca, Mg, Fe, and Cu in urine declined significantly at 6 weeks ($p=0.032$, $p=0.048$, $p=0.037$, and $p=0.026$, respectively) and 12 weeks ($p=0.028$, $p=0.031$, $p=0.023$, and $p=0.018$, respectively) of the study (Table 3). The considerable decrease in Zn concentration in urine was observed only after 12 weeks ($p=0.017$), but a slight decline in Zn elimination was evident after 6 weeks of the study.

It was found that the Ca/Mg and Zn/Cu molar ratios in serum significantly decreased at the sixth ($p=0.042$ and $p=0.032$) and twelfth weeks ($p=0.034$ and $p=0.021$) of the study (Table 4). A marked Cu/Fe molar ratio increase was present at 6 ($p=0.035$) and 12 weeks ($p=0.028$) vs. the baseline, and at 12 weeks vs. 6 weeks ($p=0.046$). In urine, significant increases in the molar ratios of Ca/Mg and Zn/Cu, as well as a significant decrease in the Fe/Zn molar ratio at 6 ($p=0.016$, $p=0.027$, and $p=0.041$, respectively) and 12 weeks ($p=0.013$, $p=0.022$, and $p=0.037$, respectively) of sibutramine treatment were observed. No marked differences in women receiving the placebo were found.

Discussion

Sibutramine hydrochloride monohydrate is a norepinephrine and serotonin reuptake inhibitor approved for the long-term management of obesity, in conjunction with a reduced calorie diet and behavior modification in patients unable to lose weight by means of diet and lifestyle changes alone. The efficacy of sibutramine has been demonstrated in randomized trials in obese patients [16–18]. Furthermore, its favorable influence on the lipid profile and glucose concentration has been shown in several studies [19–21]. In the study of Tambascia et al. [22], sibutramine demonstrated efficacy in

Table 1 Baseline values

	Baseline		6 weeks		12 weeks	
	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)
N	24	20	–	–	–	–
Sex [F]	24	20	–	–	–	–
Age [years]	40.61±12.13	40.72±12.22	–	–	–	–
Sibutramine [dose]	15 mg	–	15 mg	–	15 mg	–
BMI [kg/m ²]	38.43±8.54	38.03±5.76	37.93±7.54	37.55±6.21	36.83±7.35	37.31±4.83
SBP [mmHg]	133.52±10.44	131.82±8.37	133.18±12.42	132.87±8.91	133.57±9.74	132.38±8.31
DBP [mmHg]	85.53±5.22	85.77±5.42	85.68±5.61	85.73±5.34	85.42±4.81	85.74±4.11

Blood pressure and BMI changes at 6 and 12 weeks in the group during the study. Data are means±SD

mean the arithmetic mean, SD standard deviation, N number of subjects, BMI body mass index, SBP systolic blood pressure, DBP diastolic blood pressure

Table 2 Biochemical parameters in serum changes at 6 and 12 weeks during the study

	Baseline		6 weeks		12 weeks	
	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)
TC [mmol/l]	4.41±0.58	4.40±0.51	4.11±0.57 ^a	4.38±0.56	4.08±0.58 ^a	4.35±0.52
LDL-C [mmol/l]	2.56±0.48	2.55±0.51	2.35±0.46	2.52±0.49	2.28±0.42 ^a	2.50±0.43
HDL-C [mmol/l]	1.25±0.24	1.28±0.21	1.26±0.25	1.28±0.23	1.25±0.26	1.26±0.21
TG [mmol/l]	1.37±0.37	1.35±0.38	1.21±0.36	1.30±0.30	1.15±0.35 ^a	1.30±0.31
Glucose [mmol/l]	5.35±0.47	5.31±0.48	5.33±0.46	5.30±0.47	5.18±0.27 ^a	5.30±0.50
Ferritin [µg/l]	91.12±40.42	91.25±41.22	78.15±41.39	88.11±40.08	65.51±38.50 ^a	85.14±30.83
Nitric oxide (NO) [µmol/l]	10.60±3.01	10.67±2.67	10.83±3.08	10.57±2.68	10.67±2.99	10.60±2.68
Glutathione peroxidase [U/gHb]	15.62±6.30	15.68±7.11	18.26±8.23	16.15±7.83	19.52±8.45 ^a	16.22±6.28
Superoxide dismutase [U/gHb]	2,570±692	2,582±5.82	2,853±576	2,637±583	3,554±723 ^a	2,628±683

Data are means±SD

mean the arithmetic mean, SD standard deviation, N number of subjects, TC total cholesterol, LDL-C LDL cholesterol, HDL-C HDL cholesterol, TG triglycerides

^aSibutramine group $p < 0.05$ vs baseline

reducing weight, insulin resistance, triglycerides, and uric acid, but not glucose or cholesterol, in non-diabetic women. The results of our study after 6 weeks show a significant decrease only for TC. No change in LDL-C, TG, glucose levels were observed at that time. After 12 weeks, a significant decrease in TC, LDL-C, TG, and glucose was observed.

Some authors suggest that the use of sibutramine could raise blood pressure on account of its peripheral sympathomimetic effect [23]. Contrarily, reports of other authors and the results of our own study show no significant impact of sibutramine on blood pressure or on NO concentration in long-term or short-term treatment [3, 24]. It is suggested that sibutramine elicits a central sympatholytic effect in the

brain, counteracting, at least in part, peripheral sympathetic stimulation.

To the best of our knowledge, this is the first clinical study investigating the effect of sibutramine on mineral status in obese women. The results of the experimental studies suggest the potential influence of sibutramine on minerals. Emer et al. [25] found that a relatively high dose of sibutramine resulted in significant increases in Fe and Zn levels in the kidneys and in the adrenal glands in rats. The results of our study show significant changes in mineral levels of obese women treated with sibutramine.

The mechanism may be related to reduced energy intake as an effect of sibutramine action. It has been demonstrated that lower food intake leads to a lower supply of minerals

Table 3 Minerals in serum and urine changes at 6 and 12 weeks during the study

	Baseline		6 weeks		12 weeks	
	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)
Serum						
Ca [mmol/l]	2.54±0.13	2.52±0.12	2.65±0.42	2.60±0.21	2.52±0.25	2.50±0.33
Mg [mmol/l]	0.80±0.07	0.83±0.07	0.93±0.06 ^a	0.85±0.06	0.95±0.06 ^a	0.88±0.25
Fe [µmol/l]	17.85±5.32	17.33±5.18	16.82±6.24	16.84±5.10	16.35±5.92	16.58±5.03
Zn [µmol/l]	12.42±2.42	12.04±2.80	11.48±2.38	11.96±2.81	10.51±2.58 ^a	11.46±2.91
Cu [µmol/l]	13.61±1.81	13.03±1.71	14.94±1.78 ^a	13.41±1.80	15.68±1.92 ^a	13.63±1.21
Urine						
Ca [mmol/l]	3.84±2.53	3.21±2.31	2.97±2.09 ^a	3.25±2.30	2.89±2.35 ^a	3.18±2.18
Mg [mmol/l]	5.84±2.42	5.50±2.80	3.25±1.78 ^a	5.00±1.80	3.05±1.50 ^a	4.58±2.91
Fe [µmol/l]	9.27±5.21	9.30±4.77	6.02±4.20 ^a	8.50±4.18	5.78±3.84 ^a	7.90±5.10
Zn [µmol/l]	9.38±3.58	9.40±3.37	8.46±4.21	8.98±4.80	7.78±3.80 ^a	9.12±2.83
Cu [µmol/l]	1.02±0.51	1.11±0.68	0.68±0.34 ^a	0.98±0.80	0.61±0.35 ^a	1.08±0.56

Data are means±SD

mean the arithmetic mean, SD standard deviation, N number of subjects

^aSibutramine group $p < 0.05$ vs baseline

Table 4 Mineral molar ratio in serum and urine changes at 6 and 12 weeks during the study

	Baseline		6 weeks		12 weeks	
	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)	Sibutramine (N=24)	Placebo (N=20)
Serum						
Ca/Mg	3.16±0.21	3.04±0.20	2.85±0.27 ^a	3.06±0.25	2.66±0.23 ^a	2.84±0.20
Fe/Zn	1.44±0.12	1.44 0±.13	1.47±0.12	1.41±0.14	1.56±0.12	1.45±0.10
Cu/Fe	0.76±0.09	0.75±0.10	0.89±0.05 ^a	0.80±0.04	0.96±0.05 ^{a,b}	0.82±0.07
Zn/Cu	0.92±0.11	0.92±0.12	0.77±0.12 ^a	0.89±0.18	0.66±0.12 ^a	0.86±0.09
Urine						
Ca/Mg	0.65±0.23	0.59±0.25	0.92±0.21 ^a	0.65±0.25	0.95±0.21 ^a	0.69±0.31
Fe/Zn	0.98±0.31 ^a	0.98±0.30	0.72±0.30 ^a	0.95±0.31	0.74±0.31 ^a	0.87±0.40
Cu/Fe	0.11±0.05	0.12±0.06	0.11±0.04	0.12±0.06	0.11±0.04	0.13±0.07
Zn/Cu	9.20±3.78 ^a	8.47±3.05	12.5±4.65 ^a	9.16±5.08	12.8±5.33 ^a	8.45±3.08

Data are means±SD
mean the arithmetic mean, *SD* standard deviation, *N* number of subjects

^aSibutramine group *p*<0.05 vs baseline

^bSibutramine group *p*<0.05 vs 6 weeks

and reduced mineral status in the body. Acuirre et al. [26] reported that a 1,000 kcal/day diet decreases the intake of Fe, Zn, Cu, and Ca by 50 %, 30 %, 40 %, and 9 %, respectively.

This weight-reducing diet led to a reduction of Fe status in obese patients, as indicated by the decrease in serum ferritin. In our study, a significant decrease in ferritin, with a slight (but not significant) reduction in body mass, was also observed, though only in the women taking sibutramine, and not in the group administered the placebo.

However, in the present study, the decreased level of ferritin is connected with increased GSH-Px and SOD level in the blood, which suggests that the change in serum ferritin level is also associated with a change in inflammation in the body of obese women during the sibutramine treatment. Our results may be consistent with the observation of Zafon et al. [27], who claim that ferritin concentration in the obese is due to inflammation, rather than to the Fe status of the body.

The excretion of minerals in the urine of the studied subjects may be a reflection of a change in the mineral status of the body. Changes in the excretion of minerals are not uniform, as shown in Table 3. This is evidenced by the change in molar ratios of elements in the excreted urine. The molar ratios of minerals in the blood also changed. In each tissue of the body, the ion balance is maintained and remains constant. The evaluation of mineral status and mineral metabolism in the body is possible on the basis of the proportions found to hold between different minerals. There are synergistic and antagonistic relationships between the minerals, which directly affect the metabolism of the body. Keeping an appropriate balance between the elements is, in many cases, more important than their normal levels in the tissue. Changes in the ratio between minerals in tissues indicate metabolic disorders in the body and/or interactions between minerals. In the present study, following the

sibutramine treatment we observed a decrease in the Ca level (though insignificant) and a higher concentration of Mg in serum. Ca and Mg levels in the body are regulated by a negative-feedback system, and through competition for intestinal absorption and renal reabsorption. Ca and Mg also compete for membrane binding sites within the cell. A lower Ca/Mg ratio in the serum, and higher Ca/Mg ration in urine is probably the result of the disturbed Mg status following sibutramine treatment.

Slightly lower Fe and Zn levels, as well as higher Cu concentrations, were found in the serum after 6 and 12 weeks of the study vs. the baseline. These changes may indicate interactions between the elements due to the use of sibutramine. Known interactions of Ca–Mg and Fe–Zn–Cu can be observed. The higher Cu/Fe ratio and lower Zn/Cu ratio in the serum of patients after treatment may be affected by a disturbed Zn status. Decreasing the Zn level in the organism increases the Cu level, which is a result of the Zn–Cu antagonism.

In other studies, low plasma Zn levels were observed in obese [28], and obese type-2 diabetic individuals [29]. Similarly, rather low plasma Zn level was observed in obese women in the current study. In some studies based on hypocaloric diets and physical activity, weight loss in obese women was associated with increased plasma Zn, and this increase was not a reflection of diet [28]. In this study, we observed an inverse relationship. Moreover, Konukoglua et al. [29] found negative correlations between leptin and Zn in serum among obese diabetic subjects. The results obtained in our study may indicate the effects of sibutramine, but not of weight loss or leptin concentration decrease (not analyzed in this study), on changes in the Zn concentration in the serum of obese women. It has been shown that Zn may contribute to the modulation of serotonin uptake in the brain [30]. Sibutramine is a serotonin uptake inhibitor, and so sibutramine by influencing serotonin metabolism might also

indirectly disturb the Zn status of the body. The influence of sibutramine on neurotransmitters may also indirectly affect the level of Cu in the body. It has been found that Cu metabolism is associated with neurological functions (and with the level of serotonin in the brain) [31, 32]. Moreover, Lima et al. did not report any differences in Cu concentration in serum or erythrocytes when comparing obese and non-obese female children [33]. In the present study we observed that, along with the decrease in the concentration of TG and TC, the levels of Fe and Zn lowered significantly, while level of Cu in the serum increased. The results are consistent with the data from our previous study, where obese patients with hypertension and insulin resistance were studied [34].

Treatment with sibutramine resulted in increase in GSH-Px and SOD activity in the serum. GSH-Px is the main enzyme of the enzymatic antioxidant defense system responsible for protecting against increases in ROS production [35]. Hydrogen peroxide, formed by the catalytic reaction of SOD, is both a reactive form of oxygen and a normal cellular metabolite, and it is further detoxified by GSH-Px and catalase [36]. The erythrocyte is a relatively abundant focus of both free-radical-mediated injury by virtue of enhanced endogenous rates of production of ROS and impairment of tissue-free radical defense mechanisms [37, 38]. In addition, GSH is the substrate of GSH-Px, and the cysteine derivative GSH is synthesized in erythrocytes, and glutathione disulfide transported outside the cell to maintain a high GSH/GSSG ratio [35, 38]. Disulfide-exchange reactions occur between protein thiols and low molecular weight disulfides [36]. In our present study, GSH-Px and SOD activities in erythrocytes increased in the treatment groups after the twelfth week. The observed changes may result from an alternation of mineral concentrations during sibutramine treatment. This possible relation was reported by Vivoli et al., who found a positive relationship between serum Cu and Cu–Zn SOD [39]. Moreover Piorunskastoltzman et al. [40] reported a positive correlation between Cu and GSH-Px in serum. The increase of erythrocyte GSH-Px and SOD values in patients during sibutramine treatments has also been attributed to the inhibition of free oxygen radicals and lipid peroxidation [41], and the protective effects of sibutramine supplementation on glucose-6-phosphate dehydrogenase activity in erythrocytes.

Our study has some limitations, for example short treatment and the relatively small group of subjects. These limitations result from group selection procedure, requiring women without chronic diseases and not using additional drugs. Moreover, this study was interrupted by the withdrawal of sibutramine license in the EU. Finally, based on these results and the literature data, it is difficult to explain the precise mechanism of the mineral changes in the serum and urine during sibutramine treatment. A better

understanding of this mechanism requires further extensive biochemical studies.

Conclusion

Sibutramine favorably influences the lipid profile and glucose concentration in obese women. Additionally, treatment with sibutramine is associated with increase in GSH-Px and SOD activity. However, 12-week sibutramine therapy decreases Zn and increases Mg concentrations in the serum, leading to mineral imbalances in obese women. The control of minerals levels, particularly of Zn, and the potential need for further supplementation should be considered in patients treated with sibutramine.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Logue J, Murray HM, Welsh P, Shepherd J, Packard Ch, Macfarlane P, Cobbe S, Ford I, Sattar N (2011) Obesity is associated with fatal coronary heart disease independently of traditional risk factors and deprivation. *Heart* 97(7):564–568
2. Goodpaster BH, DeLany JP, Otto AD, Kuller L, Vockley J, South-Paul JE, Thomas SB, Brown J, McTigue K, Hames KC, Lang W, Jakicic JM (2010) Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults: a randomized trial. *JAMA* 304(16):1795–1802
3. de Lemos H, Atallah AN, de Lemos ALA (2008) Can sibutramine alter systemic blood pressure in obese patients? Systematic review and meta-analysis. *Sao Paulo Med J* 126(6):342–346
4. Smith IG, Goulder MA (2001) Sibutramine Clinical Study 1047 Team. Randomized placebo-controlled trial of long-term treatment with sibutramine in mild to moderate obesity. *J Fam Pract* 50(6):505–512
5. Svacina S, Owen K, Hendl J, Matoulek M, Brychta T (2007) Lipid lowering effect of sibutramine in statin treated patients comparison of lipid lowering effect of sibutramine in patients treated or not treated with statins—3 month follow-up. *Prague Med Rep* 108(4):333–338
6. Derosa G, Cicero AF, Murdolo G, Piccinini MN, Fogari E, Bertone G, Ciccarelli L, Fogari R (2005) Efficacy and safety comparative evaluation of orlistat and sibutramine treatment in hypertensive obese patients. *Diabetes Obes Metab* 7(1):47–55
7. Derosa G, Maffioli P, Salvadeo SA, Ferrari I, Gravina A, Mereu R, D'Angelo A, Palumbo I, Randazzo S, Cicero AF (2010) Sibutramine and L-carnitine compared to sibutramine alone on insulin resistance in diabetic patients. *Intern Med* 49(16):1717–1725
8. Shechter M, Beigel R, Freimark D, Matetzky S, Feinberg MS (2006) Short-term sibutramine therapy is associated with weight loss and improved endothelial function in obese patients with coronary artery disease. *Am J Cardiol* 97:1650–1653

9. Trapali M, Liapi C, Perelas A, Perrea D, Stroubini T, Dontas I, Couvari E, Mavri M, Galanopoulou P (2008) Effect of isocaloric diets and sibutramine on food intake, body mass variation and serum TNF-alpha levels in rats. *Pharmacology* 82(1):15–21
10. Dastych M, Prochazkova D, Pokorny A, Zdrzil L (2010) Copper and zinc in the serum, urine and hair of patients with Wilson's disease treated with penicillamine and zinc. *Biol Trace Elem Res* 133:265–269
11. Paglia DE, Valentine WN (1967) Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med* 70:158–169
12. Misra HP, Fridovich I (1972) The role of superoxide anion in the autooxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem* 247:3170–3175
13. Mahoney JJ, Vreman HJ, Stevenson DK, Van Kessel AL (1993) Measurement of carboxyhemoglobin and total hemoglobin by five specialized spectrophotometers (Co-oximeters) in comparison with reference methods. *Clin Chem* 39:1693–1700
14. National Committee for Clinical Laboratory Standards (NCCLS) (1996) *Clinical Laboratory Procedure Manuals Third Edition (GP2-A3)*
15. Miles AM, Wink DA, Cook JC, Grisham WB (1996) Determination of nitric oxide using fluorescence spectroscopy. *Methods Enzymol* 268:105–120
16. James WP, Astrup A, Finer N, Hilsted J, Kopelman P, Rössner S, Saris WH, Van Gaal LF (2000) Effect of sibutramine on weight maintenance after weight loss: a randomised trial. *STORM Study Group. Sibutramine Trial of Obesity Reduction and Maintenance. Lancet* 356:2119–2125
17. Wadden TA, Berkowitz RI, Womble LG, Sarwer DB, Phelan S, Cato RK, Hesson LA, Osei SY, Kaplan R, Stunkard AJ (2005) Randomized trial of lifestyle modification and pharmacotherapy for obesity. *N Engl J Med* 353:2111–2120
18. Finer N, Bloom SR, Frost GS, Banks LM, Griffiths J (2000) Sibutramine is effective for weight loss and diabetic control in obesity with type 2 diabetes: a randomised, double-blind, placebo-controlled study. *Diabetes Obes Metab* 2:105–112
19. McMahon FG, Fujioka K, Bramah N, Mendel CM, Rowe E, Rolston K, Johnson F, Mooradian AD (2000) Efficacy and safety of sibutramine in obese white and African American patients with hypertension a 1-year, double-blind, placebo-controlled, multicenter trial. *Arch Intern Med* 160:2185–2191
20. Derosa G, Maffioli P, Ferrari I, Palumbo I, Randazzo S, D'Angelo A, Cicero AF (2010) Effects of one year treatment of sibutramine on insulin resistance parameters in type 2 diabetic patients. *J Pharm Pharm Sci* 13(3):378–390
21. Weeke P, Andersson Ch, Fosbol EL, Brendorp B, Kober L, Sharma AM, Finer N, James P, Caterson ID, Rode RA, Torp-Pedersen Ch (2010) The weight lowering effect of sibutramine and its impact on serum lipids in cardiovascular high risk patients with and without type 2 diabetes mellitus—an analysis from the SCOUT lead-in period. *BMC Endocr Disord* 10(3):1–9
22. Tambascia MA, Geloneze B, Repetto EM, Geloneze SR, Picolo M, Magro DO (2003) Sibutramine enhances insulin sensitivity ameliorating metabolic parameters in a double-blind, randomized, placebo-controlled trial. *Diabetes Obes Metab* 5:338–344
23. Doggrel SA (2005) Clinical evidence for drug treatments in obesity-associated hypertensive patients—a discussion paper. *Methods Find Exp Clin Pharmacol* 27(2):119–125
24. GURSOY A, ERDOGAN MF, CIN MO, CESUR M, BASKAL N (2005) Effect of sibutramine on blood pressure in patients with obesity and well-controlled hypertension or normotension. *Endocr Pract* 11(5):308–312
25. Emer E, Kaplan B, Kaya S, Aliyev V, Arica E, Soylemezoglu T (2010) Effect of high dose sibutramine application on serum serotonin, plasma noradrenaline and on zinc and iron levels in liver, kidney, adrenal glands in rats. *Abstracts/Toxicol Lett* 196S:S37–S351. doi:10.1016/j.toxlet.2010.03.798
26. Aguirre ME, Ruz M, Carrasco F, Rebolledo A, Karla A, Codoceo J, Inostroza J (2007) Effect of moderate energy-restricted diets on the nutritional status of selected minerals in obese women. *Arch Latinoam Nutr* 57(3):238–247
27. Zafon C, Lecube A, Simo R (2010) Iron in obesity. an ancient micronutrient for a modern disease. *Obes Rev* 4:322–328
28. Vorugantia VS, Caib G, Klohec DM, Jordan KC, Lane MA, Freeland-Graves JH (2010) Short-term weight loss in overweight/obese low-income women improves plasma zinc and metabolic syndrome risk factors. *J Trace Elem Med Biol* 24:271–276
29. Konukoglu D, Turhana MS, Ercanb M, Serin O (2004) Relationship between plasma leptin and zinc levels and the effect of insulin and oxidative stress on leptin levels in obese diabetic patients. *J Nutr Biochem* 15:757–760
30. Garcia-Colunga J, Reyes-Haro D, Godoy-Garcia IU, Miledi R (2005) Zinc modulation of serotonin uptake in the adult rat corpus callosum. *J Neurosci Res* 80(1):145–149
31. Fujiwara N, Iso H, Kitanaka N, Kitanaka J, Eguchi H, Ookawara T, Ozawa K, Shimoda S, Yoshihara D, Takemura M, Suzuki K (2006) Effects of copper metabolism on neurological functions in Wistar and Wilson's disease model rats. *Biochem Biophys Res Commun* 349(3):1079–1086
32. Parmar P, Daya S (2001) The effect of copper on (3H)-tryptophan metabolism in organ cultures of rat pineal glands. *Metab Brain Dis* 16(3–4):199–205
33. Lima SCVC, Arrais RF, Sales CH, Almeida MG, de Sena KCM, Oliveira UTL, de Andrade AS, Pedrosa LFC (2006) Assessment of copper and lipid profile in obese children and adolescents. *Biol Trace Elem Res* 114:19–26
34. Suliburska J, Bogdanski P, Pupek-Musialik D, Krejpcio Z (2011) Dietary intake and serum and hair concentrations of minerals and their relationship with serum lipids and glucose levels in hypertensive and obese patients with insulin resistance. *Biol Trace Elem Biol* 139:137–150
35. Naziroglu M (2009) Role of selenium on calcium signaling and oxidative stress-induced molecular pathways in epilepsy. *Neurochem Res* 34:2181–2191
36. Kovacic P, Somanathan R (2008) Unifying mechanism for eye toxicity: electron transfer, reactive oxygen species, antioxidant benefits, cell signaling and cell membranes. *Cell Membr Free Radic Res* 2:56–69
37. Naziroglu M, Butterworth PJ (2005) Protective effects of dietary vitamin C and E combination on the blood antioxidative defense mechanism in rats with streptozotocin-induced diabetes. *Can J Appl Physiol* 30:172–185
38. Cimen B (2008) Free radical metabolism in human erythrocytes. *Clin Chim Acta* 390:1–11
39. Vivoli G, Bergomi M, Rovesti S, Pinotti M, Caselgrandi E (1995) Zinc, copper, and zinc- or copper-dependent enzymes in human hypertension. *Biol Trace Elem Res* 49(2–3):97–106
40. Pioruńska-Stolzmann M, Iskra M, Majewski W (2001) The activity of cholesterol esterase and ceruloplasmin are inversely related in the serum of men with atherosclerosis obliterans. *Med Sci Monit* 7(5):940–945
41. Guzman DC, Garcia EH, Mejía GB, Olguín HJ, Jimenez FT, Soto EB, Del Angel DS, Aparicio LC (2012) Effect of sibutramine on 5-hydroxyindole acetic acid levels and selected oxidative biomarkers on brain regions of female rats in the presence of zinc. *Basic Clin Pharmacol Toxicol* 110:421–426. doi:10.1111/j.1742-7843.2011.00829.x