

Erratum to: Iron Deficiency Without Anemia Is Associated with Anger and Fatigue in Young Japanese Women

Takako Sawada¹ · Aki Konomi² · Katsuhiko Yokoi³

Published online: 21 October 2015
© Springer Science+Business Media New York 2015

Erratum to: Biol Trace Elem Res (2014) 159:22–31
DOI 10.1007/s12011-014-9963-1

The authors wish to indicate the following corrections to the referenced article. Under Table 1, the IDNA value 11.6 ± 2.5 (28) and *P* value 0.307 for Menarche (years) are incorrect and should be corrected to 12.0 ± 1.1 (27) and 0.608, respectively. Also, under Table 4, the quartiles value 19 (10.7, 39.5) for Overall, A-R is incorrect and should be corrected to 19 (10.8, 39.5). The corrected tables are shown below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12011-014-9963-1>.

✉ Katsuhiko Yokoi
yokoi@seitoku.ac.jp

¹ Faculty of Food and Nutritional Sciences, Toyo University, Gunma, Japan

² Faculty of Health and Medical Sciences, Teikyo Heisei University, Tokyo, Japan

³ Department of Human Nutrition, Seitoku University Graduate School, Matsudo, Chiba 271-8555, Japan

Table 1 Characteristics of the normal, IDNA, and IDA subjects

	Normal	IDNA	IDA	ANOVA <i>P</i> value
Age (years)	19.4±0.6 ^a (36)	19.9±1.0 ^b (29)	19.8±1.1 ^{a,b} (10)	0.028
Body height (m)	1.58±0.06 (36)	1.59±0.06 (29)	1.57±0.07 (10)	0.554
Body weight (kg)	54.2±8.5 (36)	52.0±6.6 (29)	55.3±8.4 (10)	0.409
Body mass index ¹ (kg/m ²)	21.8±3.3 (36)	20.7±2.4 (29)	22.6±3.6 (10)	0.141
Systolic blood pressure (mmHg)	101.3±9.0 (35)	105.1±9.4 (28)	106.0±14.7 (10)	0.227
Diastolic blood pressure (mmHg)	65.6±8.9 (35)	65.8±7.2 (28)	67.2±7.7 (10)	0.855
Pulse (counts/min)	74.3±9.5 (35)	76.6±9.1 (28)	79.5±17.2 (10)	0.356
Sublingual temperature (°C)	36.3±0.4 (34)	36.3±0.5 (28)	36.6±0.6 (10)	0.173
Birth weight (g)	2938±561 ^a (32)	3289±631 ^b (21)	3399±314 ^b (8)	0.034
Menarche (years)	12.3±1.3 (34)	12.0±1.1 (27)	12.1±0.9 (9)	0.608
Menstrual duration (days)	5.5±1.0 (31)	5.9±1.2 (29)	6.0±1.3 (10)	0.284
Menstrual cycle (days)	29.0±4.3 (26)	29.1±3.0 (27)	28.6±4.9 (9)	0.942

Results are presented as means±SD (*n*). An overall difference between the groups was determined by one-way ANOVA. If the one-way ANOVA was significant ($P<0.05$), differences between individual groups were estimated by using Fisher's protected least significant difference test

IDNA iron deficiency without anemia, IDA iron deficiency anemia

¹ Body mass index (BMI) was calculated as weight (kg) divided by the square of height (m)

Table 4 Comparison of the somatic, mental and overall scores in the CMI-J for the normal, IDNA, and IDA subjects

	Normal (<i>n</i> =36)		IDNA (<i>n</i> =29)		IDA (<i>n</i> =10)		Kruskal-Wallis test <i>P</i> value
	Mean±SD	Median (quartiles)	Mean±SD	Median (quartiles)	Mean±SD	Median (quartiles)	
Somatic, A-L	19.1±15.1	15.5 (7.8, 25.3)	22.7±11.3	19 (16, 31)	18.0±10.7	13.5 (9, 27.5)	0.184
Mental, M-R	8.1±8.3 ^a	5 (2, 12.5)	12.5±9.3 ^b	8 (5, 19)	7.7±10.9 ^{ab}	4.5 (3, 6)	0.024
Overall, A-R	27.3±22.7	19 (10.8, 39.5)	35.2±18.6	33 (21, 42)	25.7±18.4	16.5 (13.3, 37)	0.070

IDNA iron deficiency without anemia, IDA iron deficiency anemia

^{a,b} Values not sharing common superscript letters are significantly different ($P<0.05$) by the Dwass-Steel-Chritchlow-Fligner test