

## Erratum to: NMR resonance assignment of DnaE intein from *Nostoc punctiforme*

Kimmo Heinämäki · Jesper S. Oeemig ·  
Kimmo Pääkkönen · Janica Djupsjöbacka ·  
Hideo Iwai

Published online: 20 November 2012  
© Springer Science+Business Media Dordrecht 2012

**Erratum to: Biomol NMR Assign**  
**DOI 10.1007/s12104-008-9137-1**

There was an error in labeling of “N131” in the [ $^1\text{H}$ ,  $^{15}\text{N}$ ]-HSQC spectrum (Fig. 1). The corrected Fig. 1 is presented


in this article. The third author’s name is also included after reassessment of the contributions. The corrected author list is given in this article.

---

The online version of the original article can be found under doi:[10.1007/s12104-008-9137-1](https://doi.org/10.1007/s12104-008-9137-1).

---

K. Heinämäki · J. S. Oeemig · K. Pääkkönen · J. Djupsjöbacka ·  
H. Iwai (✉)  
Research Program in Structural Biology and Biophysics, Institute  
of Biotechnology, University of Helsinki, P.O. Box 65,  
Helsinki 00014, Finland  
e-mail: [hideo.iwai@helsinki.fi](mailto:hideo.iwai@helsinki.fi)



**Fig. 1** Two-dimensional [ $^1\text{H}$ ,  $^{15}\text{N}$ ]-HSQC spectrum of 2 mM single chain *NpuDnaE* intein in 10 mM sodium phosphate buffer at pH 6.0 at 298 K. The spectrum was recorded at  $^1\text{H}$  frequency of 600 MHz. A total of  $100 (\omega_1) \times 683 (\omega_2)$  complex points were collected for this spectrum, with  $t_{1,\text{max}} = 18.2$  ms,  $t_{2,\text{max}} = 32.0$  ms. The assignments

are indicated by the *number* and *one character codes* for 20 amino acids. The side-chain resonances are marked by *sc*. The side-chain amide groups of asparagines and glutamines are connected by *horizontal lines*