

^1H , ^{13}C and ^{15}N resonance assignments of human BASP1

Leonhard Geist · Anna Zawadzka-Kazimierczuk ·
Saurabh Saxena · Szymon Żerko · Wiktor Koźmiński ·
Robert Konrat

Received: 27 June 2012 / Accepted: 31 October 2012 / Published online: 20 November 2012
© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract Brain acid-soluble protein 1 (BASP1, CAP-23, NAP-22) appears to be implicated in diverse cellular processes. An N-terminally myristoylated form of BASP1 has been discovered to participate in the regulation of actin cytoskeleton dynamics in neurons, whereas non-myristoylated nuclear BASP1 acts as co-suppressor of the potent transcription regulator WT1 (Wilms' Tumor suppressor protein 1). Here we report NMR chemical shift assignment of recombinant human BASP1 fused to an N-terminal cleavable His6-tag.

Keywords BASP1 · NMR signal assignment · Intrinsically disordered protein · Myc oncogene · WT1

Biological context

The homologues of human BASP1 were first identified as the brain-specific proteins CAP-23 (cortical cytoskeleton-associated protein) in chicken brain (Widmer and Caroni 1990) and its rat homologue NAP-22 (neuron-specific acidic protein; Maekawa et al. 1993). Human BASP1 was originally isolated from neuronal cells (Mosevitsky 2005). Interestingly it appears to fulfill quite diverse tasks in the cell. N-myristoylated BASP1 has been described to be

involved in neurite outgrowth and plasma membrane organization (Korshunova et al. 2008). It is able to interact with the inner leaflet of the plasma membrane via its myristoyl-anchor and sequesters Phosphatidylinositol-4,5-diphosphate (PIP2) into lipid rafts (Epand et al. 2004; Shaw et al. 2006). Recently it has been shown that liposomes containing anionic phospholipids induce oligomerization of BASP1. Interaction with calmodulin is followed by dissociation of BASP1 from the membrane and disruption of the oligomers (Zakharov and Mosevitsky 2010). Additionally, BASP1 is under the control of protein kinase C (PKC), which phosphorylates BASP1 at Ser5. It is hypothesized that phosphorylation leads to the disruption of the interaction of the N-terminal positive effector domain of BASP1 with anionic phospholipids (Laux et al. 2000).

Furthermore, non-myristoylated BASP1 appears to influence transcription regulation in the nucleus, greatly affecting the differentiation pathway of a cell. It has been discovered as a co-suppressor of WT1 function (Wilms' Tumor suppressor protein 1) exerting its function by interacting with an N-terminal suppression domain of WT1 (Carpenter et al. 2004; Green et al. 2009). WT1 itself is a potent transcriptional regulator that activates or represses target genes including those for growth factors and regulators of cell division (Wagner and Roberts 2004). Aberrant expression of WT1 is associated with several childhood and adult cancers (Rivera and Haber 2005; Yang et al. 2007). Additionally, a recent study discovered BASP1 to be downregulated in v-Myc-transformed chicken fibroblasts. Strikingly, ectopic expression of BASP1 renders fibroblasts resistant to subsequent cell transformation by v-Myc and it has been shown that the inhibition of v-Myc-induced cell transformation by BASP1 affects the transcriptional regulation of Myc target genes (Hartl et al. 2009). Other findings, reporting the frequent down-regulation of BASP1

L. Geist · R. Konrat (✉)
Department of Computational and Structural Biology, Max F. Perutz Laboratories, University of Vienna, Campus Vienna Biocenter 5, 1030 Vienna, Austria
e-mail: robert.konrat@univie.ac.at

A. Zawadzka-Kazimierczuk · S. Saxena · S. Żerko ·
W. Koźmiński
Faculty of Chemistry, University of Warsaw, Pasteura 1, 02-093
Warsaw, Poland

Table 1 Maximum evolution times (t_{max}, ms) and spectral width (sw, kHz) used for acquisition of spectra for H6-hBASPI

	3D HNCO	4D HabCabCONH	5D HNCOCACB	5D HN(CA)CONH	5D H(CC-tocsy)CONH	5D (H)NCO(NCA)CONH	5D (HACA)CON(CA)CONH
Number of points	700	550	550	550	650	610	610
Experiment duration (hours)	4	13	13	13	17	14	14
sw ₁	2	4	14	6	8	2.5	3.8
sw ₂	2.5	14	14	2.5	18	2	2
sw ₃		2	2	2	2	2	2
sw ₄		2.5	2.5	2.5	2.5	2.5	3.8
t ₁ ^{max}	100	20	10	25	10	50	50
t ₂ ^{max}	150	7.1	10	50	10	45	45
t ₃ ^{max}		45	45	50	45	45	45
t ₄ ^{max}		75	75	75	75	75	75
Sampling density versus conventional	9.3 × 10 ⁻³	4.1 × 10 ⁻⁶	1.7 × 10 ⁻⁶	1.6 × 10 ⁻⁶	2.7 × 10 ⁻⁶	3.2 × 10 ⁻⁸	1.4 × 10 ⁻⁷

expression in ALL (acute lymphocytic leukaemia) and CLL (chronic lymphocytic leukaemia) (Yeoh et al. 2002; Wang et al. 2004), as well as apoptosis-induced cleavage of BASP1 and its subsequent translocation to the cytoplasm (Ohsawa et al. 2008), again highlight the importance of BASP1 in transcription regulation.

To provide molecular information about this potential tumour suppressor protein we have started the NMR structure determination of recombinant human BASP1. The near complete chemical shift assignment reveals that BASP1 belongs to the class of intrinsically disordered proteins.

Methods and results

Protein expression and purification

The coding region for hBASP1 (human BASP1) was amplified by PCR from the mammalian expression vector Flag-hBASP1-pTKX3 (Ohsawa et al. 2008) introducing a 5' NcoI and 3' NotI site. Subsequently the fragment was inserted in-frame into the NcoI and NotI sites of the bacterial expression vector pET-M11 (Pinotsis et al. 2006), yielding pET-M11-hBASP1, encoding hBASP1 fused to an N-terminal His6-tag plus the TEV-cleavage site (H6-hBASP1). ¹⁵N/¹³C labeled H6-hBASP1 was expressed in the *E. coli* strain Rosetta(DE3)pLysS following a new expression protocol for efficient isotopic labeling of recombinant proteins using a fourfold cell concentration in isotopically labeled minimal medium (Marley et al. 2001). The cells were collected after 4 h of expression at 37 °C by centrifugation at 5,000 rpm for 15 min and resuspended in 40 ml of ice-cold lysis buffer (20 mM Na_xH_(3-x)PO₄, 50 mM NaCl, 10 mM imidazole, pH

7.2) per liter of the original bacterial culture. Bacteria were lysed by passing through a French press, and the cell lysate was cleared by centrifugation at 18,000 rpm for 20 min. The supernatant containing the soluble protein fraction was loaded onto a Ni²⁺ loaded HiTrap 5 ml affinity column (GE Healthcare), washed with 2 column volumes of high salt buffer (20 mM Na_xH_(3-x)PO₄, 1.5 M NaCl, 10 mM imidazole, pH 7.2) and eluted with high imidazole buffer (20 mM Na_xH_(3-x)PO₄, 50 mM NaCl, 0.5 M imidazole, pH 7.2) using a linear gradient of 15 column volumes. The H6-hBASP1 containing fractions were collected and the buffer was exchanged by 4 steps of concentration in an Amicon Ultra-15 centrifugal filter device 10 K NMWL (Amicon) and subsequent dilution in target buffer (20 mM Na_xH_(3-x)PO₄, 50 mM NaCl, pH 6.0). NMR samples contain 1.5 mM uniformly ¹⁵N/¹³C labeled protein in 20 mM sodium phosphate (pH 6.0, in 90 % H₂O and 10 % D₂O), 50 mM NaCl and 0.2 % sodium azide.

NMR experiments

All spectra were acquired at 298 K on an Agilent Direct Drive 700 MHz spectrometer using the standard 5 mm ¹H-¹³C-¹⁵N triple-resonance probe head.

The backbone ¹H, ¹³C and ¹⁵N resonances were assigned using sparse random sampling of indirectly detected time domains, in order to increase resolution. A 3D HNCO experiment was used as a base spectrum for SMFT (Sparse Multidimensional Fourier Transform) processing of higher dimensionality experiments (Kazimierczuk et al. 2010). Backbone assignment was achieved using 5D HN(CA)CONH (Kazimierczuk et al. 2010), (HACA)CON(CA)CONH (Zawadzka-Kazimierczuk et al. 2012b), (H)NCO(NCA)CONH (Zawadzka-Kazimierczuk et al. 2012b) and HNCOCACB (Zawadzka-Kazimierczuk et al. 2012b)





Fig. 2 2D spectral planes for consecutive amino acids in H6-hBASPI obtained by SMFT processing of the 5D randomly sampled signal. 2D cross-sections of **a** 5D (HACA)CON(CA)CONH (N_i - CO_{i-1} & N_{i-1} - CO_{i-2}) and **b** 5D HN(CA)CONH (HN_i - N_i & HN_{i+1} - N_{i+1})

Fig. 3 Secondary chemical shifts for **a** $^{13}C'$, **b** $^{13}C^\alpha$, and **c** $^1H^\alpha$ using sequence-specific random coil chemical shifts of intrinsically disordered proteins (Tamiola et al. 2010)



resonance assignment in a 5D (HACA)CON(CA)CONH and HN(CA)CONH experiment. Secondary chemical shifts for $^{13}C'$, $^{13}C^\alpha$, $^1H^\alpha$ (Fig. 3) show only minor deviations

from random coil chemical shift values. Interestingly, the N-terminus appears to harbour stretches with slight α -helical structure propensities, whereas the rest of the

protein seems to adopt a rather extended conformation indicated by positive $^1\text{H}^\alpha$ chemical shift differences.

The ^1H , ^{13}C and ^{15}N chemical shifts have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu/>) under the BMRB accession number 18417.

Acknowledgments The authors thank Professor Masayuki Miura for providing the Flag-hBASP1-pTKX3 plasmid (Ohsawa et al. 2008). All NMR experiments were carried out in the Structural Research Laboratory at the Faculty of Chemistry, University of Warsaw. This work was supported by the EAST-NMR project (contract no. 228461) inside of a transnational access program (proposal acronym: SEQUASSIGNIUPBASP1) and by the Grant P 20549-N19 from the Austrian Science Foundation FWF. A.Z.-K. thanks the Foundation for Polish Science for supporting her with the MPD Programme, S.S., S.Ż., and W.K. thank the Foundation for Polish Science for support with the TEAM Programme. MPD and TEAM programs were co-financed by the EU European Regional Development.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Carpenter B, Hill KJ, Charalambous M, Wagner KJ, Lahiri D, James DI, Andersen JS, Schumacher V, Royer-Pokora B, Mann M, Ward A, Roberts SG (2004) BASP1 is a transcriptional cosuppressor for the Wilms' tumor suppressor protein WT1. *Mol Cell Biol* 24(2): 537–549
- Epand RM, Vuong P, Yip CM, Maekawa S, Epand RF (2004) Cholesterol-dependent partitioning of PtdIns(4,5)P₂ into membrane domains by the N-terminal fragment of NAP-22 (neuronal axonal myristoylated membrane protein of 22 kDa). *Biochem J* 379(Pt 3):527–532. doi:10.1042/BJ20040204
- Green LM, Wagner KJ, Campbell HA, Addison K, Roberts SG (2009) Dynamic interaction between WT1 and BASP1 in transcriptional regulation during differentiation. *Nucleic Acids Res* 37(2): 431–440. doi:10.1093/nar/gkn955
- Hartl M, Nist A, Khan MI, Valovka T, Bister K (2009) Inhibition of Myc-induced cell transformation by brain acid-soluble protein 1 (BASP1). *Proc Natl Acad Sci USA* 106(14):5604–5609. doi:10.1073/pnas.0812101106
- Kazimierczuk K, Zawadzka A, Kozminski W (2009) Narrow peaks and high dimensionalities: exploiting the advantages of random sampling. *J Magn Reson* 197(2):219–228. doi:10.1016/j.jmr.2009.01.003
- Kazimierczuk K, Zawadzka-Kazimierczuk A, Kozminski W (2010) Non-uniform frequency domain for optimal exploitation of non-uniform sampling. *J Magn Reson* 205(2):286–292. doi:10.1016/j.jmr.2010.05.012
- Korshunova I, Caroni P, Kolkova K, Berezin V, Bock E, Walmod PS (2008) Characterization of BASP1-mediated neurite outgrowth. *J Neurosci Res* 86(10):2201–2213. doi:10.1002/jnr.21678
- Laux T, Fukami K, Thelen M, Golub T, Frey D, Caroni P (2000) GAP43, MARCKS, and CAP23 modulate PI(4,5)P(2) at plasmalemmal rafts, and regulate cell cortex actin dynamics through a common mechanism. *J Cell Biol* 149(7):1455–1472
- Maekawa S, Maekawa M, Hattori S, Nakamura S (1993) Purification and molecular cloning of a novel acidic calmodulin binding protein from rat brain. *J Biol Chem* 268(18):13703–13709
- Marley J, Lu M, Bracken C (2001) A method for efficient isotopic labeling of recombinant proteins. *J Biomol NMR* 20(1):71–75
- Mosevitsky MI (2005) Nerve ending “signal” proteins GAP-43, MARCKS, and BASP1. *Int Rev Cytol* 245:245–325. doi:10.1016/S0074-7696(05)45007-X
- Ohsawa S, Watanabe T, Katada T, Nishina H, Miura M (2008) Novel antibody to human BASP1 labels apoptotic cells post-caspase activation. *Biochem Biophys Res Commun* 371(4):639–643. doi:10.1016/j.bbrc.2008.04.056
- Pinotsis N, Petoukhov M, Lange S, Svergun D, Zou P, Gautel M, Wilmanns M (2006) Evidence for a dimeric assembly of two titin/telethonin complexes induced by the telethonin C-terminus. *J Struct Biol* 155(2):239–250. doi:10.1016/j.jsb.2006.03.028
- Rivera MN, Haber DA (2005) Wilms' tumour: connecting tumorigenesis and organ development in the kidney. *Nat Rev Cancer* 5(9):699–712. doi:10.1038/nrc1696
- Shaw JE, Epand RF, Sinnathamby K, Li Z, Bittman R, Epand RM, Yip CM (2006) Tracking peptide-membrane interactions: insights from in situ coupled confocal-atomic force microscopy imaging of NAP-22 peptide insertion and assembly. *J Struct Biol* 155(3):458–469. doi:10.1016/j.jsb.2006.04.015
- Tamiola K, Acar B, Mulder FA (2010) Sequence-specific random coil chemical shifts of intrinsically disordered proteins. *J Am Chem Soc* 132(51):18000–18003. doi:10.1021/ja105656t
- Wagner KJ, Roberts SG (2004) Transcriptional regulation by the Wilms' tumour suppressor protein WT1. *Biochem Soc Trans* 32(Pt 6):932–935. doi:10.1042/BST0320932
- Wang J, Coombes KR, Highsmith WE, Keating MJ, Abruzzo LV (2004) Differences in gene expression between B-cell chronic lymphocytic leukemia and normal B cells: a meta-analysis of three microarray studies. *Bioinformatics* 20(17):3166–3178. doi:10.1093/bioinformatics/bth381
- Widmer F, Caroni P (1990) Identification, localization, and primary structure of CAP-23, a particle-bound cytosolic protein of early development. *J Cell Biol* 111(6 Pt 2):3035–3047
- Goddard TD, Kneller DG (2002) SPARKY 3. University of California, San Francisco
- Yang L, Han Y, Suarez Saiz F, Minden MD (2007) A tumor suppressor and oncogene: the WT1 story. *Leukemia* 21(5): 868–876. doi:10.1038/sj.leu.2404624
- Yeoh EJ, Ross ME, Shurtleff SA, Williams WK, Patel D, Mahfouz R, Behm FG, Raimondi SC, Relling MV, Patel A, Cheng C, Campana D, Wilkins D, Zhou X, Li J, Liu H, Pui CH, Evans WE, Naevae C, Wong L, Downing JR (2002) Classification, subtype discovery, and prediction of outcome in pediatric acute lymphoblastic leukemia by gene expression profiling. *Cancer Cell* 1(2):133–143
- Zakharov VV, Mosevitsky MI (2010) Oligomeric structure of brain abundant proteins GAP-43 and BASP1. *J Struct Biol* 170(3): 470–483. doi:10.1016/j.jsb.2010.01.010
- Zawadzka-Kazimierczuk A, Kozminski W, Billeter M (2012a) TSAR a program for automatic resonance assignment using cross-sections of high dimensionality, high-resolution spectra. *J Biomol NMR* 54:81–95. doi:10.1007/s10858-012-9652-3
- Zawadzka-Kazimierczuk A, Kozminski W, Sanderova H, Krasny L (2012b) High dimensional and high resolution pulse sequences for backbone resonance assignment of intrinsically disordered proteins. *J Biomol NMR* 52:329–337. doi:10.1007/s10858-012-9613-x