

Erratum to: Comparison of Microlife BP A200 Plus and Omron M6 Blood Pressure Monitors to Detect Atrial Fibrillation in Hypertensive Patients

Giuseppe Marazzi · Ferdinando Iellamo · Maurizio Volterrani · Mauro Lombardo ·
Francesco Pelliccia · Daniela Righi · Fabrizia Grieco · Luca Cacciotti ·
Luigi Iaia · Giuseppe Caminiti · Giuseppe Rosano

To view enhanced content go to www.advancesintherapy.com
Published online: December 13, 2014
© Springer Healthcare 2014

Erratum to: Adv Ther (2012) 29(1):64–70
DOI 10.1007/s12325-011-0087-0

The authors of the above-mentioned paper noticed some errors subsequent to publication and would like to make the following corrections.

On page 65 under the heading **Introduction** the sentence “Commercially-available

The online version of the original article can be found under doi:[10.1007/s12325-011-0087-0](https://doi.org/10.1007/s12325-011-0087-0).

G. Marazzi (✉) · F. Iellamo · M. Volterrani ·
G. Caminiti · G. Rosano
Department of Medical Sciences, IRCCS San Raffaele
Pisana, Roma, Italy
e-mail: Giuseppe.marazzi@sanraffaele.it

F. Iellamo · M. Lombardo
University of Rome, Tor Vergata, Italy

F. Pelliccia
University of Rome, La Sapienza, Italy

D. Righi
Cardian Massaia Hospital, Asti, Italy

F. Grieco
University of Naples, Federico II, Italy

L. Cacciotti
Vannini Hospital, Rome, Italy

L. Iaia
Fatebenefratelli Hospital, Isola Tiberina, Rome, Italy

automated devices for self-home BP monitoring with integrated algorithms for detection of AF...” should read “Automated devices for self-home BP monitoring with integrated algorithms for detection of AF...”.

On page 68 under the heading **Discussion** the sentence “The findings of present study on the capability of the OMRON M6 BP monitor of diagnosing 100% AF confirm those of a previous study employing an early version of Omron device (8) and extend those findings to a large population of patients with high BP” should read “The findings of present study on the capability of the OMRON M6 BP monitor of diagnosing 100% AF confirm those of a previous study (8) and extend those findings to a large population of patients with high BP”.