


Erratum to: Early epipelagic life-history characteristics of the North Pacific armorhead *Pentaceros wheeleri*

Chisato Murakami¹  · Shiroh Yonezaki¹ · Satoshi Suyama² · Masayasu Nakagami^{2,3} · Takehiro Okuda¹ · Masashi Kiyota¹

Published online: 10 November 2016
© Japanese Society of Fisheries Science 2016

Erratum to: Fish Sci (2016) 82:709–718 DOI 10.1007/s12562-016-1002-z

In the original publication, in Abstract, the sentence starting with, “Standard length-at-age...respectively”. was incorrectly published as:

Standard length-at-age data were fitted to the mixed model of the von Bertalanffy growth curves, and growth coefficients L_{∞} , k , and t_0 were estimated to be 308 and 290 mm, 0.909 and 1.055, and 0.183 and 0.256 years for males and females, respectively.

The corrected sentence is given below:

Standard length-at-age data were fitted to the mixed model of the von Bertalanffy growth curves, and growth coefficients L_{∞} , k , and t_0 were estimated to be 308 and 290 mm, 0.909 and 1.055, and 0.183 and 0.260 years for males and females, respectively.

The online version of the original article can be found under doi:[10.1007/s12562-016-1002-z](https://doi.org/10.1007/s12562-016-1002-z).

✉ Chisato Murakami
chisato32@affrc.go.jp

- ¹ National Research Institute of Far Seas Fisheries, Japan Fisheries Research and Education Agency, 2-12-4 Fukuura, Kanazawa-ku, Yokohama 236-8648, Japan
- ² Hachinohe Station, Tohoku National Fisheries Research Institute, Japan Fisheries Research and Education Agency, 25-259 Shimomekurakubo, Same, Hachinohe, Aomori 031-0841, Japan
- ³ Research Management Department, Japan Fisheries Research and Education Agency, 15F Queen's Tower B, 2-3-3 Minato Mirai, Nishi-ku, Yokohama, Kanagawa 220-6115, Japan