



Correction to: Predictors of bowel necrosis in patients with acute mesenteric ischemia: systematic review and meta-analysis

Sameh Hany Emile¹ · Sualeh Muslim Khan² · Samer Hani Barsoum¹

Published online: 6 August 2020
© Italian Society of Surgery (SIC) 2020

Correction to: Updates in Surgery
<https://doi.org/10.1007/s13304-020-00857-9>

Table 4: The Column “Biochemical” the entire row “Hyperamylasemia” is missing the updated Table 4 has been copied below

The original article has been updated.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s13304-020-00857-9>.

✉ Sameh Hany Emile
Sameh200@hotmail.com
Sualeh Muslim Khan
sualeh.muslim@yahoo.com
Samer Hani Barsoum
d.samer.barsoum@gmail.com

¹ General Surgery Department, Mansoura University Hospitals, Mansoura University, PO: 35516, Mansoura, Egypt

² Dow Medical College, Dow University of Health Sciences, Karachi, Pakistan

Table 4 Prognostic score for prediction of bowel necrosis in acute mesenteric ischemia

Parameter		Odds ratio (95% CI)	Weight (%)	Points
Clinical	High body mass index	1.11 (1.01–1.22)	0.8	1
	Pulmonary embolism	3.0 (1.2–7.7)	2.2	1
	Coronary heart disease	3.33 (1.17–9.46)	2.4	1
	Shock	3.39 (1.09–10.5)	2.4	1
	Time between symptoms and operation > 90 h	4.8 (1.36–9.89)	3.5	1
	Organ failure	8.47 (2.75–26.1)	6.1	1
	Mesenteric arterial occlusion	26.5 (1.55–453.3)	19.2	2
	Systemic inflammatory response syndrome	28.94 (2.29–365.19)	20.9	3
Biochemical	Leukocytosis > 18.1 × 10 ⁹ /L	1.22 (1.14–1.37)	0.88	1
	Hyperamylasemia	4.22 (1.46–12.16)	3.05	1
	Serum lactate > 2 mmol/L	4.34 (1.7–9.2)	3.1	1
	Platelet-lymphocyte ratio > 156.26	5.04 (1.64–15.49)	3.6	1
	Hemoconcentration Hb > 16 g	6 (1.58–22.75)	4.3	1
	Acidosis with pH < 7.35	6.4 (1.18–58.4)	4.6	1
	Neutrophil-lymphocyte ratio > 11.05	7.57 (2.47–23.16)	5.5	1
Radiological	Splenic vein thrombosis in CT	1.6 (1.2–2.1)	1.1	1
	SMV thrombosis in CT	2.27 (1.22–4.19)	1.6	1
	Bowel loop dilatation in CT	2.29 (1–5.25)	1.6	1
	Portal vein thrombosis in CT	2.7 (1.6–4.4)	1.9	1
	Free intraperitoneal fluid in CT	3.94 (2.41–6.46)	2.8	1
	Pneumatosis intestinalis in CT	10.91 (5.24–22.7)	7.9	1

*0–7=Low probability of bowel necrosis; 8–15=Moderate probability of bowel necrosis; 16–24=High probability of bowel necrosis