

## RESEARCH ARTICLE


# GC-MS Study of Mono- and Bis-haloethylphosphonates Related to Schedule 2.B.04 of the Chemical Weapons Convention: The Discovery of a New Intramolecular Halogen Transfer

Nerea Picazas-Márquez,<sup>1</sup> María Sierra,<sup>1</sup> Clara Nova,<sup>1</sup> Juan Manuel Moreno,<sup>2</sup> Nuria Aboitiz,<sup>1</sup> Gema de Rivas,<sup>2</sup> Miguel A. Sierra,<sup>3</sup> Roberto Martínez-Álvarez,<sup>3</sup> Esther Gómez-Caballero<sup>2</sup>

<sup>1</sup>Ingeniería de Sistemas para la Defensa de España (ISDEFE), Beatriz de Bobadilla 3, E-28040, Madrid, Spain

<sup>2</sup>Laboratorio de Verificación de Armas Químicas (LAVEMA), Área de Defensa Química, Subdirección General de Sistemas Terrestres, INTA, Campus La Marañosa, San Martín de la Vega, E-28330, Madrid, Spain

<sup>3</sup>Departamento de Química Orgánica, Facultad de Ciencias Químicas, Universidad Complutense, E-28040, Madrid, Spain



**Abstract.** A new class of compounds, mono- and bis-haloethylphosphonates (HAPs and bisHAPs, respectively), listed in Schedule 2.B.04 of the Chemical Weapons Convention (CWC), has been synthesized and studied by GC-MS with two aims. First, to improve the identification of this type of chemicals by the Organization for the Prohibition of Chemical Weapons, (OPCW). Second, to study the synergistic effect of halogen and silicon atoms in molecules undergoing mass spectrometry. Fragmentation patterns of trimethylsilyl derivatives of HAPs were found to depend on the nature of the halogen atom; this was in agreement with DFT-calculations. The data suggest that a novel intramolecular halogen transfer takes place during the fragmentation process.

**Keywords:** Iodine transfer, Haloethylphosphonates, Chemical weapons convention

Received: 10 February 2016/Revised: 19 May 2016/Accepted: 24 May 2016/Published Online: 14 June 2016

## Introduction

Recent use of chemical warfare agents (CWAs) during the Syrian conflict has caused international apprehension [1]; in 1997, the CWC was created to protect the world from any CWA attack; it prohibits the development, production, acquisition, stockpiling, retention, transfer, or use of chemical weapons [2]. The OPCW, the compliance body for CWC, holds inspections that may include sampling and on-site or


off-site analysis [3]. The chemicals subject to control by the OPCW are listed in an Annex of the CWC under various schedules relating to military and nonmilitary purposes, chemical nature, etc. To carry out the mandatory CWAs verifications, the OPCW maintains an analytical database, the Central Analytical Database (OCAD), which is a collection of gas chromatography retention indexes (GC(RI)), infrared (IR), mass spectrometry (MS), and nuclear magnetic resonance (NMR) spectra for scheduled chemicals and related compounds [4]. The primary on-site analytical tool is gas chromatography/mass spectrometry (GC/MS). Therefore, GC(RI) and MS data in the OCAD are the primary analytical references [5]. Because these efforts are of critical international importance, a large number of scientific articles regarding GC/MS analysis of CWAs and related compounds has been reported [6–26].

The compounds of the present study belong to Schedule 2.B.04, which includes “all chemicals, except for those listed in

Dedicated to Professor Michael Hanack (University of Tübingen, Germany) on the occasion of his 85th birthday.

**Electronic supplementary material** The online version of this article (doi:10.1007/s13361-016-1430-0) contains supplementary material, which is available to authorized users.

Correspondence to: Roberto Martínez-Álvarez; e-mail: rma@quim.ucm.es



**Scheme 1.** Synthesis of HAPs, bisHAPs, and TMS derivatives investigated

Schedule 1, containing a phosphorus atom to which is bonded one methyl, ethyl, or propyl (normal or *iso*) group but no additional carbon atoms” [2]. Since the degradation of Scheduled 1 chemicals may lead to the formation of this kind of organophosphorous compounds, some of them are considered environmental markers of nerve agents [19, 24, 26]. Schedule 2.B.04 is open to many compounds. Therefore, an understanding of the different classes of compounds in 2.B.04 is of paramount importance. In this study, we report the EI-MS data for new bis-haloethylphosphonates (bisHAPs) and trimethylsilyl derivatives of mono-haloethylphosphonic acids (HAPs). Through a combination of experimental and computational methods, we have discovered new, unprecedented fragmentation patterns of some of these compounds. Specifically, an iodine transfer may be of relevance in the spectral interpretation of compounds in the OCAD and related data bases.

## Experimental

### General

All the chemicals required for the synthesis of HAPs and bisHAPs as well as the solvents and derivatizing agent *N,N*-bis(trimethylsilyl)trifluoroacetamide (BSTFA) are commercially available and were used without further purification.

### Synthesis of bisHAPs and HAPs and Derivatization with BSTFA (Scheme 1)


*N,N*-dicyclohexylcarbodiimide (DCC, 0.15 mmol) and 4-dimethylaminopyridine (DMAP, 0.04 mmol) were added to a solution of the corresponding alkoxyphosphonic acid (APA, 0.1 mmol) in dichloromethane (1 mL) at room temperature. The mixture was stirred at 60 °C for 1 h. Immediately following, the corresponding haloalcohol (0.44 mmol) was poured into the reaction and stirred at 60 °C for 3 h. Stock solutions were prepared by dilution to the desired concentration with dichloromethane and by derivatization with 200  $\mu$ L of BSTFA at 60 °C for 30 min in 2 mL sealed glass vials when necessary. The corresponding bisHAPs and silylated HAPs were analyzed by GC/MS.

### GC/MS(EI) Analysis

The GC/MS analyses were conducted in electron ionization (EI) mode (70 eV) using an Agilent 6890 gas chromatograph, equipped with a 5973 MSD model (Agilent Technologies, Palo Alto, CA, USA). An HP5 column (5% phenyl 95% methylpolysiloxane; 30 m length, 0.25 mm i.d., and 0.25 mm film thickness) was used at a temperature program of 40 °C (1 min),

**Table 1.** Significant Peaks 25–29 (*m/z*, Relative Peak Intensities) in the Mass Spectra of Monohalogenated TMS Derivatives 1–12

Compound	R	X	25	26	27	28	29
<b>1</b>	Me	Cl	215 (30)	195(55)		153 (100)	151 (40)
C <sub>6</sub> H <sub>16</sub> ClPO <sub>3</sub> Si							
<b>2</b>	Me	Br	259 (20)	195 (65)		153 (100)	151 (40)
C <sub>6</sub> H <sub>16</sub> BrPO <sub>3</sub> Si							
<b>3</b>	Me	I	307 (<10)	195 (75)	279 (100)	153 (65)	151 (50)
C <sub>6</sub> H <sub>16</sub> IPO <sub>3</sub> Si			306.9439	195.0446	278.8776	152.9995	151.0257
<b>4</b>	Et	Cl	229 (30)	209 (72)		167 (100)	165 (38)
C <sub>7</sub> H <sub>18</sub> ClPO <sub>3</sub> Si							
<b>5</b>	Et	Br	275 (20)	209 (72)		167 (100)	165 (50)
C <sub>7</sub> H <sub>18</sub> BrPO <sub>3</sub> Si							
<b>6</b>	Et	I	321 (<10)	209 (80)	293 (100)	167 (60)	165 (50)
C <sub>7</sub> H <sub>18</sub> IPO <sub>3</sub> Si				209.0660	292.8932	167.0181	165.0417
<b>7</b>	Pr	Cl	243 (45)	223 (80)		181 (100)	
C <sub>8</sub> H <sub>20</sub> ClPO <sub>3</sub> Si							
<b>8</b>	Pr	Br	287 (30)	223 (95)		181 (100)	179 (60)
C <sub>8</sub> H <sub>20</sub> BrPO <sub>3</sub> Si							
<b>9</b>	Pr	I	335 (<10)	223 (100)	307 (98)	181 (50)	179 (70)
C <sub>8</sub> H <sub>20</sub> IPO <sub>3</sub> Si				223.0793	306.9146		179.0565
<b>10</b>	<i>i</i> -Pr	Cl	243 (50)	223 (100)		181 (100)	
C <sub>8</sub> H <sub>20</sub> ClPO <sub>3</sub> Si							
<b>11</b>	<i>i</i> -Pr	Br	287 (35)	223 (100)		181 (85)	179 (70)
C <sub>8</sub> H <sub>20</sub> BrPO <sub>3</sub> Si							
<b>12</b>	<i>i</i> -Pr	I	335 (10)	223 (90)	307 (100)	181 (60)	179 (80)
C <sub>8</sub> H <sub>20</sub> IPO <sub>3</sub> Si				223.9081	306.9513	181.0529	179.0773



**Scheme 2.** Proposed fragmentation pattern for TMS derivatives of (2-haloethyl)alkylphosphonates **1–12**

then raised at 10 °C/min to 280 °C, where it was held for 5 min. Helium was used as the carrier gas at a constant flow of 1 mL/min. The samples were injected in splitless mode at an injection temperature of 250 °C and 1  $\mu\text{L}$  volume. The spectra were acquired at an EI source temperature of 230 °C and a quadrupole analyzer temperature of 150 °C, with unit mass resolution, scan range  $m/z$  40–500 and a scan cycle of 3.15 scans/s. The


GC/MS plots as well as the retention indexes (RIs) of the investigated compounds are included in the [Supplementary Material](#). HRMS spectra were recorded using a TOF GCT Premier from Waters (Manchester, UK) Micromass coupled with a gas chromatograph 7890A from Agilent with continuous internal calibration, scan time of 0.9 s, and inter-scan delay of 0.1 s, and are included in the [Supplementary Material](#).

### GC/MS(CI) Analysis

The molecular mass of the compounds subject to this study was confirmed by GC/MS analysis conducted in chemical ionization (CI) mode using a Varian (Palo Alto, CA, USA) 3400 gas chromatograph, equipped with a Saturn 2000 ion trap. Chromatographic conditions were identical to those described for above. The spectra were acquired at a CI source temperature of 200 °C and acetonitrile as reactant gas, with unit mass resolution, scan range  $m/z$  80–500 and a scan cycle of 2.5 scans/s.

### Computational Details

All calculations were performed at the DFT level using the uB3LYP functional [27, 28] as implemented in Gaussian 09 [29]. Cl, Br, and I atoms were described using the lan12dz pseudopotential [30]. The 6-31G\*\* basis set was used for the H, C, N, and O atoms [31, 32]. All structures of the reactants, intermediates, transition states, and products were fully optimized in vacuo [31]. Transition states were identified by having one imaginary frequency in the Hessian matrix. It was



**Figure 1.** EI Mass spectrum of 2-iodoethyl trimethylsilyl methylphosphonate **3**






Figure 2. EI Mass spectrum of 2-chloroethyl trimethylsilyl methylphosphonate **1**

confirmed that transition states connect with the corresponding intermediates by means of application of the eigenvector corresponding to the imaginary frequency

and subsequent optimization of the resulting structures. All energies collected in the text are Gibbs energies at 298 K.



Scheme 3. Calculated reaction pathway for the fragmentation of **I1**.  $\Delta G^\ddagger$  values are given in  $\text{kcal mol}^{-1}$



**Scheme 4.** Proposed fragmentation pattern for 3-iodopropyl trimethylsilyl methylphosphonate **30**

## Results and Discussion

Unmodified mono-haloethylphosphonates (HAPs) cannot be analyzed by GC since their high polarity and nonvolatile nature preclude their submission without degradation. To overcome this problem, these compounds were treated with BSTFA to form the corresponding monosilylated derivatives **1–12**. Table 1 compiles the most significant ions observed in the EI mass spectra of the prepared TMS derivatives while the corresponding mass spectra can be found in the [Supplementary Material](#) (SF1–16). Peaks for the molecular ions were not observed. The molecular mass of **1–12** was confirmed, however, by GC/MS(CI) ([Supplementary Material SF17–19](#)).

Data in Table 1 show that the first fragmentation corresponds to a methyl radical elimination, probably from the trimethylsilyl group, to form the corresponding fragments **25**. The loss of a methyl radical bonded to the silicon atom is a well known

fragmentation in the mass spectrometry of these derivatives [33]. This elimination is independent of the nature of the halogen atom (chlorine or bromine) in the molecule. Variation of the alkyl group attached to phosphorus atom produces no significant effect in the general fragmentation of these TMS derivatives. The loss of a methyl radical is followed by the elimination of a vinyl halide molecule ( $\text{CH}_2=\text{CHX}$ ) to form the base peak of the spectra (**28**), except when the halogen is iodine (compounds **3**, **6**, **9**, and **12**). The formation of fragments such as **26** for chlorine and bromine derivatives can be explained as a result of a direct elimination of a halogen radical from the undetected molecular ion (Scheme 2). On the other hand, iodine derivatives behave differently; the loss of a methyl group produces fragments of very low abundance, which subsequently undergo an ethylene molecule elimination leading to fragments such as **27**. Accurate mass measurements confirm these two consecutive eliminations. Figure 1 (compound **3**) is representative of the behavior of these iodinated TMS derivatives, whereas Figure 2 shows the spectra of a chlorine homologue (**1**) for comparison.

To explore the remarkable differences between iodides (**3**, **6**, **9**, and **12**) and their chloro- and bromo-substituted congeners, DFT calculations were carried out for compounds **1**, **2**, and **3** ( $\text{R} = \text{Me}$ ,  $\text{X} = \text{Cl}$ ,  $\text{Br}$ ,  $\text{I}$ ) (see computational procedures and Scheme 3). Regardless of the halogen, the loss of one Me-group attached to the silicon easily occurs from the initially formed radical cation **11**. This process is assisted by the  $\text{P}=\text{O}$  bond forming the cyclic intermediates **12** through **TS1**. This step is more difficult for bromide **2** ( $\Delta G^\ddagger = 19.8 \text{ kcal mol}^{-1}$ ) and for chloride **1** ( $\Delta G^\ddagger = 15.5 \text{ kcal mol}^{-1}$ ) being easiest for iodide **3** ( $\Delta G^\ddagger = 10.7 \text{ kcal mol}^{-1}$ ). Nevertheless, the differences encountered between the three halogens have to be found lately in the reaction mechanism leading to **27**, since the evolution of **12** to **28** or **29** takes place irrespective of the halogen. Subsequently, **12** evolves to **13** through **TS2**, which involves the attack of the halogen ( $\text{Cl}$ ,  $q = -0.03$ ;  $\text{Br}$ ,  $q = 0.01$ ;  $\text{I}$ ,  $q = 0.12$ ) to the electron deficient silicon center (when  $\text{X} = \text{Cl}$ ,  $q = 1.58$ ;  $\text{X} = \text{Br}$ ,  $q = 1.58$ ;  $\text{X} = \text{I}$ ,  $q = 1.58$ ). In this step, the oxygen tethered to the silicon in **12** is displaced, forming the cyclic intermediate **13**. Now, the

**Table 2.** Significant Peaks **31–36** ( $m/z$ , Relative Peak Intensities) in the Mass Spectra of Chloride and Bromide bisHAPs

Compound	R	X	31	32	33	34	35	36
<b>13</b> $\text{C}_5\text{H}_{11}\text{Cl}_2\text{PO}_3$	Me	Cl	185 (100)	159 (32)	97 (34)	63 (42)	141 (74)	79 (62)
<b>16</b> $\text{C}_6\text{H}_{13}\text{Cl}_2\text{PO}_3$	Et	Cl	199 (100)	173 (11)	111 (29)	63 (26)	155 (83)	93 (30)
<b>19</b> $\text{C}_7\text{H}_{15}\text{Cl}_2\text{PO}_3$	Pr	Cl	213 (100)	-	125 (23)	63 (40)	169 (76)	107 (<2)
<b>22</b> $\text{C}_7\text{H}_{15}\text{Cl}_2\text{PO}_3$	<i>i</i> Pr	Cl	213 (100)	187 (19)	125 (15)	63 (32)	169 (73)	107 (<2)
<b>14</b> $\text{C}_5\text{H}_{11}\text{Br}_2\text{PO}_3$	Me	Br	229 (81)	203 (21)	97 (41)	107 (100)	185 (48)	79 (74)
<b>17</b> $\text{C}_6\text{H}_{13}\text{Br}_2\text{PO}_3$	Et	Br	243 (92)	217 (24)	111 (34)	107 (100)	199 (64)	93 (44)
<b>20</b> $\text{C}_6\text{H}_{13}\text{Br}_2\text{PO}_3$	Pr	Br	257 (59)	229 (25)	125 (18)	107 (100)	213 (32)	107 (25)
<b>23</b> $\text{C}_6\text{H}_{13}\text{Br}_2\text{PO}_3$	<i>i</i> Pr	Br	257 (100)	231 (18)	125 (20)	107 (100)	213 (62)	107 (<2)

**Table 3.** Significant Peaks 37–41 ( $m/z$ , Relative Peak Intensities) in the Mass Spectra of Iodine bisHAPs

Compound	R	37	38	39	40	41
<b>15</b>	Me	250 (38)	97 (18)	123 (100)	79 (22)	155 (84)
$C_5H_{11}I_2PO_3$		249.9190		123.0165		154.9340
<b>18</b>	Et	264 (28)	111 (24)	137 (100)	93 (16)	155 (75)
$C_6H_{13}I_2PO_3$		263.9308		137.0324		154.9340
<b>21</b>	Pr	278 (22)	125 (25)	151 (100)	107 (13)	155 (70)
$C_7H_{15}I_2PO_3$		277.9475				
<b>24</b>	<i>i</i> Pr	278 (26)	125 (25)	151 (100)	107 (9)	155 (65)
$C_7H_{15}I_2PO_3$		277.9755		151.0617		154.9444


energies for this displacement are very similar for the three halogens with the iodine being slightly disfavored. Cyclic cation **13** evolves to **14** by an intramolecular attack of the oxygen tethered to the halogenoethenyl chain. The energies involved are again similar for the three halogens with iodine being slightly disfavored compared with chlorine or bromine. The final step of the mechanism consists of the extrusion of an ethylene molecule from **13** to give the observed **27** (M-43). This process is analogous to one reported for the fragmentation of halogen-substituted amines via intramolecular halogen transfer [34]. This rate-determining step is clearly favored for the more nucleophilic iodine ( $\Delta G^\ddagger = 48.2 \text{ kcal mol}^{-1}$ ) compared with chlorine ( $\Delta G^\ddagger = 68.8 \text{ kcal mol}^{-1}$ ) or bromine ( $\Delta G^\ddagger = 57.3 \text{ kcal mol}^{-1}$ ). These energies are congruent with the experimentally observed behavior of compounds **1**–**3**.

To discard other possible fragmentation pathways, we have computed the direct halogen transfer to the P=O bond in **11** as an alternative hetero McLafferty type six-membered rearrangement to explain both the Me and ethylene eliminations, together with the reactivity differences between halogens. So far, we have been unable to find a reaction pathway connecting **11** with the product resulting from the transfer of the iodine atom to the P=O bond. However, a high energy pathway connecting **11**

with **27** (M-43) was found by attacking the silicon atom of **11** with the oxygen tethering the halogenoethenyl chain. This pathway is similar to the one depicted in Scheme 3. Although this pathway is also suitable to explain the different behavior of iodine compared with chlorine or bromine, the energies involved in the key step are  $>100 \text{ kcal mol}^{-1}$ . Therefore, it seems less probable than the proposed mechanism pathway in Scheme 3. It is remarkable that although some heteroatom transfers have been reported as examples of the hetero McLafferty rearrangement [35–39], no data for similar iodine consecutive transfers have been described in the literature.

The proposed mechanism for the formation of fragment **27** (Scheme 2) is supported by the mass spectra of 3-iodopropyl trimethylsilyl methylphosphonate **30** (M = 336, SF20) which was prepared (see Experimental) using 3-iodopropanol as the corresponding alcohol. The spectrum reveals a base peak at  $m/z$  153, the proposed formation of which begins with Me-cleavage. Formation of an oxetane ring, however, is disfavored since its transition state homologous to **TS2** in Scheme 3 would have an eight-membered ring. Instead, the elimination of methyl radical is followed by the loss of a propenyl iodide molecule (168 Da) to form the main fragment at  $m/z$  153 (Scheme 4). Nevertheless, the fragment at  $m/z$  279, generated as a consequence of consecutive eliminations of methyl radical and propylene molecule (42 Da), involving a iodine atom transfer similar to the calculated in Scheme 2, is formed albeit in low abundance ( $<10\%$ ). All the attempts to compute a reaction pathway connecting **12** (Scheme 3) to the corresponding **13** for the homologous **30** were fruitless.

Finally, a comparative study was made with the corresponding ethyl trimethylsilyl alkylphosphonate acids [4, 40] (SF21–24). Similar to the results for compounds **1**–**12**, the elimination of the methyl radical (M-15) that forms product-ions as **25** (Scheme 2, X = H) followed by the loss of an ethylene molecule leads to the base peaks with structure identical to **28**.

**Scheme 5.** Proposed fragmentation pattern for bis(2-chloroethyl) and bis(2-bromoethyl) alkylphosphonates




Figure 3. EI Mass spectrum of bis(2-chloroethyl)methylphosphonate **13**

The EI mass spectra of all bisHAPs studied (**13–24**) lacked the odd-electron molecular ions (Tables 2 and 3, SF 25–40). The molecular masses of these compounds were confirmed by GC/MS(CI) (SF 41–43). Furthermore, fragments from the cleavage of the P–C bond were also absent from the spectra, whereas this dissociation was detected in the mass spectra of  $\alpha$ -amino phosphonates [41] and spiro alkylphosphonates [42]. Again,

the nature of the halogen atom (X) plays a crucial role in the fragmentation pathway of compounds **13–24**.

Thus, the homolytic cleavage of the C–halogen bond with the subsequent loss of the corresponding halogen radical is the main fragmentation for chlorinated bisHAPs to form the ion fragments **31** (Table 2 and Scheme 5). Figure 3 shows the EI mass spectrum of bis(2-chloroethyl)methylphosphonate **13** ( $M$





Figure 4. EI Mass spectrum of bis(2-bromoethyl)methylphosphonate **14**




**Scheme 6.** Proposed fragmentation pattern for bis(2-iodoethyl) alkylphosphonates

= 220) with the base peak at *m/z* 185 (fragment type **31**, R = Me, X = Cl). Although brominated bisHAPs also eliminate bromine radical to form an ion at *m/z* 229, this ion is not the base peak. The main peak for the brominated compounds corresponds to the three membered cyclic ion  $[C_2H_4Br]^+$  (**34**, X = Br, *m/z* 107). Figure 4 shows the EI mass spectrum of bis(2-bromoethyl)methylphosphonate **14** (M = 308).

Fragmentations involving a double hydrogen atom transfer to the P=O bond (McLafferty + 1 rearrangement) can also be observed for chlorine and bromine bisHAPs. Thus, the elimination of a halogen vinylidene radical ( $XCH=CH^\bullet$ ) leads to the formation of fragments type **32**, which evolve either by loss of a water molecule, forming **35**, or by elimination of a vinylhalogenide molecule leading to **33**. The subsequent elimination of a vinyl halide molecule ( $CH_2=CHX$ ) from **35** generates the fragment ion **36** (Scheme 5).

To better understand the mass spectrometry of these halogenated compounds, a comparative study with the corresponding diethyl alkylphosphonates (dEAPs) was carried out [4, 40, 43]. Similar to the observed fragmentations of bisHAPs, diethyl methylphosphonate and diethyl ethylphosphonate (SF 44–45) undergo a McLafferty type-rearrangement with a double hydrogen atom transfer to the P=O bond with loss of a halogen vinylidene radical ( $XCH=CH^\bullet$ ) forming ions at *m/z* 125 and 139, respectively [4, 40, 43]. Two subsequent eliminations of ethylene and water molecules, respectively, lead to the formation of the most significant peaks of the spectra.

Differences were found, however, between the fragmentations of bisHAP chlorides, bromides, and iodides (Table 3 and Scheme 6). The reason could be attributed to the C-I bond's lower dissociation energy compared with those for the C-Cl and C-Br bonds [44, 45]; this would lead to a different fragmentation mechanism. The mass spectra of the iodides lack  $M^+$  and  $[M - I]^+$  ions. Instead, a molecule of vinyl iodide is promptly eliminated, affording fragments such as **37**, which evolve either by undergoing a McLafferty rearrangement with double hydrogen transfer to form **38**, or by loss of an iodine radical to form **39**. Additionally, a direct elimination of



**Figure 5.** EI Mass spectrum of bis(2-iodoethyl)methylphosphonate **15**



iodoethylene radical (155 Da) produces fragments in low abundance. The EI mass spectrum of bis(2-iodoethyl)methylphosphonate **15** (Figure 5) reveals peaks at  $m/z$  155 [ $C_2H_4I$ ]<sup>+</sup> and the product of elimination of 155 Da (M-155,  $m/z$  249). The presence of both fragments indicates that a competition process to retain the charge between these fragments is taking place. The fragment at  $m/z$  155 (**41**) shows a higher abundance because it can retain the charge more easily than the other fragment (M-ICH<sub>2</sub>CH<sub>2</sub>•).

Although it was mentioned above that the nature of the alkyl group attached at the phosphorus atom (methyl and ethyl) plays no significant role in the fragmentation patterns, the small differences in the pathways between compounds bearing propyl and *isopropyl* groups are worth of discussion. These differences are more easily observed in the chlorine derivatives **19** and **22** (SF 27, 30). Thus, **22** with an *isopropyl* group attached at the P atom undergoes a fragmentation pathway where both the elimination of the chlorine radical and the McLafferty fragmentation with double hydrogen transfer (ML + 1) produces peaks at  $m/z$  213 (**31**, R = C<sub>3</sub>H<sub>7</sub>, X = Cl) and  $m/z$  187 (**32**, R = C<sub>3</sub>H<sub>7</sub>, X = Cl), respectively. Elimination of a water molecule from **32** produces the fragment **35** at  $m/z$  169 (R = C<sub>3</sub>H<sub>7</sub>, X = Cl). By contrast, the linear propyl group (**19**) induces a different pathway where **43** (M-Cl,  $m/z$  213) undergoes the elimination of an ethylene molecule (28 Da) from the linear propyl moiety forming a fragment at  $m/z$  185, which cannot be observed in the mass spectrum of **22**. The absence of fragment **32** ( $m/z$  187) indicates that in this case, the double H rearrangement does not take place for **22**.

## Conclusions

Two new classes of mono- and bis-haloethylphosphonates listed in Schedule 2.B.04 of the CWC were synthesized and studied by GC-MS. For trimethylsilylated HPAs a clear halogen-dependent differential behavior in their fragmentation patterns was observed. Thus, chlorine and bromine derivatives fragment by losing a Me group bonded to the silicon, followed by extrusion of a vinylhalide molecule. In contrast, iodine substituted HPAs follow a much more complex fragmentation pattern, including three iodine migrations. This fragmentation behavior has been further examined by DFT calculations, which are fully consistent with the experimental data. The newly described behavior of iodine derivatives of Schedule 2.B.04 may be important in the analysis of these compounds.

Fragmentation of bis-HPAs is also halogen-dependent. Thus, the main fragmentation of dichloro and dibromo bis-HPAs is the cleavage of the carbon-halogen bond with loss of the corresponding halogen radical. In contrast, diiodo HPAs present as the base peak fragment originated through two consecutive eliminations of vinyl iodide and a iodine radical, respectively.

## Acknowledgments

Financial support from the Spanish MINECO (CTQ2013-46459-C2-1P; and CTQ2014-51912-REDC) is acknowledged. The authors are very grateful for the help of Professor John Almy (CSU, Stanislaus, CA, USA) for fruitful discussions and careful editing of the manuscript.

## References

- United Nations Mission to Investigate Allegations of the Use of Chemical Weapons in the Syrian Arab Republic: Report on the Alleged Use of Chemical Weapons in the Ghouta Area of Damascus on 21 August 2013. United Nations, New York (2013)
- Convention on the Prohibition of the Development, Production, Stockpiling, and Use of Chemical Weapons and their Destruction, Technical Secretariat of the Organization for Prohibition of Chemical Weapons. The Hague (1997). Available at: <http://www.opcw.nl>. Accessed 6 June 2016
- Rautio, M.: Recommended Operating Procedures for Sampling and Analysis in the Verification of Chemical Disarmament. Ministry for Foreign Affairs of Finland, Helsinki (1994)
- Central OPCW Analytical Database, version 18, OCAD v.18\_2016 Technical Secretariat of the OPCW: The Hague (2016)
- Wils, E.R.J.: Gas chromatography/mass spectrometry in analysis of chemicals related to the Chemicals Weapons Convention. In: Meyers, R.A. (ed.) Encyclopedia of Analytical Chemistry. John Wiley, Chichester (2000)
- Bauer, G., Vogt, W.: Gas chromatographic determination of acids derived from phosphorus by trimethylsilylation with N, O-bis(trimethylsilyl)trifluoroacetamide. Anal. Chem. **53**, 917–919 (1981)
- Rohrbaugh, D.K., Sarver, E.W.: Detection of alkyl methylphosphonic acid in complex matrices by gas-chromatography-tandem mass spectrometry. J. Chromatogr. A **809**, 141–150 (1998)
- Purdon, J.G., Pagotto, J.G., Miller, R.K.: Preparation, stability, and quantitative analysis by gas chromatography and gas chromatography-electron impact mass spectrometry of *tert*-butyldimethylsilyl derivatives of some alkylphosphonic and alkyl methylphosphonic acids. J. Chromatogr. A **475**, 261–272 (1989)
- Riches, J., Morton, I., Read, R.W., Black, R.W.: The trace analysis of alkylphosphonic acids in urine using gas chromatography-ion trap negative ion tandem mass spectrometry. J. Chromatogr. B **816**, 251–258 (2005)
- Jang, Y.J., Kim, K., Tsay, O.G., Atwood, D.A., Churchill, D.G.: Update 1 of destruction and detection of chemical warfare agents. Chem. Rev. **115**, PR1–PR76 (2015)
- Blum, M.M., Murty Mamidanna, R.V.S.: Analytical chemistry and the chemical weapons convention. Anal. Bioanal. Chem. **406**, 5067–5069 (2014)
- Harvey, D.J., Hoving, M.G.: Derivatives for the characterization of alkyl- and aminoalkylphosphonates by gas chromatography and gas chromatography-mass spectrometry. J. Chromatogr. A **79**, 65–74 (1973)
- Black, R.M., Harrison, J.: In: Hartley, F.R. (ed.) The chemistry of organophosphorus compounds, vol. 4, pp. 781–890. Wiley-Interscience, Chichester (1996)
- Timperley, C.M.: In: Timperley, C.M. (ed.) Best synthetic methods. Organophosphorus (V) chemistry. Academic Press, London (2015)
- Sass, S., Fisher, T.L.: Chemical ionization and electron impact mass spectrometry of some organophosphonate compounds. Org. Mass Spectrom. **14**, 257–264 (1979)
- Wils, E.R.J., Hulst, A.G.: Mass spectra of some pinacolyl-containing organophosphorus compounds. Org. Mass Spectrom. **21**, 763–765 (1986)
- Borrett, V.T., Mathews, R.J., Mattsson, E.R.: Verification of the chemical weapons convention: mass spectrometry of alkyl methylphosphonofluoridates. Aust. J. Chem. **47**, 2065–2074 (1994)
- Chen, H., Zheng, X., Cooks, R.G.: Ketalization of phosphonium ions by 1,4-dioxolane: selective detection of the chemical warfare agent stimulant DMMP in mixtures using ion/molecule reactions. J. Am. Soc. Mass Spectrom. **14**, 182–188 (2003)
- D'Agostino, P.A., Provost, L.R.: Mass spectrometric identification of products formed during degradation of dimethylphosphoramidocyanidate (tabun). J. Chromatogr. A **598**, 89–95 (1992)
- Witkiewicz, Z., Mazurek, M., Szulc, J.: Chromatographic analysis of chemical warfare agents. J. Chromatogr. A **503**, 293–357 (1990)

21. Reddy, T.J., Mirza, S.P., Vijaya Saradhi, U.V.R., Rao, V.J., Mairamani, M.: Mass spectral studies of *N,N*-dialkylaminoethanols. *Rapid Commun. Mass Spectrom.* **17**, 746–752 (2003)
22. Saeidian, H., Ashrafi, D., Sarabadani, M., Naseri, M.T., Babri, M.: Mass spectrometric study of O(S)-alkyl N, N-dimethylamino alkylphosphonates (alkylphosphonothiolates) for chemical weapons convention verification purposes. *Int. J. Mass Spectrom.* **319**, 9–16 (2012)
23. Gupta, A.K., Pardasani, D., Kanaujia, P.K., Tak, V., Dubey, D.K.: Gas chromatography/mass spectrometric analysis of N, N-dialkylaminoethyl-2-chlorides and trimethylsilyl derivatives of N, N-dialkylaminoethan-2-ols for verification of the Chemical Weapons Convention. *Rapid Commun. Mass Spectrom.* **20**, 2115–2119 (2006)
24. Yang, Y.C., Baker, J.A., Ward, J.R.: Decontamination of chemical warfare agents. *Chem. Rev.* **92**, 1729–1743 (1992)
25. Hooijschuur, E.W.J., Kientz, C.E., Brinkman, U.A.T.: Analytical separation techniques for the determination of chemical warfare agents. *J. Chromatogr. A* **982**, 177–322 (2002)
26. Black, R.M., Muir, B.: Derivatization reactions in the chromatographic analysis of chemical warfare agents and their degradation products. *J. Chromatogr. A* **1000**, 253–281 (2003)
27. Becke, A.D.: Density functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **98**, 5648–5652 (1993)
28. Lee, C., Yang, W., Parr, R.G.: Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B* **37**, 785–789 (1988)
29. Frisch, M.J., Trucks, G.W., Schlegel, H.B., Scuseria, G.E., Robb, Cheeseman, J.R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G.A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H.P., Izmaylov, A.F., Bloino, J., Zheng, G., Sonnenberg, J.L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, H., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery Jr., J.A., Peralta, J.E., Ogliaro, F., Bearpark, M., Heyd, J.J., Brothers, E., Kudin, K.N., Staroverov, V.N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J.C., Iyengar, S.S., Tomasi, J., Cossi, M., Rega, N., Millam, J.M., Klene, M., Knox, J.E., Cross, J.B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R.E., Yazyev, O., Austin, A.J., Cammi, R., Pomelli, C., Ochterski, J.W., Martin, R.L., Morokuma, K., Zakrzewski, V.G., Voth, G.A., Salvador, P., Dannenberg, J.J., Dapprich, S., Daniels, A.D., Farkas, Ö., Foresman, J.B., Ortiz, J.V., Cioslowski, J., Fox, D.J.: *Gaussian 09, Revision B.02*. Gaussian, Inc., Wallingford (2009)
30. Hay, P.J., Wadt, W.R.: Ab initio effective core potentials for molecular calculations. Potentials for K to Au including outermost core orbitals. *J. Chem. Phys.* **82**, 299–310 (1985)
31. McIver Jr., J.W., Komornicki, A.: Structure of transition state in organic reactions. General theory and an application to the cyclobutene-butadiene isomerization using a semi-empirical molecular orbital method. *J. Am. Chem. Soc.* **94**, 2625–2633 (1972)
32. Hehre, W.J., Radom, L., Schelyer, P. v. R., Pople, J.A.: *Ab Initio Molecular Orbital Theory*. Wiley: New York, p. 76 and references therein. (1986)
33. Harvey, D.J., Horning, M.G.: The mass spectra of trimethylsilyl derivatives of some alkyl and aminoalkyl phosphonates. *Org. Mass Spectrom.* **9**, 111–124 (1974)
34. Chai, Y., Xiong, X., Yue, L., Jiang, Y., Pan, Y., Fang, X.: Intramolecular halogen transfer via halonium ion intermediates in the gas phase. *J. Am. Soc. Mass Spectrom.* **27**, 161–167 (2016)
35. Weber, W.P., Felix, R.A., Willard, A.K.: Mass spectral rearrangements. Silyl McLafferty rearrangement. *J. Am. Chem. Soc.* **92**, 1420–1421 (1970)
36. Petersson, G.: A McLafferty-type rearrangement of a trimethylsilyl group in silylated hydroxyl carbonyl compounds. *Org. Mass Spectrom.* **6**, 577–592 (1972)
37. Arseniyadis, S., Goré, J., Guenot, P., Carrié, R.: Fragmentation behavior of  $\omega$ -functionalized allenes and their isomeric acetylenes under electron impact. Study on the isomerization of molecular ions and structural assignment of heteroatom migration related to a McLafferty rearrangement by CAD-MIKE spectrometry. *J. Chem. Soc. Perkin Trans. 2*, 1413–1418 (1985)
38. Hwu, J.R., Wu, I.-F., Shiao, S.-S., Tseng, W.N., Huang, J.J., Chen, B.L.: New fragmentation process in mass spectrometry of carbonyl compounds with a  $\delta$ -silyl group. *Organometallics* **18**, 2314–2320 (1999)
39. Rontani, J.-F., Aubert, C.: Hydrogen and trimethylsilyl transfers during EI mass spectral fragmentation of hydrocarboxylic and oxocarboxylic acid trimethylsilyl derivatives. *J. Am. Soc. Mass Spectrom.* **19**, 66–75 (2007)
40. NIST/EPA/NIH Mass spectral library with search program, Data version NIST 14: Gaithersburg, MD, USA
41. Gao, Y., Xu, J., He, Y., Tang, G., Lin, Z., Liu, H., Gao, X., Zhao, Y.: Cleavage of phosphorus-carbon (P-C) bonds of  $\alpha$ -amino phosphonates with intramolecular hydrogen migration in the gas phase using electrospray ionization tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **28**, 1964–1970 (2014)
42. Saeidian, H., Babri, M., Mirjafary, Z., Naseri, M.T., Sarabadani, M., Ashrafi, D., Faray, S.S.M.: Fragmentations mechanism in mass spectrometry of Chemical Weapons Convention related spiro alkylphosphonates and alkylidioxaphosphinane oxides. *Int. J. Mass Spectrom.* **369**, 59–70 (2014)
43. Wensing, M.W., Synder, A.P., Harden, C.S.: Energy resolved mass spectrometry of diethyl alkyl phosphonates via an atmospheric pressure ionization tandem mass spectrometer. *J. Mass Spectrom.* **30**, 1539–1545 (1995)
44. Glockler, G.: Carbon-halogen bond energies and bond distances. *J. Phys. Chem.* **63**, 828–832 (1959)
45. McLafferty, F.W.: Mass spectrometry analysis. Aliphatic halogenated compounds. *Anal. Chem.* **34**, 2–15 (1962)