

On the Borsuk conjecture concerning homotopy domination

R. Komendarczyk¹ · S. Kwasik¹ · W. Rosicki²

Received: 17 February 2015 / Accepted: 20 November 2015 / Published online: 26 December 2015
© Tbilisi Centre for Mathematical Sciences 2015

Abstract In the seminal monograph *Theory of retracts*, Borsuk raised the following question: suppose two compact ANR spaces are h -equal, i.e. mutually homotopy dominate each other, are they homotopy equivalent? The current paper approaches this question in two ways. On one end, we provide conditions on the fundamental group which guarantee a positive answer to the Borsuk question. On the other end, we construct various examples of compact h -equal, not homotopy equivalent continua, with distinct properties. The first class of these examples has trivial all known algebraic invariants (such as homology, homotopy groups etc.) The second class is given by n -connected continua, for any n , which are infinite CW -complexes, and hence ANR spaces, on a complement of a point.

Keywords ANR spaces · Homotopy domination · Homotopy type

Mathematics Subject Classification Primary 55P55; Secondary 55P15 · 54C56

Communicated by Jim Stasheff.

The R. Komendarczyk acknowledges the support of DARPA YFA N66001-11-1-4132 and NSF DMS 1043009. The S. Kwasik acknowledges the support of the Simons Foundation Grant No. 281810.

✉ R. Komendarczyk
rako@tulane.edu

S. Kwasik
kwasik@tulane.edu

W. Rosicki
wrosicki@mat.ug.edu.pl

¹ Department of Mathematics, Tulane University, New Orleans, USA

² Faculty of Mathematics, University of Gdansk, Gdansk, Poland

1 Introduction

Given two topological spaces X and Y , X is *homotopy dominated* by Y ; denoted by $X \leq_h Y$, if and only if there exist maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$, such that $g \circ f \simeq \text{id}_X$. If $X \leq_h Y$ and $Y \leq_h X$, the spaces X and Y are called *h -equal*, the latter denoted by $X =_h Y$. In particular if X is homotopy equivalent to Y , i.e. $X \simeq Y$, then they are *h -equal*. In the homotopy theory of Borsuk's ANR spaces, c.f. [5], two basic problems are raised. Paraphrasing Borsuk [5], the first one can be stated as follows:

(1) Is every compact ANR space homotopy equivalent to a finite CW-complex?

and the second one:

(2) Are two h -equal compact ANR spaces homotopy equivalent? In other words, given compact ANR spaces X and Y , does $X =_h Y$ imply $X \simeq Y$?

Both questions become less challenging if the compactness condition is relaxed, since the answer to the first question is positive [19], and negative for the second one, [22]. *Problem 1* (with the compactness assumption) became known as the *Borsuk conjecture* and attracted a considerable interest (c.f. [8, 15, 19, 21]) which culminated in the positive solution by West in [24]. In contrast, for the second question surprisingly little progress has been made over the years. One of the goals of the current paper is to renew interest in *Problem 2*.

The paper consists of essentially two parts. In the first part, which is mostly of expository nature, we make some comments on the role of the fundamental group in *Problem 2*. By analogy to Hopfian groups, we define a notion of a *Hopfian pair* for h -equal spaces and make the following observation.

Observation 1.1 *The pair of ANR spaces X, Y is a Hopfian pair, if and only if, X and Y are homotopy equivalent.*

This observation is in essence a reformulation of the classical Whitehead theorem, but it helps to put the Borsuk problem (*Problem 2*) in a proper perspective. In particular, it yields the following consequence

Theorem 1.2 *Suppose X , and Y are h -equal ANR spaces, such that $\pi_1(X)$ or $\pi_1(Y)$ is Hopfian, where X is compact, or more generally has finitely generated homology groups $H_k(X)$ for all k . If one of the group rings Λ is a Noetherian ring, then X and Y is a Hopfian pair, and hence X and Y are homotopy equivalent.*

The following corollary is well known to the experts [11, 16, 17] in this research area:

Corollary 1.3 *Suppose X and Y are compact, h -equal ANR spaces with the polycyclic-by-finite fundamental groups, then X and Y is a Hopfian pair.*

In Sect. 2, we also make several related observations in the context of Hopfian pairs, Poincaré complexes and H -spaces.

The second part, the main part of the paper, is where we construct 2-dimensional continua which are h -equal but not homotopy equivalent, see Theorem 3.5, these

constructions are inspired by [14] and [22]. A basic building block of these examples is a well known “topological broom” pictured on Fig. 1. An interesting feature of these constructions is that these spaces have trivial all basic known algebraic invariants, such as singular or Čech homology groups, homotopy groups etc. Consequently, to prove that the spaces are not homotopy equivalent requires a more direct, approach via techniques of set theoretic topology. Further, in Theorem 3.5, we provide examples of pairs $\mathcal{S}_0, \mathcal{S}_1$ of $2n$ -dimensional continua (for $n \geq 2$), modeled on the *Hawaiian earrings*, c.f [9], and satisfying:

- (a) $\mathcal{S}_0, \mathcal{S}_1$ are *singular ANR spaces*, i.e. for specific points $s_0 \in \mathcal{S}_0$, and $s_1 \in \mathcal{S}_1$, complements $\mathcal{S}_0^\circ = \mathcal{S}_0 - \{s_0\}$, $\mathcal{S}_1^\circ = \mathcal{S}_1 - \{s_1\}$ are countable disjoint¹ sums of connected ANR spaces.
- (b) $\mathcal{S}_0, \mathcal{S}_1$ are $(n - 1)$ -connected, and each connected component of \mathcal{S}_0° and \mathcal{S}_1° is locally contractible.
- (c) $\mathcal{S}_0 =_h \mathcal{S}_1$ but $\mathcal{S}_0 \not\cong \mathcal{S}_1$.

The point of this construction is to obtain examples of *compact* spaces which are as close as possible to ANR spaces. The construction is a generalization of the earlier result in [22] and relies on the fairly recent work in [9].

2 On a role of the fundamental group in Borsuk’s problem.

2.1 Hopfian pairs

We recall that a finitely presented group G is called *Hopfian*, if every epimorphism $h : G \rightarrow G$ is an isomorphism. Analogously, given a ring R , a finitely generated R -module M is *Hopfian*, if any module epimorphism $h : M \rightarrow M$ is an isomorphism. Let X and Y be a pair of h -equal spaces then, from definition, there are maps

$$\begin{aligned} f : X &\rightarrow Y, & i : Y &\rightarrow X, & f \circ i &\simeq \text{id}_Y, \\ g : Y &\rightarrow X, & j : X &\rightarrow Y, & g \circ j &\simeq \text{id}_X. \end{aligned} \tag{2.1}$$

In particular it implies that induced homomorphisms f_*, g_* on the fundamental group and homology groups, are epimorphisms.

Definition 2.1 A pair of spaces X, Y is called *Hopfian pair*, if and only if $X =_h Y$ and one of the epimorphisms $(g \circ f)_*$ or $(f \circ g)_*$ induced on the fundamental groups and homology modules from maps in (2.1) is an isomorphism.

Note that, in the above definition, if one of the epimorphisms is an isomorphism the second one is an isomorphism as well. For convenience, let us restate Observation 1.1:

Observation *The pair of ANR spaces: X, Y is Hopfian pair, if and only if, X and Y are homotopy equivalent.*

¹ A simple modification of this construction (see Sect. 3.2) yields a path connected complement of analogs of \mathcal{S}_0 and \mathcal{S}_1 .

Proof From the above definition, both compositions

$$\begin{aligned} g_* f_* &= (g \circ f)_* : \pi_1(X) \longrightarrow \pi_1(X), \\ f_* g_* &= (f \circ g)_* : \pi_1(Y) \longrightarrow \pi_1(Y), \end{aligned}$$

are isomorphisms. Thus, f_* and g_* are monomorphisms and consequently they have to be isomorphisms as well. The same reasoning applies to the module homomorphisms $f_* : H_k(X; \Lambda) \longrightarrow H_k(Y; \Lambda)$, $g_* : H_k(Y; \Lambda) \longrightarrow H_k(X; \Lambda)$, $\Lambda = \mathbb{Z}[\pi]$. As a consequence, maps f and g induce isomorphisms on π_1 , and all homology with local coefficients, and the Whitehead Theorem implies that f and g are homotopy equivalences. Let $f : X \longrightarrow Y$ be a homotopy equivalence with the inverse $g : Y \longrightarrow X$. Then, $g \circ f \simeq \text{id}_X$ and $f \circ g \simeq \text{id}_Y$, then obviously the pair X, Y is a Hopfian pair. \square

Remark 2.2 Note that in the above observation, it suffices to only have one ANR space; X or Y , then the result of Milnor [19], implies that the other space (homotopy dominated by the former) is also an ANR, up to homotopy.

Proof of Theorem 1.2 and Corollary 1.3 Since the group rings are Noetherian rings, and modules $H_*(X; \mathbb{Z}[\pi_1(X)])$ and $H_*(Y; \mathbb{Z}[\pi_1(Y)])$ are finitely generated, they in turn are Hopfian, c.f. [18]. Given that X and Y are h -equal they must form a Hopfian pair, implying Theorem 1.2. Since polycyclic-by-finite groups are Hopfian and their group rings Noetherian, Corollary 1.3 is a special case of Theorem 1.2.

The class of Hopfian groups is considerably larger than the polycyclic-by-finite groups. In particular, the following question is a weaker form of *Problem 2*.

Question 2.3 *Let X, Y be finite CW-complexes (compact ANR's) such that $X =_h Y$, suppose further $\pi_1(X)$ (and hence $\pi_1(Y)$) is Hopfian. Is X, Y a Hopfian pair?*

The following example, guided by the results of [2, 10], illustrates a delicate nature of the above question. Indeed, if $G = \pi_1(X) \cong \pi_1(Y)$ and G is Hopfian, it is not necessarily the case that $X =_h Y$, even for 2-dimensional CW-complexes X and Y .

Example 2.4 Let $G = \langle x, y \mid x^2 = y^3 \rangle$ be the standard presentation for the fundamental group of the trefoil knot, and let

$$G_i = \langle x, y, \bar{x}, \bar{y} \mid x^2 = y^3, \bar{x}^2 = \bar{y}^3, x^{2i+1} = \bar{x}^{2i+1}, y^{3i+1} = \bar{y}^{3i+1} \rangle, \quad i \in \mathbb{N},$$

be different presentations of G (c.f. [2, 10]). For infinitely many i , there are 2-dimensional CW-complexes K_i of distinct homotopy type, with $\pi_1(K_i) \cong G_i \cong G$, c.f. [10]. Note that the commutator subgroup $[G, G]$ of G is isomorphic to F_2 , i.e. free group on two generators, and $G/[G, G] \cong H_1(G; \mathbb{Z}) \cong \mathbb{Z}$.

Also, G is Hopfian, since both $[G, G] \cong F_2$ and $G/[G, G] \cong \mathbb{Z}$ are Hopfian and it is well known that G is not polycyclic-by-finite. We claim that there are infinitely many pairs i and $j, i \neq j$, such that $K_i \neq_h K_j$.

Recall that it is shown in [2, 10] that there are infinitely many pairs i and $j, i \neq j$, such that K_i is not homotopy equivalent to K_j , because $H_2(\tilde{K}_i; \mathbb{Z})$ and $H_2(\tilde{K}_j; \mathbb{Z})$ are

not isomorphic as $\mathbb{Z}[G]$ -modules. More precisely, for some prime number p , there are infinitely many distinct i and j such that $\mathbb{Z}_p \otimes_{\mathbb{Z}} H_2(\tilde{K}_i; \mathbb{Z})$ has just one generator and $\mathbb{Z}_p \otimes_{\mathbb{Z}} H_2(\tilde{K}_j; \mathbb{Z})$ has at least two generators. Suppose $K_i =_h K_j$, for $i \neq j$, where the above holds, then one obtains an epimorphism $f_* : H_2(K_i; \mathbb{Z}[G]) \rightarrow H_2(K_j; \mathbb{Z}[G])$ (see Definition 2.1), and therefore an obvious epimorphism

$$\text{id} \otimes f_* : \mathbb{Z}_p \otimes_{\mathbb{Z}} H_*(K_i; \mathbb{Z}[G]) \rightarrow \mathbb{Z}_p \otimes_{\mathbb{Z}} H_*(K_j; \mathbb{Z}[G]),$$

this however contradicts that $H_*(K_i; \mathbb{Z}[G])$ has just one generator versus $H_*(K_j; \mathbb{Z}[G])$ having two generators. Thus by contradiction, we conclude that $K_i \neq_h K_j$ and hence the pair K_i and K_j is not a Hopfian pair.

The above considerations lead one to a surprising outcome when one considers spaces $K_i \vee S^2$ and $K_j \vee S^2$ in place of K_i and K_j . By work in [2, 10] we know that

$$K_i \vee S^2 \simeq K_j \vee S^2, \quad \text{thus} \quad K_i \vee S^2 =_h K_j \vee S^2.$$

Note that $G \cong \pi_1(K_i \vee S^2) \cong \pi_1(K_j \vee S^2)$, and the pair $K_i \vee S^2, K_j \vee S^2$ is Hopfian. Indeed the modules $H_2(K_i \vee S^2; \mathbb{Z}[G]), H_2(K_j \vee S^2; \mathbb{Z}[G])$ are Hopfian as both are isomorphic to the free $\mathbb{Z}[G]$ -module $\mathbb{Z}[G] \oplus \mathbb{Z}[G]$ (c.f. [2]). This shows a difficulty in dealing with modules $H_*(X; \mathbb{Z}[G])$ and $H_*(Y; \mathbb{Z}[G])$, in the context of Question 2.3, even if the group $\pi_1(X) \cong \pi_1(Y)$ is a “nice” group.

2.2 Poincaré complexes

Now, let M^n be a closed n -dimensional manifold and Y any space (see Remark 2.2), such that $M^n =_h Y$, then M^n, Y is a Hopfian pair [3, 17]. More generally, let X be a finite Poincaré complex of formal dimension n , c.f. [23]. To be specific, X has a homotopy type of a finite CW-complex and there exists a class $[X] \in H_n(X; \mathbb{Z})$, such that for all r the cap product with $[X]$ induces an isomorphism

$$[X] \cap \cdot : H^r(X; \Lambda) \rightarrow H_{n-r}(X; \Lambda), \quad \Lambda = \mathbb{Z}[\pi_1(X)].$$

If Y is any space, such that $X =_h Y$ then X, Y is a Hopfian pair, [17].

Theorem 2.5 *Suppose X is a homology manifold of formal dimension n , i.e. X is a finite dimensional ANR space such that*

$$H_*(X, X - \{pt\}) \cong H_*(\mathbb{R}^n, \mathbb{R}^n - \{pt\}) = \begin{cases} \mathbb{Z}, & * = n, \\ 0, & * \neq n. \end{cases}$$

Then X is a finite Poincaré complex of formal dimension n .

The above theorem is stated without a proof in [13, p. 5099]. It is a well known fact that X satisfies the Poincaré duality with integer coefficients, [4]. The only argument we are aware of, that shows X is a Poincaré complex, is based on the existence of a spectral sequence for the identity map $\text{id}_X : X \rightarrow X$ in sheaf homology giving a

very general version of Poincaré duality in Theorem 9.2 of [7]. It should be noted that if X is *polyhedral homology manifold* then a much simpler argument shows that X is a Poincaré complex (see Theorem 2.1 in [23]).

Corollary 2.6 *Let X be a homology manifold of formal dimension n and Y any space with $X =_h Y$, then X, Y is a Hopfian pair.*

Recall, that the well known conjecture asserts that finite dimensional homogeneous ANRs are homology manifolds [8].

Following [6], recall that X is *locally isotopic* if for each path $\lambda : [0, 1] \rightarrow X$, there is a neighborhood N of $\lambda(0)$ in X and a map $H : I \times N \rightarrow X$, such that $H(t, \lambda(0)) = \lambda(t)$ and such that each $H(t, \cdot)$ is a homeomorphism of N onto a neighborhood of $\lambda(t)$. Clearly, manifolds are locally isotopic. Suppose X is a compact finite dimensional ANR space which is locally isotopic. By Theorem 4.6 of [6], X is a homology manifold of some formal dimension n . Therefore, we obtain

Corollary 2.7 *Let X be a compact finite dimensional ANR space which is locally isotopic, and let Y any space such that $X =_h Y$. Then X, Y is a Hopfian pair.*

In the case X admits an H -space structure, $\pi_1(X)$ is abelian, in particular polycyclic-by-finite, thus if $H_*(X)$ to be finitely generated in each degree (where we allow the degree to go to infinity), we obtain

Proposition 2.8 *Let X be an H -space, such that $H_k(X)$ is finitely generated for each k , and Y any space such that $X =_h Y$. Then X, Y is a Hopfian pair.*

Clearly, if X is a compact H -space the above homological condition holds. Curiously enough, compact H -spaces are also Poincaré complexes, as can be deduced from the work in [1].

3 About h -equal but not homotopy equivalent spaces

Looking for a counterexample to *Problem 2*, one may consider the following problem in the combinatorial group theory; suppose G and H , $G \not\cong H$ are two finitely presented groups and retracts of each other, which would make such pair of groups “strongly” non-Hopfian. If both G and H are finite dimensional, i.e. $K(G; 1)$ and $K(H; 1)$ are chosen to be finite CW -complexes, then the functoriality of the construction of $K(\pi; 1)$ -spaces would imply the existence of a counterexample to *Problem 2*, namely

$$K(G; 1) =_h K(H; 1), \quad \text{and} \quad K(G; 1) \not\cong K(H; 1).$$

Consequently, the following algebraic question is of crucial importance and of an independent interest.

Question 3.1 *Find two finitely presented groups G and H , such that $G \not\cong H$ which are retracts of each other.*

If one considers a more general class of spaces, then the answer to *Problem 2* is negative, as first observed by Stewart in [22], who provided examples of noncompact ANR spaces. The remainder of this paper is devoted to a construction of compact examples with particular properties as described in the introduction, Sect. 1.




Fig. 1 Topological broom denoted by \mathcal{B} , with the center point v , and a location of the point u , and its neighborhood U , considered in the proof of Lemma 3.8

3.1 Infinite wedges of “hairy disks”

Our example is inspired by constructions of both [14,22], and based on the “hairy disk” depicted in Fig. 2. First, consider a double broom \mathcal{B} as shown on Fig. 1. \mathcal{B} is a well known space which is not contractible but has all trivial known algebraic invariants, such as homology and homotopy groups etc. [12, p. 295]. Denote the *center point* of the broom \mathcal{B} by v and the left and right sequence of broom’s endpoints converging to v by $\{a_n\}$ and $\{b_n\}$ respectively. Generally, J_x , provided it is uniquely determined, will refer to a segment of \mathcal{B} containing $x \in \mathcal{B}$. An exception to this are the following cases: for $x = v, a_0, b_0$, we set


$$J_v = [v, a_0] \cup [v, b_0], \quad J_{a_0} = [v, a_0], \quad J_{b_0} = [v, b_0]. \tag{3.1}$$

In particular,

$$J_{a_n} = [a_n, a_0], \quad J_{b_n} = [b_n, b_0]. \tag{3.2}$$

Naturally, we may view \mathcal{B} as a wedge product of two pieces A and B , containing sequences $\{a_n\}$ and $\{b_n\}$, i.e.

Fig. 2 The “hairy disk” \mathcal{H} from [14, p. 286], with countably many copies of \mathcal{B} densely attached along the boundary of the unit disk in \mathbb{R}^2



$$\mathcal{B} = A \vee B, \quad A = J_{a_0} \cup \bigcup_n J_{a_n}, \quad B = J_{b_0} \cup \bigcup_n J_{b_n}. \tag{3.3}$$

Further, we order points in \mathcal{B} along segments J_x ; simply assuming the order is “increasing” from the bottom to top, for instance any $z \in J_{a_n}$ satisfies $a_n \leq z \leq a_0$. In particular, if $x, y \in J_w$, and $x \leq y$ with respect to this order, then $[x, y]$ will denote a portion of the segment J_w containing all z such that $x \leq z \leq y$.

Definition 3.2 Given a space X , we say $x \in X$ is *homotopically fixed in X* , if x is fixed under any homotopy f_t , where $f_0 = \text{id}_X$. The set of homotopically fixed points in X is denoted by

$$hf(X) = \{x \in X \mid x \text{ is homotopically fixed in } X\}. \tag{3.4}$$

The set $hf(X)$ is a closed subset of X , in particular we have the following fact about \mathcal{B} :

Lemma 3.3 *Suppose $f : \mathcal{B} \rightarrow \mathcal{B}$ fixes v , i.e. $f(v) = v$, and v is a limit point for both sets: $Z \cap A$ and $Z \cap B$, $Z = f(\mathcal{B})$. Then, any homotopy $f_t : \mathcal{B} \rightarrow \mathcal{B}$, $f_0 = f$ keeps v fixed, i.e. $f_t(v) = v$.*

Proof (Sketch of Proof) Choose sequences $\{u_n\}, u_n \in J_{a_n}$ in $Z \cap A$ and $\{w_n\}, w_n \in J_{b_n}$ in $Z \cap B$ respectively, such that

$$u_n \rightarrow v, \quad \text{and} \quad w_n \rightarrow v \quad \text{in } \mathcal{B}.$$

Let $v_t = f_t(v)$, by continuity, for each t :

$$f_t(u_n) \rightarrow v_t, \quad \text{and} \quad f_t(w_n) \rightarrow v_t.$$

Note that a point u_n can only move up along the arm $J_{u_n} \subset A$ of \mathcal{B} and w_n move up along $J_{w_n} \subset B$ (for large enough n). Thus $v_t = v_0$ for all t , because $A \cap B = \{v\}$. □

The above lemma is completely analogous to [14, Lemma 2.3], where a similar topological broom is considered.²

Corollary 3.4 $hf(\mathcal{B}) = \{v\}$.

Proof We already know that $v \in hf(\mathcal{B})$ by Lemma 3.3. It is easy to rule out other points in \mathcal{B} as homotopically fixed, with an exception of possibly a_0 and b_0 . Observe however, that a_0 and b_0 cannot be homotopically fixed as we may construct a homotopy which lets a_0 or b_0 to “flow out” along one of the arms of \mathcal{B} , e.g J_{a_1} and J_{b_1} respectively. □

² We choose the broom \mathcal{B} , shown on Fig. 1, over the one constructed in [14] to simplify certain arguments of this section.

Before introducing relevant spaces we make the following convenient definition of a *wedge product* Υ of spaces X and Y disjointly embedded in \mathbb{R}^N (for some N):

$$X \Upsilon_{x,y} Y := X \cup [x, y] \cup Y, \quad x \in X, y \in Y, \tag{3.5}$$

where $[x, y]$ is an arc in \mathbb{R}^N connecting points x and y , with its interior (x, y) disjoint from X and Y .

The disk \mathcal{H} is constructed by densely attaching brooms \mathcal{B} , Fig. 1, along the boundary of the unit disk D^2 in \mathbb{R}^2 , c.f. [14]. More precisely, let

$$M = \{m_i\}_{i=1}^\infty \tag{3.6}$$

to be a countable dense subset of the boundary of D^2 . Then \mathcal{H} is obtained by attaching to each point m_i the broom \mathcal{B} at the center vertex v , so that the copies of \mathcal{B} do not intersect each other and their diameters tend to zero as $i \rightarrow \infty$. Denote by c the center of interior disk D^2 in \mathcal{H} . Following the ideas of [22] we consider a countable wedge product of \mathcal{H} , with center points at $c(k) = (-\frac{1}{k}, 0, 0)$ along the x -axis of \mathbb{R}^3 . Each copy of \mathcal{H} is denoted by $\mathcal{H}(k)$ and contained in the translated yz -plane to the point $c(k)$, together with the connecting segments $[c(k), c(k+1)]$ along the x -axis. In addition, each factor $\mathcal{H}(k)$ is scaled down to have the diameter $\frac{1}{k}$. This process yields a non-compact space we denote by \mathcal{WH}° . The second space, denoted by \mathcal{WBH}° is obtained from \mathcal{WH}° by wedging a copy of \mathcal{B} at the first factor. Using the notation in (3.5), we express these spaces as follows

$$\begin{aligned} \mathcal{WH}^\circ &= \mathcal{H}(1) \Upsilon_{c(1),c(2)} \mathcal{H}(2) \Upsilon_{c(2),c(3)} \dots \Upsilon_{c(k-1),c(k)} \mathcal{H}(k) \Upsilon_{c(k),c(k+1)} \dots, \\ \mathcal{WBH}^\circ &= \mathcal{B} \Upsilon_{b,c'(1)} \mathcal{H}'(1) \Upsilon_{c'(1),c'(2)} \dots \Upsilon_{c'(k-1),c'(k)} \mathcal{H}'(k) \Upsilon_{c'(k),c'(k+1)} \dots, \end{aligned} \tag{3.7}$$

where b denotes the center point v of the \mathcal{B} factor, and \mathcal{H} -factors of \mathcal{WBH}° are denoted by $\mathcal{H}'(k)$ for clarity. The countable dense subset M of \mathcal{H} given in (3.6) will be further denoted by $M(k)$ for each $\mathcal{H}(k)$ in \mathcal{WH}° , and by $M'(k)$ for each $\mathcal{H}'(k)$ in \mathcal{WBH}° . Including the origin of \mathbb{R}^3 in both \mathcal{WH}° and \mathcal{WBH}° yields

$$\mathcal{WH} = \mathcal{WH}^\circ \cup \{(0, 0, 0)\}, \quad \mathcal{WBH} = \mathcal{WBH}^\circ \cup \{(0, 0, 0)\}. \tag{3.8}$$

Observe that \mathcal{WH} and \mathcal{WBH} are homeomorphic with one-point compactifications of \mathcal{WH}° and \mathcal{WBH}° . We claim \mathcal{WH}° and \mathcal{WBH}° are h -equal, in particular there exist retractions³

$$r_{\mathcal{WH}} : \mathcal{WBH}^\circ \longrightarrow \mathcal{WH}^\circ, \quad r_{\mathcal{WBH}} : \mathcal{WH}^\circ \longrightarrow \mathcal{WBH}^\circ. \tag{3.9}$$

³ Thus, \mathcal{WH}° and \mathcal{WBH}° are r -equal, c.f. [5].




Fig. 3 Infinite wedge products of hairy disks: $WB\mathcal{H}^\circ$ (left) and $W\mathcal{H}^\circ$ (right) embedded in \mathbb{R}^3 . The segment connecting $c(1)$ and m_k , in the right picture, indicates a choice of embedding $WB\mathcal{H}^\circ \hookrightarrow W\mathcal{H}^\circ$

Since $W\mathcal{H}$ is naturally a subset of $WB\mathcal{H}$, $r_{W\mathcal{H}}$ can be chosen as a quotient projection mapping the \mathcal{B} -factor, together with the segment $[b, c'(1)]$, to the point $c(1) \in \mathcal{H}(1)$ of $W\mathcal{H}^\circ$. The retraction $r_{WB\mathcal{H}}$ of $W\mathcal{H}^\circ$ onto $WB\mathcal{H}^\circ$ can be defined once we choose an embedding $\iota : WB\mathcal{H}^\circ \hookrightarrow W\mathcal{H}^\circ$. Once a point $m_k \in M(1)$ is selected, the embedding ι can be chosen to map the \mathcal{B} -factor of $WB\mathcal{H}$ to the factor \mathcal{B}_{m_k} of $\mathcal{H}(1)$ and identifying the segment $[b, c'(1)]$ with the segment $[m_k, c(1)]$ in $\mathcal{H}(1)$, as shown on Fig. 3(right). Having identified $WB\mathcal{H}^\circ$ with a subset of $W\mathcal{H}^\circ$ we may define the retraction $r_{WB\mathcal{H}}$ next. It will be done in two stages; first we retract all the broom factors of $\mathcal{H}(1)$, except \mathcal{B}_{m_k} to the boundary circle of the interior disk D^2 in $\mathcal{H}(1)$ via the following obvious map

$$z : W\mathcal{H}^\circ \longrightarrow W\mathcal{H}^\circ, \quad z(x) = \begin{cases} m_i & x \in \mathcal{B}_{m_i} \subset \mathcal{H}'(1), \quad i \neq k \\ x, & \text{otherwise.} \end{cases}$$

Continuity of z is a direct consequence of the ‘‘hairy disk’’ construction. Indeed given a convergent sequence of points $\{h_n\}$ in the complement of \mathcal{B}_{m_k} , i.e. $\{h_n\} \subset \mathcal{H}(1) - \mathcal{B}_{m_k}$. If the limit of $\{h_n\}$ is in \mathcal{B}_{m_k} then it has to be m_k , proving continuity of z . For the second stage, in the construction of $r_{WB\mathcal{H}}$, we define a map $y : z(W\mathcal{H}^\circ) \longrightarrow WB\mathcal{H}^\circ$ on the image of z , simply by collapsing the interior disk $D^2 \subset \mathcal{H}(1)$ onto the segment $[m_k, c(1)]$ in $\mathcal{H}(1)$. The required retraction $r_{WB\mathcal{H}}$ can be now defined as $r_{WB\mathcal{H}} = y \circ z$.

Theorem 3.5 *Both pairs: $W\mathcal{H}^\circ, WB\mathcal{H}^\circ$ and $W\mathcal{H}, WB\mathcal{H}$ are h-equal but not homotopy equivalent.*

The proof requires the following two lemmas.

Lemma 3.6 *We have the following homeomorphisms*

$$\begin{aligned} hf(W\mathcal{H}^\circ) &\cong \bigsqcup_{i=1}^\infty hf(\mathcal{H}(k)), & hf(WB\mathcal{H}^\circ) &\cong \{b\} \sqcup \bigsqcup_{i=1}^\infty hf(\mathcal{H}'(k)), \\ hf(W\mathcal{H}) &\cong hf(W\mathcal{H}^\circ) \sqcup \{(0, 0, 0)\}, & hf(WB\mathcal{H}) &\cong hf(WB\mathcal{H}^\circ) \sqcup \{(0, 0, 0)\}, \end{aligned} \tag{3.10}$$

where each $hf(\mathcal{H}(k))$ or $hf(\mathcal{H}(k))$ is just a boundary of the interior disk D^2 in each factor $\mathcal{H}(k)$ of \mathcal{WH}° (\mathcal{WH}) or $\mathcal{H}'(k)$ in \mathcal{WBH}° (\mathcal{WBH}), and therefore homeomorphic to S^1 . The topology is the subspace topology induced from \mathbb{R}^3 via the embeddings constructed in (3.7).

Corollary 3.7 *In particular, $hf(\mathcal{WH}^\circ)$ ($hf(\mathcal{WH})$) is not homeomorphic to $hf(\mathcal{WBH}^\circ)$ ($hf(\mathcal{WBH})$).*

Proof of Lemma 3.6 Lemma 3.4 and the construction of the hairy disk \mathcal{H} imply⁴

$$M \subset hf(\mathcal{H}).$$

Since M is dense in the boundary $S^1 = \partial D^2 \subset \mathcal{H}$, we obtain

$$S^1 = \overline{M} \subset hf(\mathcal{H}).$$

Since, none of the interior points in $D^2 \subset \mathcal{H}$ is homotopically fixed, and by Lemma 3.4, for each \mathcal{B}_{m_k} -factor of \mathcal{H} , m_k is the only homotopically fixed point of \mathcal{B}_{m_k} , we conclude

$$hf(\mathcal{H}) = S^1.$$

It in turn implies equalities in (3.10), note that $\{(0, 0, 0)\}$ is homotopically fixed as a limit of points in $hf(\mathcal{H}(k))$ fixed points from the \mathcal{H} -factors of \mathcal{WH} or \mathcal{WBH} .

Further, we obtain the following key lemma,

Lemma 3.8 *Let f be the homotopy equivalence between \mathcal{WH}° , and \mathcal{WBH}° , and g its inverse. Then,*

$$f(hf(\mathcal{WH}^\circ)) \subset hf(\mathcal{WBH}^\circ), \quad g(hf(\mathcal{WBH}^\circ)) \subset hf(\mathcal{WH}^\circ). \tag{3.11}$$

The same inclusions holds for the compactifications: \mathcal{WH} and \mathcal{WBH} .

Proof We will prove the first inclusion in (3.11), as the proof of the second is analogous. It suffices to prove for each k :

$$f(M(k)) \subset hf(\mathcal{WBH}^\circ). \tag{3.12}$$

Then the claim follows from continuity of f , and the fact that the closure of $\bigcup_k M(k)$ in \mathcal{WH}° is equal to $hf(\mathcal{WH}^\circ)$ (see Lemma 3.6). (Note that for the second inclusion in (3.11), the only difference is the point b (the center of the first broom factor of \mathcal{WBH}°) which needs to be added to the union $\bigcup_k M(k)$).

To prove (3.12), consider a point v in $M(k)$. By definition it has to be the center point of one of a broom factors in $\mathcal{H}(k) \subset \mathcal{WH}^\circ$, see (3.7). We further denote this factor by

⁴ It is easy to see that broom centers along \mathcal{H} -factors cannot be moved, by a homotopy, to the interior of the disk $D^2 \subset \mathcal{H}$, c.f. [14, (v) on p. 288].

\mathcal{B} (i.e. $v \in \mathcal{B} \subset \mathcal{H}(k)$). Let $u = f(v)$, and suppose by contradiction $u \notin hf(\mathcal{WBH}^\circ)$, then

- either 1° , \mathcal{WBH}° is locally path connected at u ;
- or 2° , \mathcal{WBH}° is not locally path connected at u . □

Observation 3.13 Since $v \in hf(\mathcal{WH})$, we must have $g \circ f(v) = v$ (as $g \circ f \simeq \text{id}_{\mathcal{WH}}$). Consider sequences $a_n \rightarrow v, b_n \rightarrow v$ of points in \mathcal{B} (see Fig. 1). Denote by $\tilde{a}_n = g \circ f(a_n), \tilde{b}_n = g \circ f(b_n)$, then we have $\tilde{a}_n \rightarrow v$ and $\tilde{b}_n \rightarrow v$. We claim that for large n :

$$\tilde{a}_n \in J_{a_n}, \quad \tilde{b}_n \in J_{b_n}. \tag{3.13}$$

Proof Indeed, denoting the homotopy $g \circ f \simeq \text{id}_{\mathcal{WH}}$ by $h_t = h(t, \cdot)$, $h : I \times \mathcal{WH} \rightarrow \mathcal{WH}$ we observe that for every n : $\gamma_{a_n}(t) = h_t(a_n)$ defines a path in \mathcal{WH} connecting $a_n = \gamma_{a_n}(1)$ and $\tilde{a}_n = \gamma_{a_n}(0) = g \circ f(a_n)$ (analogously for the sequence $\{b_n\}$). Since for the limit point $v = \lim a_n, \gamma_v$ is a constant path, for a small ε -ball $B_v(\varepsilon)$ around v , the inverse image $h^{-1}(B_v(\varepsilon)) \subset I \times \mathcal{WH}$ contains $I \times \{v\}$ and therefore some small neighborhood $I \times B_v(\delta)$ is also in $h^{-1}(B_v(\varepsilon))$. For large enough n, a_n 's are in $B_v(\delta)$ and hence the paths γ_{a_n} have image in $B_v(\varepsilon)$. It follows that each γ_{a_n} is contained in the connected component $J_{a_n} \cap B_v(\varepsilon) \subset \mathcal{B}$ of $B_v(\varepsilon)$. Hence, for small positive ε we obtain $J_{a_k} \cap J_{a_j} \cap B_v(\varepsilon) = \emptyset$ and the first part of (3.13). The second part follows analogously. □

Now we consider Case 1° and Case 2° .

Case 1° : Suppose \mathcal{WBH}° is locally path connected at $u = f(v)$. Choose a small path connected ball $B_u(\tilde{\varepsilon})$ around u , such that $f(B_v(\delta)) \subset B_u(\tilde{\varepsilon})$ then $g(B_u(\tilde{\varepsilon})) \subset B_v(\varepsilon)$ with δ and ε chosen as in the proof of Observation (3.13) above. Since $g(B_u(\tilde{\varepsilon}))$ is connected, and all $\{\tilde{a}_n\}$ for large n are contained in $g(B_u(\tilde{\varepsilon}))$, $\{\tilde{a}_n\}$ would have to belong entirely to one of the arms $J_{a_k} \cap B_v(\varepsilon)$ of \mathcal{B} . But, this leads to a contradiction with Observation (3.13).

Case 2° : Suppose \mathcal{WBH}° is not locally connected at $u = f(v)$. Since $u \notin hf(\mathcal{WBH}^\circ), u$ belongs to one of the broom factors of \mathcal{WBH}° , we denote by \mathcal{B}' (i.e. \mathcal{B}' is either the \mathcal{B} -factor of \mathcal{WBH}° or belongs to one of the $\mathcal{H}'(k)$ -factors). We also endow \mathcal{B}' with decorations of Fig. 1, where v' stands for the center of \mathcal{B}' , and a'_n, b'_n correspond to a_n and b_n , etc. Note that the set of points where \mathcal{B}' is not locally path connected is given by $V' = J_{v'} - (\{a'_0\} \cup \{b'_0\})$ and therefore $u \in V'$. Since $u \notin hf(\mathcal{WBH}^\circ)$ and also $u \neq v'$, without loss of generality, we assume $u \in J_{a'_0} - \{a'_0\}$. Further, continuity of f implies $f(a_n) \rightarrow u$ and $f(b_n) \rightarrow u$ and for large n , both sequences $\{f(a_n)\}$ and $\{f(b_n)\}$ belong to a small neighborhood U of u consisting of infinitely many disjoint segments accumulating on $J_{v'} \cap U$ (see Fig. 1 for the illustration). Consider the shortest piece-wise linear paths $\alpha_n : I \rightarrow \mathcal{B}'$, joining $\alpha_n(0) = f(a_n) \in U$ and $\alpha_n(1) = u; \beta_n : I \rightarrow \mathcal{B}'$, joining $\beta_n(0) = f(b_n) \in U$ and u . Clearly, both α_n and β_n trace segments respectively:

$$\alpha_n = [f(a_n), a'_0] \cup [u, a'_0] \subset \mathcal{B}', \quad \beta_n = [f(b_n), a'_0] \cup [u, a'_0] \subset \mathcal{B}'$$

(we identify α_n and β_n with their images for simplicity). In turn, the paths $g \circ \alpha_n$ and $g \circ \beta_n$, join points $\tilde{a}_n = g(f(a_n))$ and $v = g(f(u))$, see Eq. (3.13). Since points \tilde{a}_n

(resp. \tilde{b}_n) are close to a_n (resp. b_n), and belong to J_{a_n} (resp. J_{b_n}). The image of $g \circ \alpha_n$ contains $[\tilde{a}_n, a_0] \cup [v, a_0]$ and the image of $g \circ \beta_n$ contains segments $[\tilde{b}_n, b_0] \cup [v, b_0]$. Therefore, for n large enough, we can find $s_n \in \alpha_n$, and $t_n \in \beta_n$, such that

$$g(s_n) = a_0, \quad g(t_n) = b_0. \tag{3.14}$$

Moreover, for each n we can choose minimal such s_n and t_n (i.e. closest to the initial point of the paths α_n and β_n). Passing to subsequences, if necessary, we have $s_n \rightarrow s$, $t_n \rightarrow t$, and both limits belong to $J_{a'_0}$. Clearly,

$$g(s) = a_0, \quad g(t) = b_0. \tag{3.15}$$

By (3.14), we have $s \neq t$, and both s and t are above u , i.e. $s > u$ and $t > u$, according to the order defined after Eq. (3.3).

Suppose $s > t > u$: the initial points $f(a_n)$ of α_n converge to u , and $s_n \in \alpha_n$ (or a subsequence) converges to s as $n \rightarrow \infty$. Since, $s > t$ we can find points $\{e_n\}$ with $e_n \in \alpha_n$ and

$$f(a_n) \leq e_n \leq s_n,$$

(as points ordered along α_n) and such that $e_n \rightarrow t$, as a consequence $g(e_n) \rightarrow g(t) = b_0$. However, s_n is the first point on α_n mapped to a_0 under g . Further $f(a_n)$, the initial point of α_n , is mapped to \tilde{a}_n . By (3.13), we conclude that $g(e_n) \in J_{a_n}$ for large enough n , and therefore the limit of $\{g(e_n)\}$ has to belong to J_{a_0} , contradicting the fact that $b_0 \notin J_{a_0}$.

In the case $t > s > u$, analogously considering paths β_n , we may find a sequence of points $\{h_n\}$, converging to s , and such that for large n :

$$f(b_n) \leq h_n \leq t_n.$$

Then, again points $g(h_n)$ can only accumulate on J_{b_0} , contradicting

$$g(h_n) \rightarrow g(s) = a_0 \notin J_{b_0}.$$

This proves (3.12) and concludes the proof of (3.11) for \mathcal{WH}° and \mathcal{WBH}° . To prove the inclusion for the compactifications \mathcal{WH} and \mathcal{WBH} , we see that Claim (3.12) follows because the point at ∞ , i.e. $\{(0, 0, 0)\}$, is in the closure of $\bigcup_k M(k)$. \square

Proof of Theorem 3.5 Suppose that \mathcal{WH}° and \mathcal{WBH}° are homotopy equivalent and take a homotopy equivalence $f : \mathcal{WH}^\circ \rightarrow \mathcal{WBH}^\circ$, with inverse $g : \mathcal{WBH}^\circ \rightarrow \mathcal{WH}^\circ$ (in the notation of Lemma 3.8.) Their restrictions to the sets of homotopically fixed points are defined by

$$\tilde{f} = f|_{h_f(\mathcal{WH}^\circ)}, \quad \tilde{g} = g|_{h_f(\mathcal{WBH}^\circ)}$$

By Lemma 3.8, compositions $\tilde{g} \circ \tilde{f}$ and $\tilde{f} \circ \tilde{g}$ are well defined and by Definition 3.2 they satisfy

$$\tilde{g} \circ \tilde{f} = \text{id}_{hf(\mathcal{WH}^\circ)}, \quad \tilde{f} \circ \tilde{g} = \text{id}_{hf(\mathcal{WBH}^\circ)}.$$

Thus \tilde{f} , defines a homeomorphism between $hf(\mathcal{WH}^\circ)$ and $hf(\mathcal{WBH}^\circ)$, and \tilde{g} its inverse, contradicting Lemma 3.6. For one-point compactifications: \mathcal{WH} and \mathcal{WBH} , points at infinity are homotopically fixed therefore the statement follows analogously.

3.2 Infinite wedges of products of n -spheres

For arbitrarily high connected examples, we follow a similar pattern as in the previous section. Let \mathcal{S} be an n -sphere, and $\mathcal{S}^2 = \mathcal{S} \times \mathcal{S}$, define the following countable wedge products at a common basepoint s :

$$\mathcal{S}_0 = \mathcal{S}^2 \vee \mathcal{S}^2 \vee \dots \vee \mathcal{S}^2 \vee \dots = \bigvee_{j=1}^\infty \mathcal{S}_0(j), \quad \mathcal{S}_1 = \mathcal{S} \vee \mathcal{S}_0 = \bigvee_{j=1}^\infty \mathcal{S}_1(j). \quad (3.16)$$

Thus \mathcal{S}_0 and \mathcal{S}_1 differ just by the first factor, further we consider both \mathcal{S}_0 and \mathcal{S}_1 to be metrically embedded in \mathbb{R}^{2n+2} , with the basepoint at the origin, and the diameters of factors $\mathcal{S}_*(j)$ tending to zero as $j \rightarrow \infty$, see Fig. 4. Both \mathcal{S}_0 and \mathcal{S}_1 are compact in the topology induced from the embedding. Clearly, \mathcal{S}_0 and \mathcal{S}_1 are $(n - 1)$ -connected in this topology since each factor is $(n - 1)$ -connected. Further, let \mathcal{S}_* stand for either \mathcal{S}_0 or \mathcal{S}_1 .

The obvious retraction $r_{\mathcal{S}} : \mathcal{S} \times \mathcal{S} \rightarrow \mathcal{S}$, can be extended by the identity (and rescaling) to a retraction $r_{\mathcal{S}_1} : \mathcal{S}_0 \rightarrow \mathcal{S}_1$. A retraction $r_{\mathcal{S}_0} : \mathcal{S}_1 \rightarrow \mathcal{S}_0$ can be obtained by simply collapsing the \mathcal{S} -factor of \mathcal{S}_1 to the basepoint. Thus we obtain

$$\mathcal{S}_0 =_h \mathcal{S}_1, \quad \mathcal{S}_0 =_h \mathcal{S}_1.$$

The strategy for proving that \mathcal{S}_0 and \mathcal{S}_1 are not homotopy equivalent is a little different than before and based on some homological considerations.

Both \mathcal{S}_0 and \mathcal{S}_1 are a special case of the generalized Hawaiian earrings construction considered in [9]. Following [9] consider the following homomorphisms defined on $\pi_*(\mathcal{S}_*, s)$ in dimension n (for $n > 1$):




Fig. 4 Hawaiian earrings \mathcal{S}_0 (left) and \mathcal{S}_1 (right) based on $\mathcal{S} \times \mathcal{S}$ and \mathcal{S} , for $n = 1$

$$h_0 : \pi_n(\mathcal{S}_0, s) \longrightarrow \prod_{j=1}^{\infty} \pi_n(\mathcal{S}_0(j), s), \quad h_1 : \pi_n(\mathcal{S}_1, s) \longrightarrow \prod_{j=1}^{\infty} \pi_n(\mathcal{S}_1(j), s),$$

and induced by the product of obvious coordinate retractions $r_{j,*} : \mathcal{S}_* \longrightarrow \mathcal{S}_*(j)$ onto each factor of \mathcal{S}_* . The main theorem of [9, p. 18] implies that both h_0 and h_1 are isomorphisms. By the Hurewicz Theorem

$$\begin{aligned}
 H_n(\mathcal{S}_0) &\cong \pi_n(\mathcal{S}_0, s) \cong \prod_{j=1}^{\infty} H_n(\mathcal{S}_0(j); \mathbb{Z}) \cong \prod_{j=1}^{\infty} (\mathbb{Z} \times \mathbb{Z}) = (\mathbb{Z} \times \mathbb{Z})^{\omega} \cong \mathbb{Z}^{\omega}, \\
 H_n(\mathcal{S}_1) &\cong \pi_n(\mathcal{S}_1, s) \cong H_n(\mathcal{S}; \mathbb{Z}) \times \prod_{j=2}^{\infty} H_n(\mathcal{S}_1(j); \mathbb{Z}) \cong \mathbb{Z} \times (\mathbb{Z} \times \mathbb{Z})^{\omega} \cong \mathbb{Z}^{\omega}.
 \end{aligned}
 \tag{3.17}$$

Since both \mathcal{S}_0 and \mathcal{S}_1 are $(n - 1)$ -connected, for $n > 1$, the Universal Coefficients Theorem for cohomology and (3.17) implies additively

$$H^n(\mathcal{S}_*) \cong \text{Hom}(H_n(\mathcal{S}_*; \mathbb{Z}); \mathbb{Z}) \cong \bigoplus_{j=1}^{\infty} H^n(\mathcal{S}_*(j); \mathbb{Z}) \cong \bigoplus_{k=1}^{\infty} \mathbb{Z},
 \tag{3.18}$$

(c.f. [20, p. 67] for the second isomorphism).

Theorem 3.9 \mathcal{S}_0 and \mathcal{S}_1 are h -equal but not homotopy equivalent.

Proof The h -equality has been already argued at the beginning of this section. For the second claim, first we note that the graded ring structures of each factor $H^*(\mathcal{S}_*(j); \mathbb{Z})$ are well known, i.e.

$$\begin{aligned}
 H^*(\mathcal{S}_0(j)) &\cong \mathbb{Z}[x_j, y_j] / \langle x_j^2 = 0, y_j^2 = 0 \rangle, \\
 H^*(\mathcal{S}_1(1)) &\cong \mathbb{Z}[w] / \langle w^2 = 0 \rangle, \quad H^*(\mathcal{S}_1(k)) \cong \mathbb{Z}[x_k, y_k] / \langle x_k^2 = 0, y_k^2 = 0 \rangle, \quad k > 1,
 \end{aligned}
 \tag{3.19}$$

where x_i, y_i and w are of degree n . Observe that the graded ring $H^*(\mathcal{S}_0)$ has the following property:

(*) For any nontrivial p in $H^n(\mathcal{S}_0)$ there exists q in $H^n(\mathcal{S}_0)$ such that $p \cdot q \neq 0$.

Indeed, from (3.18) any $p \in H^n(\mathcal{S}_0)$ is given as

$$p = \sum_{i=1}^{\infty} (a_i x_i + b_i y_i), \quad a_i, b_i \in \mathbb{Z},$$

where only finitely many a_i 's and b_i 's are nonzero. Let $r : \mathcal{S}_0 \longrightarrow \mathcal{S}_0(1) \vee \dots \vee \mathcal{S}_0(k)$ be a retraction on k first factors of \mathcal{S}_0 , and $r^* : H^*(\bigvee_{l=1}^k \mathcal{S}_0(l)) \longrightarrow H^*(\mathcal{S}_0)$ the




Fig. 5 Bouquets \mathcal{WS}_0 (left) and \mathcal{WS}_1 (right), for $n = 1$

induced monomorphism. Choosing k large enough, and using the same symbols for the generators of $H^n(\bigvee_{l=1}^k \mathcal{S}_0(l))$ as in (3.19), we have

$$p = r^*(p') = r^*\left(\sum_{i=1}^k (a_i x_i + b_i y_i)\right),$$

for some $p' \in H^n(\bigvee_{l=1}^k \mathcal{S}_0(l))$. Note that $H^*(\bigvee_{l=1}^k \mathcal{S}_0(l)) \cong \bigoplus_{l=1}^k H^*(\mathcal{S}_0(l))$ as graded rings. Let $q' = y_i$, for i such that $a_i \neq 0$, then $p' \cdot q' = a_i x_i y_i \neq 0$ and we obtain

$$p \cdot q = r^*(p' \cdot q') \neq 0,$$

by the injectivity of r^* . Clearly, the property (*) is preserved under the graded ring isomorphisms. Note that for $H^*(\mathcal{S}_1)$ the property (*) does not hold. Indeed, let $p = w \in H^n(\mathcal{S}_1)$, which generates the cohomology of the \mathcal{S} -factor of \mathcal{S}_1 , (3.19). By (3.18), any $q \in H^n(\mathcal{S}_1)$ is represented by

$$q = c_0 w + \sum_{i=1}^{\infty} (a_i x_i + b_i y_i), \quad c_0, a_i, b_i \in \mathbb{Z},$$

where only finitely many a_i 's and b_i 's are nonzero. Again, choosing an appropriate retraction r of \mathcal{S}_1 onto finitely many factors, we have $p = r^*(p')$, $q = r^*(q')$ for some p' and q' in $H^n(\bigvee_{l=1}^k \mathcal{S}_1(l))$, and therefore

$$p \cdot q = r^*(p' \cdot q') = r^*(w \cdot q') = 0.$$

We conclude, as graded rings

$$H^*(\mathcal{S}_0) \not\cong H^*(\mathcal{S}_1),$$

and consequently \mathcal{S}_0 and \mathcal{S}_1 cannot be homotopy equivalent. □

Modifying slightly the construction of \mathcal{S}_0 and \mathcal{S}_1 , one may consider an infinite compact bouquets \mathcal{WS}_0 and \mathcal{WS}_1 embedded in \mathbb{R}^{2n+2} as pictured on Fig. 5, having the same factors as \mathcal{S}_0 and \mathcal{S}_1 . We claim that \mathcal{WS}_0 and \mathcal{WS}_1 are homotopy equivalent to their respective counterparts \mathcal{S}_0 and \mathcal{S}_1 . For this purpose, let us consider just the case of \mathcal{WS}_0 and \mathcal{S}_0 (for \mathcal{WS}_1 and \mathcal{S}_1 the claim follows analogously).




Fig. 6 Homotopy equivalences P , Q and C on spaces S_0 , WS_0 and MS_0

For simplicity, we just consider the case S is a circle ($n = 1$), then all spaces can be embedded in \mathbb{R}^3 as shown⁵ on Fig. 6. The following construction can be conducted for any n , without major changes. Before we describe the intermediate space MS_0 of Fig. 6, let us define the map

$$Q : WS_0 \longrightarrow S_0, \tag{3.20}$$

which is the required homotopy equivalence. Denote by $a_i \subset WS_0$ arcs connecting successive wedge points in WS_0 , and $A = \bigcup_i a_i$ their union. The map Q is the quotient projection identifying A with the wedge point x_0 of S_0 .

Next, let us define an intermediate space MS_0 . It is built from a contractible broom $\mathcal{M} = \bigcup_{i=0}^\infty c_i$ of a carefully chosen sequence of segments $\{c_i\}_{i=0}^\infty$ in \mathbb{R}^3 , all having the common right endpoint, i.e. the wedge point x_0 of \mathcal{M} . At the left endpoints of c_i 's we attach S^2 -factors, as shown on Fig. 6. We assume that the sequence $\{c_i\}_{i=0}^\infty$ starts with a horizontal segment $c_0 = [-1, 0]$ contained in the xz -plane of \mathbb{R}^3 . As $i \rightarrow \infty$, we require the lengths of c_i 's, and the diameters of S^2 -factors tend to zero. As a result, c_i 's and the S^2 -factors of MS_0 are converging to the point $x_0 \in MS_0$ in the limit. The map

$$C : S_0 \longrightarrow MS_0, \tag{3.21}$$

in Fig. 6 is defined as follows; let c_i be a small contractible neighborhood of x_0 in the i th S^2 -factor of S_0 . The map C collapses each c_i to the corresponding segment in MS_0 , also called c_i . Clearly, S_0 and MS_0 are homotopy equivalent under C , the inverse homotopy equivalence simply contracts the broom \mathcal{M} to the wedge point x_0 in MS_0 .

Now, let \mathcal{M}' be an extension of the broom \mathcal{M} in MS_0 , given by

$$\mathcal{M}' = c_0 \cup (a_1 \cup c_1) \cup (a_2 \cup c_2) \cup \dots = c_0 \cup \bigcup_{i=1}^\infty (a_i \cup c_i),$$

⁵ Note that Fig. 4 shows a different embedding of S_0 in \mathbb{R}^3 .

where each c_i , for $i > 0$, is extended by adjoining an arc a_i to the left endpoint of c_i . In order to define the map

$$P : \mathcal{MS}_0 \longrightarrow \mathcal{WS}_0, \tag{3.22}$$

of Fig. 6, first, consider a retraction $p : \mathcal{M}' \longrightarrow c_0$ of \mathcal{M}' onto the c_0 segment of \mathcal{M}' . The arcs $a_i \cup c_i$ can be arranged appropriately, so that p can be induced by restriction of the projection $(x, y, z) \longrightarrow (x, 0, 0)$ in \mathbb{R}^3 to \mathcal{M}' , mapping each $a_i \cup c_i$ of the broom onto a portion of c_0 . Clearly, p defines a deformation retraction⁶ of \mathcal{M}' onto c_0 . The map P identifies the segment $c_0 \subset \mathcal{MS}_0$ with the arc $A = \bigcup_i a_i$ in \mathcal{WS}_0 . The map P is equal to p along $\mathcal{M}' \subset \mathcal{MS}_0$, and identifies the \mathcal{S}^2 -factors of \mathcal{MS}_0 with the corresponding factors in \mathcal{WS}_0 .

Proposition 3.10 *The Hawaiian earrings \mathcal{S}_0 and \mathcal{S}_1 of Fig. 4 are homotopy equivalent to infinite wedges \mathcal{WS}_0 and \mathcal{WS}_1 of Fig. 5.*

Proof (Sketch of Proof) Having the maps C and P defined, we set $\widehat{Q} = P \circ C$, and claim that it defines a homotopy inverse of Q . Indeed, the composition $Q \circ \widehat{Q} : \mathcal{S}_0 \longrightarrow \mathcal{S}_0$ maps the inverse image $C^{-1}(\mathcal{M})$ to the point x_0 , and is an embedding on $\mathcal{S}_0 - C^{-1}(\mathcal{M})$. The set $C^{-1}(\mathcal{M})$ is the union of $c_i \subset \mathcal{S}_0$ and arcs $C^{-1}(a_i)$, thus $C^{-1}(\mathcal{M})$ can be continuously contracted to the point $x_0 \in \mathcal{S}_0$, implying $Q \circ \widehat{Q} \simeq \text{id}_{\mathcal{S}_0}$. The composite $\widehat{Q} \circ Q : \mathcal{WS}_0 \longrightarrow \mathcal{WS}_0$, maps small contractible neighborhoods $U(a_i)$ of arcs a_i in \mathcal{WS}_0 together with paths⁷ $g_i = Q^{-1}(C^{-1}(a_i))$ to the arc $A = \bigcup_i a_i$. Both A and the union of subsets $U(a_i) \cup g_i$ are contractible within \mathcal{WS}_0 , implying again $\widehat{Q} \circ Q \simeq \text{id}_{\mathcal{WS}_0}$. □

Remark 3.11 Observe that, contrary to \mathcal{S}_* , the complement of the infinity point in \mathcal{WS}_* , denoted by \mathcal{WS}_*° is an $(n - 1)$ -connected, locally contractible, ANR. Note that if \mathcal{WS}_* were locally $2n$ -connected, it would imply it was an ANR. In this sense, \mathcal{WS}_0 and \mathcal{WS}_1 are “close” to being ANR spaces. In [22], Stewart shows \mathcal{WS}_0° and \mathcal{WS}_1° for $n = 1$ are examples of noncompact h -equal ANRs, which are not homotopy equivalent. The proof of [22], relies heavily on the non-triviality of the fundamental group \mathcal{S}_0 and \mathcal{S}_1 and on the structure of isomorphisms between free products of groups, [22]. We wish to point out that the computation of cohomology rings $H^*(\mathcal{WS}_0^\circ)$, $H^*(\mathcal{WS}_1^\circ)$ is basic and the argument of Theorem 3.9 can be easily adapted to show $H^*(\mathcal{WS}_0^\circ) \not\cong H^*(\mathcal{WS}_1^\circ)$, which implies $\mathcal{WS}_0^\circ \not\cong \mathcal{WS}_1^\circ$. Moreover, the argument is valid for all $n \geq 1$, giving $2n$ -dimensional examples which are $(n - 1)$ -connected for $n \geq 1$.

In examples, from the last two sections, homotopy dominations are given by retractions. Consequently, these are examples of r -equal continua, which are not homotopy equivalent (c.f. [5]).

Acknowledgements We wish thank the anonymous referee for remarks and corrections which improved quality of the paper.

⁶ Via e.g. a piecewise linear homotopy.

⁷ Q is an embedding away from the arc A .

References

1. Bauer, T., Kitchloo, N., Notbohm, D., Pedersen, E.K.: Finite loop spaces are manifolds. *Acta Math.* **192**(1), 5–31 (2004)
2. Berridge, P.H., Dunwoody, M.J.: Nonfree projective modules for torsion-free groups. *J. London Math. Soc. (2)* **19**(3), 433–436 (1979)
3. Berstein, I., Ganea, T.: Remark on spaces dominated by manifolds. *Fund. Math.* **47**, 45–56 (1959)
4. Borel, A.: The Poincaré duality in generalized manifolds. *Michigan Math. J.* **4**, 227–239 (1957)
5. Borsuk, K.: Theory of Retracts. *Monografie Matematyczne, Tom 44.* Państwowe Wydawnictwo Naukowe, Warsaw (1967)
6. Bredon, G.E.: Generalized manifolds, revisited. In: *Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, GA, 1969)*, pp. 461–469. Markham, Chicago (1970)
7. Bredon, G.E.: *Sheaf Theory, Volume 170 of Graduate Texts in Mathematics*, 2nd edn. Springer, New York (1997)
8. Bryant, J., Ferry, S., Mio, W., Weinberger, S.: Topology of homology manifolds. *Ann. Math. (2)* **143**(3), 435–467 (1996)
9. Eda, K., Kawamura, K.: Homotopy and homology groups of the n -dimensional Hawaiian earring. *Fund. Math.* **165**(1), 17–28 (2000)
10. Harlander, J., Jensen, J.A.: Exotic relation modules and homotopy types for certain 1-relator groups. *Algebra Geom. Topol.* **6**, 2163–2173 (electronic) (2006)
11. Hausmann, J.C.: Geometric Hopfian and non-Hopfian situations. In: *Geometry and topology (Athens, GA, 1985)*, volume 105 of *Lecture Notes in Pure and Appl. Math.*, pp. 157–166. Dekker, New York (1987)
12. Hilton, P.J., Wylie, S.: *Homology theory: an introduction to algebraic topology.* Cambridge University Press, New York (1960)
13. Johnston, H., Ranicki, A.: Homology manifold bordism. *Trans. Am. Math. Soc.* **352**(11), 5093–5137 (2000)
14. Karimov, U., Repovš, D., Rosicki, W., Zastrow, A.: On two-dimensional planar compacta not homotopically equivalent to any one-dimensional compactum. *Topol. Appl.* **153**(2–3), 284–293 (2005)
15. Kirby, R.C., Siebenmann, L.C.: *Foundational essays on topological manifolds, smoothings, and triangulations.* Princeton University Press, Princeton; University of Tokyo Press, Tokyo (1977). With notes by John Milnor and Michael Atiyah. *Annals of Mathematics Studies*, No. 88
16. Kołodziejczyk, D.: Homotopy dominations by polyhedra with polycyclic-by-finite fundamental groups. *Topol. Appl.* **153**(2–3), 294–302 (2005)
17. Kwasik, S.: On homotopy domination. *Can. Math. Bull.* **27**(4), 448–451 (1984)
18. Lam, T.Y.: *Lectures on Modules and Rings.* Graduate Texts in Mathematics, vol. 189. Springer, New York (1999)
19. Milnor, J.: On spaces having the homotopy type of a CW-complex. *Trans. Am. Math. Soc.* **90**, 272–280 (1959)
20. Nunke, R.J.: On direct products of infinite cyclic groups. *Proc. Am. Math. Soc.* **13**, 66–71 (1962)
21. Quinn, F.: Ends of maps. I. *Ann. Math. (2)* **110**(2), 275–331 (1979)
22. Stewart, T.E.: On R -equivalent spaces. *Nederl. Akad. Wetensch. Proc. Ser. A 61 = Indag. Math.* **20**, 460–462 (1958)
23. Wall, C.T.C.: Finiteness conditions for CW-complexes. *Ann. Math.* **2**(81), 56–69 (1965)
24. West, J.E.: Mapping Hilbert cube manifolds to ANR's: a solution of a conjecture of Borsuk. *Ann. Math. (2)* **106**(1), 1–18 (1977)