

Erratum to: Open Flow Microperfusion as a Dermal Pharmacokinetic Approach to Evaluate Topical Bioequivalence

Manfred Bodenlenz¹ · Katrin I. Tiffner¹ · Reingard Raml¹ · Thomas Augustin¹ · Christian Dragatin¹ · Thomas Birngruber¹ · Denise Schimek¹ · Gerd Schwagerle² · Thomas R. Pieber^{1,2} · Sam G. Raney³ · Isadore Kanfer^{4,5} · Frank Sinner^{1,2}

Published online: 21 November 2016
© Springer International Publishing Switzerland 2016

Erratum to: Clin Pharmacokinet DOI 10.1007/s40262-016-0442-z

In the original publication, some errors are found and the following corrections should be noted:

Abstract, ‘Methods’ section, 8th sentence which previously read:

...curve and maximum plasma concentration) where the...

Should read:

...curve and maximum dermal concentration) where the...

Table 1, the notes which previously read:

... C_{\max} maximum plasma concentration,...

Should read:

... C_{\max} maximum dermal concentration,...

Table 2, the notes which previously read:

... C_{\max} maximum plasma concentration,...

Should read:

... C_{\max} maximum dermal concentration,...

The online version of the original article can be found under doi:[10.1007/s40262-016-0442-z](https://doi.org/10.1007/s40262-016-0442-z).

✉ Frank Sinner
frank.sinner@joanneum.at

- ¹ HEALTH-Institute for Biomedicine and Health Sciences, Joanneum Research Forschungsgesellschaft mbH, Neue Stiftingtalstrasse 2, 8010 Graz, Austria
- ² Division of Endocrinology and Diabetology, Department of Internal Medicine, Medical University of Graz, Graz, Austria
- ³ Division of Therapeutic Performance, Office of Research and Standards, Office of Generic Drugs, US Food and Drug Administration, Silver Spring, MD, USA
- ⁴ Faculty of Pharmacy, Rhodes University, Grahamstown, South Africa
- ⁵ Present Address: Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON, Canada