

Recent Translational Findings on Impulsivity in Relation to Drug Abuse

Jessica Weafer · Suzanne H. Mitchell · Harriet de Wit

Published online: 18 September 2014
© Springer International Publishing AG 2014

Abstract Impulsive behavior is strongly implicated in drug abuse, as both a cause and a consequence of drug use. To understand how impulsive behaviors lead to and result from drug use, translational evidence from both human and non-human animal studies is needed. Here, we review recent (2009 or later) studies that have investigated two major components of impulsive behavior, inhibitory control and impulsive choice, across preclinical and clinical studies. We concentrate on the stop-signal task as the measure of inhibitory control and delay discounting as the measure of impulsive choice. Consistent with previous reports, recent studies show greater impulsive behavior in drug users compared with non-users. Additionally, new evidence supports the prospective role of impulsive behavior in drug abuse, and has begun to identify the neurobiological mechanisms underlying impulsive behavior. We focus on the commonalities and differences in findings between preclinical and clinical studies, and suggest future directions for translational research.

Keywords Impulsive behavior · Inhibitory control · Stop-signal task · Impulsive choice · Delay discounting · Drug abuse

J. Weafer · H. de Wit (✉)
Department of Psychiatry and Behavioral Neuroscience, University of Chicago, 5841 S. Maryland Avenue MC3077, Chicago, IL 60637, USA
e-mail: hdew@uchicago.edu

J. Weafer
e-mail: jweafer@uchicago.edu

S. H. Mitchell
Departments of Behavioral Neuroscience and Psychiatry, L470, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239, USA
e-mail: mitchesu@ohsu.edu

Introduction

Impulsive behavior is strongly implicated in drug abuse, as both a cause and a consequence of drug use. The role of impulsive behavior in drug abuse has been investigated extensively, as described in several recent reviews [1–4]. These reviews confirm that impulsive behavior is multifaceted, and that specific components of impulsivity relate to distinct stages of drug abuse. There is now convincing evidence that impulsive behavior is both a determinant and consequence of drug abuse, but there is still a need to identify which factors predispose individuals to use, and which result from drug use, either after acute administration or after extended periods of use. There is also a need to investigate the underlying neurobiological mechanisms involved in impulsivity to develop effective prevention and treatment strategies.

To understand how impulsive behaviors lead to, and result from, drug use, translational evidence from both human and non-human animal studies is needed, and translational research requires valid and sensitive behavioral models of impulsivity. Here, we critically examine recent studies that have investigated two major components of impulsive behavior, inhibitory control and impulsive choice, across preclinical (i.e., studies involving nonhumans) and clinical studies (i.e., studies involving humans). To reduce overlap with previous reviews, we focus on studies published in 2009 or later, and concentrate our review on the stop-signal task as the measure of inhibitory control and delay discounting as the measure of impulsive choice. We will first review the task methodologies. We will then review recent studies comparing task performance in drug users and non-users, as well as recent preclinical and clinical studies that prospectively predict drug use from these impulsive behaviors. We will also present results of studies investigating the acute effects of drugs on these tasks, because drug consumption can produce state-level disruption of impulsive behavior, and may provide some indirect

information regarding chronic drug effects. Finally, we review findings on the neurobiological mechanisms of impulsive behavior in both non-human animals and humans. Our focus is on the commonalities and differences in findings between preclinical and clinical studies, and an assessment of the strengths and limitations of current approaches.

Task Descriptions

Stop-Signal Task The stop-signal task is one of the most commonly used measures of inhibitory control. This task measures the ability to inhibit an instigated or ‘prepotent’ response [5]. In human versions of this task, subjects are instructed to respond as quickly as possible to Go signals by making a key press, but occasionally to inhibit their response when a Stop signal occurs (typically an auditory tone). The Stop signal is presented shortly after the Go signal and the experimenter measures the time an individual needs between the Go and Stop signal to successfully inhibit a response. In one commonly used version of the task, the onset of the Stop signal is adjusted to target a 50 % successful inhibition rate. A stop signal reaction time (SSRT) may be calculated (i.e., the difference between mean Go reaction time and mean stop signal delay), providing a quantitative index of inhibitory control. Greater SSRT values indicate poorer response inhibition. In animal studies, the stop-signal task can be arranged in a similar manner by adjusting the onset of the stop signal based on performance [6]. However, more commonly, the stop signal onset is varied in a pre-planned sequence across blocks and the percent of trials on which stopping occurs is measured. SSRT is obtained by interpolating the time at which 50 % stopping would occur [7].

Delay Discounting Task Delay discounting tasks are the gold standard to measure impulsive choice. These tasks measure the degree to which the value of a reward decreases as a function of the delay in time to its delivery [8–11]. In human versions of the task, subjects make choices between small rewards (usually money) delivered immediately or larger rewards delivered after a delay. A curve is plotted based on the subject’s points of indifference between immediate and delayed rewards, and steeper discounting curves indicate greater delay discounting. These tasks can be administered as both paper and pencil or computerized versions. In preclinical studies, several variants are commonly used. Similar to tasks used with human subjects, some identify indifference points by titrating the length of the delay to the large reward [12] or the size of the immediate reward [13], until an indifference point is reached. Other procedures assess choices of the larger, later reward over a series of fixed delays

presented during the course of a session [14] and use percent choice at the various delays as their measure of discounting.

Studies Comparing Drug Users and Non-Users

Studies comparing drug users and non-users provide important information about behavioral tendencies that make it difficult for users to resist drugs during attempts to abstain. However, the origin of these differences is difficult to discern because users and nonusers differ on many other dimensions (e.g., trait differences or differences in experience), and because it is not possible to determine whether differences pre-date, or result from, the drug use. Nevertheless, evidence that drug users are more impulsive on certain measures may guide development of treatment strategies, and guide preclinical prospective studies examining behavioral risk factors.

Stop-signal Task Drug users have poorer inhibitory control as measured by the stop-signal task compared with non-users [1, 4]. Recent findings generally support this conclusion, although there are some interesting exceptions, and the findings vary across drug types. Alcohol-dependent individuals (inpatient and outpatient) exhibited poorer performance on the stop-signal task than healthy controls [15, 16], but among nondependent drinkers the findings are less clear. Smith and Mattick [17] found that longer SSRTs predicted binge drinking and alcohol problems in women, but two other studies observed no differences between binge and non-binge drinkers [18, 19], and a third study reported that stop-signal task performance did not predict consumption in non-treatment-seeking problem drinkers [20]. Additional studies have reported longer SSRTs in cannabis users [18] and cocaine-dependent individuals [21], although two other studies failed to observe these group differences [22, 23]. The inconsistent results across studies remain to be explained, but these studies provide some evidence that drug users have poor inhibitory control. The association appears to be most pronounced for dependent alcohol users, and less consistent for cannabis and cocaine users.

Delay Discounting Substance users also exhibit greater delay discounting than non-users [1, 4, 24, 25], and this conclusion is supported by findings in the last 5 years. In most studies examining delay discounting in relation to habitual alcohol consumption, heavier drinkers discounted more [20, 15, 26, 18, 27]; although see [28]. One study found an association in African Americans but not European Americans [29], and another study found that discounting predicted drinking in females but not males [30]. In studies of cigarette smokers, discounting was also related to severity of nicotine

dependence [31–35], but see [36], and individuals who were both drinkers and smokers discounted more than drinkers or smokers only [37]. Similarly, both cocaine- and opiate-dependent individuals discounted more than healthy controls [21, 38–40], although see [41], and delay discounting was greater for early or intermediate onset opiate dependence compared with late-onset [42]. In regard to cannabis, discounting predicted use in recreational users [18], but not dependent individuals [33]. Finally, among users of multiple substances, discounting was greater in users of cocaine, nicotine, or both, compared with controls [43], although no difference was observed in polysubstance-using rave attenders compared with controls [44]. Taken together, these studies add to a large body of evidence that steeper discounting is greater in substance users, including alcohol, nicotine, cannabis, cocaine, and opiate users, compared with non-users.

Summary In sum, studies reviewed in this section provide further evidence that drug users exhibit both poor inhibitory control and steep delay discounting. The associations are observed across drug classes, including alcohol, stimulants, and opiates, and they tend to be more pronounced in individuals meeting criteria for substance dependence. The following sections provide updated preclinical and clinical evidence regarding the causality of this relationship and whether drug users are likely to be more impulsive even before drug use, or whether the behaviors change as a consequence of drug use, or whether other factors (e.g., smoking) contribute to the relationship between impulsivity and drug use.

Impulsive Behavior as a Determinant of Drug Use

Inhibitory Control: Preclinical Studies Our literature search produced no relevant preclinical studies examining stop-signal task performance as a determinant of drug use. However, a number of studies have been conducted using the 5-choice serial reaction time task (5-CSRTT [45]), which examines responses occurring prematurely to a Go signal. Several studies have shown that ‘high’ impulsivity, based on the number of premature responses, is a predictor of drug self-administration in laboratory animals (e.g., [46–48]).

Inhibitory Control: Clinical Studies We found only one recent prospective study that examined stop-signal task performance as a predictor of drug use in humans. This study, done by Fernie et al. [49•], assessed inhibitory control and delay discounting as predictors of alcohol consumption every 6 months over a period of 2 years, in adolescents aged 12–13 years. Prospective analyses showed that stop-signal task performance consistently predicted alcohol consumption

6 months later across the 2-year period, consistent with the idea that poor inhibition may precede and perhaps play a causal role in increased alcohol use. This is consistent with two earlier reports showing that stop-signal task performance predicted future drug and alcohol use in adolescents and young adults [50, 51]. However, as in any prospective study of this type, it is difficult to rule out other individual difference factors that might covary with stop-signal task performance and mediate the observed relationship [4].

Delay Discounting: Preclinical Studies Consistent with earlier studies [1, 4], recent experiments suggest a relationship between steeper discounting and drug self-administration in laboratory animals. Anker et al. [52] reported that rats selected for high levels of choice impulsivity acquired cocaine self-administration more quickly than animals that discounted less. Compatible with these results, Koffarnus and Woods [53] demonstrated that rats with steeper discounting functions were less sensitive to increasing response costs to earn cocaine infusions (low elasticity of demand), but they were not less sensitive to increasing cost of nondrug, sucrose reinforcers. Diergaarde et al. [54•] observed the same relationship using a different stimulant, nicotine, but not for ethanol as the reinforcer. In a somewhat different paradigm, Yates et al. [55] reported that rats selected for high levels of impulsive choice developed a conditioned place preference for amphetamine (several doses), whereas low impulsive animals did not develop a conditioned place preference for amphetamine at any dose; suggesting that steeper discounting was predictive of amphetamine reward, as measured by the place preference procedure. In sum, recent preclinical studies suggest that steeper discounting is a predictor of indices of drug use, but this association is only robust for stimulants.

Delay Discounting: Clinical Studies Fernie et al. [49•], mentioned earlier, found that steeper discounting prospectively predicted greater alcohol consumption among adolescents over the following 2-year period. This result differs from the Diergaarde et al. [54•] finding that discounting in rats did not predict alcohol consumption, but many differences between the clinical and preclinical studies could account for this dissimilar outcome. In another study, Brody et al. [56] examined delay discounting as a predictor of drug use 1 year later in 19–20-year-old African Americans. Although delay discounting did not predict drug use at follow-up in the sample as a whole, steeper discounting did predict drug use among men with high catecholamine levels (i.e., epinephrine and norepinephrine). To our knowledge, these are the first studies to prospectively assess delay discounting as a predictor of drug use in humans, and together they provide some preliminary evidence suggesting delay discounting may have a prospective role in drug abuse.

Summary of Preclinical and Clinical Studies In sum, recent preclinical and clinical studies add to a growing literature supporting the prospective role of delay discounting in drug abuse. The stop-signal task has not been thoroughly tested as a predictor of drug reward in laboratory animals, although there is some evidence that another measure of inhibitory control, the 5-CSRTT, does predict drug self-administration. A relatively small number of prospective studies with humans suggest that both poor inhibitory control and delay discounting predict drug use in children and adolescents. However, there are substantial methodological and conceptual challenges in identifying common predictors of drug reward in humans and laboratory animals, and for now we can only speculate about the causal determinants in humans.

Impulsivity as a Consequence of Drug Use

It has been proposed that drug users become more impulsive as a result of chronic exposure to drugs [57]. Although the effects of chronic drug use can readily be studied in animal models, it is extremely difficult to demonstrate that cognitive dysfunction is a result of chronic drug use in humans. However, some indirect evidence may be gleaned from studies with acute drug administration in humans. Single doses of drugs have marked effects on impulsive behaviors, which provide important information about safety, and may contribute to risk for continued use within a drug-using occasion. The acute effects may also shed light on the effects of chronic exposure to the drugs.

Inhibitory Control (Acute Drug Effects): Preclinical Studies In their previous review, Perry and Carroll [4] concluded that stimulant drugs improve stop-signal task performance in laboratory animals, especially in animals with poor baseline inhibition, and that alcohol and other sedatives impair performance. Only two recent preclinical studies have examined the acute effects of drugs on the stop-signal task, both with rats (see Table 1). Pattij et al. [58] reported that neither morphine nor naloxone altered stop-signal task performance but, consistent with earlier reports, Eagle et al. [59] reported that d-amphetamine dose-relatedly decreased SSRT. Thus, similar to earlier reports, these preclinical studies provide further evidence that stimulants improve inhibitory control, and provide new evidence suggesting that opioid drugs do not affect inhibition.

Inhibitory Control (Acute Drug Effects): Clinical Studies Consistent with preclinical studies, previous reviews concluded that stimulant drugs tend to improve stop-signal task performance, especially in individuals with poor baseline inhibition, whereas sedative drugs, including alcohol and

THC, impair stop-signal task performance [1, 4]. Recent reports have examined the effects of alcohol, marijuana smoking, and oxycodone on inhibitory control in humans (Table 1). Caswell et al. [60] found that alcohol dose-dependently increased SSRT in young adult social drinkers. Metrik et al. [61] found that smoked marijuana significantly impaired performance on the stop-signal task in young adult marijuana users. By contrast, oxycodone did not alter stop-signal task performance among healthy young adults [62]. In sum, consistent with prior reports of acute drug effects on stop-signal task performance [4], alcohol and THC appear to impair inhibitory control while oxycodone does not.

Delay Discounting (Acute Drug Effects): Preclinical Studies Although results have been mixed, prior preclinical studies indicate that acute doses of stimulant drugs decrease delay discounting whereas alcohol increases it [4]. Numerous studies have since examined the effects of drugs of abuse on delay discounting performance (see Table 2 for details). The results are complex and it appears that the effects of drugs depend on genetic factors; for example, Wooters and Bardo [63] observed that methylphenidate decreased delay discounting in Wistar-Kyoto rats but not Sprague-Dawley or Spontaneously Hypertensive rats. Further, for some drugs, the behavioral effects are closely linked to dose; for example, Huskinson and Anderson [64] reported that in Fischer 344 rats low doses of diazepam reduced delay discounting but high doses increased discounting, while in Lewis rats diazepam did not affect discounting at any dose. Pre-drug levels of discounting may also be important, as was seen in the “Impulsive behavior as a determinant of drug use” section, but this variable is as yet unexamined.

Delay Discounting (Acute Drug Effects): Clinical Studies As reported in earlier studies, there are relatively few reports that acute administration of drugs change delay discounting in humans [1, 4]. This pattern is continued in recent studies with alcohol, marijuana, d-amphetamine and oxycodone (Table 2). In one exception, Reed et al. [65] found that alcohol increased delay discounting in both heavy and light drinking women tested in the follicular phase of the menstrual cycle. Bidwell et al. [66] found no effect of alcohol on delay discounting in social drinkers, although this study did not include a placebo condition. Other studies found no effects of smoked marijuana [61], d-amphetamine [67], or oxycodone [62] on delay discounting in healthy subjects. It has been proposed that the delay discounting task typically used with humans is not sensitive to the effects of drugs [68], perhaps because of differences in time frame of the delay (i.e., seconds in rats and weeks or months in humans) or because of differences in the rewards (i.e., food vs money).

Summary of Preclinical and Clinical Studies There are discordant findings on the acute effects of drugs on discounting

Table 1 Acute drug effects on stop-signal task performance

Subjects (M:F)	Drug (dose)	Dose	Effect on SSRT	Reference
Preclinical studies				
Lister-hooded rats (n=24)	D-amphetamine	0, 0.3, 1 mg/kg	↓	[59]
Wistar rats (n=16/group)	Morphine	0, 0.3, 1, 3 mg/kg	–	[58]
	Naloxone	0, 0.3, 1, 3 mg/kg	–	
Human studies				
Healthy adults (24:24)	Alcohol	0, 0.4, 0.8 g/kg	↑ (0.8 g/kg)	[60]
Healthy marijuana smokers (88:48)	Marijuana cigarettes	0, 2.8 % THC	↑	[61]
Healthy adults (n=7)	Oxycodone	0, 5, 10, 20 mg	–	[62]

Note. All preclinical studies used males only. *SSRT* Stop signal reaction time, ↑ SSRT increased, ↓ SSRT decreased, – no drug effect.

in humans and laboratory animals. Few drugs affect discounting in humans, whereas in laboratory animals drugs have more consistent effects, although the findings depend on strain and dose. As discussed previously, it is likely that methodological differences in human and nonhuman discounting tasks (e.g., time of reward delivery and type of reward) influence the differences in findings across species.

Neurobiological Mechanisms of Impulsive Behavior and Drug Abuse

Recent advances in neurobiological research techniques have improved our understanding of neural mechanisms of impulsive behavior in relation to drug abuse. Preclinical studies have focused on identifying neuroanatomical correlates of task performance, and could provide important information regarding neurobiological mechanisms related to impulsive behavior that might serve as a predictor of drug use. By contrast, clinical studies have examined differences between drug users and non-users in brain activation during task performance, and thus provide initial information regarding correlations between drug use and brain function.

Inhibitory Control: Preclinical Studies Several strategies have been used to examine the neurobiological regions implicated in stop-signal task performance in laboratory animals: inactivating or lesioning areas of interest, or recording activity in areas of interest. The neurochemical basis of inhibition has also been studied in laboratory animals, but because there is no comparable research method in humans, we do not review these studies. To validate a mouse model of the stop-signal task, one recent study performed excitotoxic lesions of mouse medial prefrontal cortex and reported impaired stopping, as has been reported in earlier research with rats [69]. Two others recorded activation patterns using multiunit cellular recordings to provide localization information. Chen et al. [70] reported activation using intracranial local field potentials in

the supplementary motor area (SMA), and Schmidt et al. [71] used multiunit recordings from rats to describe a circuit in which neurons of the subthalamic nucleus respond to stop cues, regardless of whether stopping subsequently occurs. Only activity of cells downstream of the subthalamic nucleus, in the substantia nigra pars reticulata, was reliably coupled to successful stopping.

Inhibitory Control: Clinical Studies In humans, imaging techniques have investigated brain activation during stop-signal task performance in substance users and non-users. These studies take into account differences in behavioral task performance, because poorer performance on the tasks may explain differences in brain activation. Fortunately, from the point of view of interpreting the results, only two of the studies reviewed below found group differences in task performance [72, 73]. In general, substance abusers appear to exhibit less prefrontal activation during response inhibition than control subjects, even in the absence of performance differences.

Brain activation during stop-signal task performance is typically studied using fMRI. Li et al. [74] showed that alcohol-dependent individuals had less activation in the dorsolateral prefrontal cortex (DLPFC) during response inhibition on the stop-signal task compared with controls. Similarly, among nontreatment-seeking problem drinkers, alcohol-use severity was associated with less functional connectivity in fronto-striatal networks during the stop-signal task [75]. Among smokers, heaviness of smoking was negatively correlated with PFC activation during response inhibition in adolescents [76], and heavy-smoking adults displayed decreased dorsomedial PFC activation relative to controls [77]. Stimulant-dependent individuals displayed decreased activation in the ventrolateral PFC (VLPFC) during successful inhibitions relative to controls [73], as well as reduced white matter integrity in this region, which was inversely related to SSRT [72]. Finally, Elton et al. [78] observed reduced activation during response inhibition in cocaine-dependent individuals relative to controls in a frontal-parietal network (including DLPFC, VLPFC and insula) and in a dorsomedial

Table 2 Acute drug effects on delay discounting task performance

Subjects (M:F)	Drug	Dose	Effects on delay discounting	Reference
Preclinical studies				
Wistar rats (n=30)	Amphetamine	0, 0.5 mg/kg	↓	[98•]
Lewis rats (n=8)	Amphetamine	0, 0.1, 0.3, 1.0, 1.7 mg/kg	↓ (dose-related)	[96]
Fischer 344 rats (n=8)	Amphetamine	0, 0.1, 0.3, 1.0, 1.7 mg/kg	↑ (1, 1.7 mg/kg)	[96]
Sprague-Dawley rats (n=24)	Amphetamine	0, 0.032, 0.1, 0.32, 1.0 mg/kg	No systematic effect	[99]
Sprague-Dawley rats (n=8)	Amphetamine	0, 0.1, 0.3, 1.0, 1.8 mg/kg	↓	[100]
Sprague-Dawley rats (n=8)	Amphetamine	0, 0.32, 1.0, 1.78 mg/kg	↑ (1, 1.78 mg/kg; only descending delay order)	[101]
High alcohol preferring 1 mice (15:15)	Amphetamine	0, 0.4, 0.8, 1.2 mg/kg	↓ (dose-related; dependent on length of delay)	[102]
Sprague-Dawley rats (n=16)	Amphetamine	0, 0.1, 0.3, 1.0, 1.7, 3.0 mg/kg	↑ (1, 1.7 mg/kg)	[103]
Sprague-Dawley rats (n=12)	Amphetamine	0, 0.1, 0.32, 0.56, 1.0, 1.78 mg/kg	Dose-related; ↑ (for increasing delays) ↓ (for decreasing delays)	[104]
Spontaneously hypertensive rats (n=8)	Amphetamine	0, 0.1, 0.3, 0.56, 1.0 mg/kg	–	[63]
Wistar Kyoto rats (n=8)	Amphetamine	0, 0.1, 0.3, 0.56, 1.0 mg/kg	–	[63]
Sprague-Dawley rats (n=8)	Amphetamine	0, 0.1, 0.3, 0.56, 1.0 mg/kg	–	[63]
Spontaneously hypertensive rats (n=8)	Methylphenidate	0, 1, 3, 5.6, 10 mg/kg	↓ (0.3 mg/kg)	[105]
Sprague-Dawley rats (n=5)	Methylphenidate	0, 1, 3, 10 mg/kg (d-, l-, dl-isomers)	↓ (dose-related; especially for d- and dl-isomers)	[106]
Sprague-Dawley rats (n=12)	Methylphenidate	0, 1, 3.2, 10, 17.8 mg/kg	Dose-related; ↑ (for increasing delays) ↓ (for decreasing delays)	[104]
Spontaneously hypertensive rats (n=8)	Methylphenidate	0, 1, 3, 5.6, 10 mg/kg	–	[63]
Wistar Kyoto rats (n=8)	Methylphenidate	0, 1, 3, 5.6, 10 mg/kg	↓ (5.6, 10 mg/kg)	[63]
Sprague-Dawley rats (n=8)	Methylphenidate	0, 1, 3, 5.6, 10 mg/kg	–	[63]
Lewis rats (n=8)	Nicotine	0, 0.1, 0.3, 1.0 mg/kg	↓ (0.3 mg/kg)	[107]
Fischer 344 rats (n=8)	Nicotine	0, 0.1, 0.3, 1.0 mg/kg	↓ (1.0 mg/kg)	[107]
Long Evans rats (n=5)	Nicotine	0, 0.8 mg/kg	–	[108]
Long Evans rats (n=5)	Nicotine	0, 1.2 mg/kg	↑	[108]
Fischer 344 rats (n=8)	Diazepam	0, 0.3, 0.56, 1.0, 3.0, 10 mg/kg	↑ (10 mg/kg)	[64]
Rhesus macaque monkeys (n=2; female)	Diazepam	0, 0.1, 0.32, 1.0 mg/kg	–	[109]
Sardinian alcohol-preferring rats (n=8)	Ethanol	0, 0.25, 0.5 g/kg	–	[110]
Sardinian non-alcohol-preferring rats (n=8)	Ethanol	0, 0.25, 0.5 g/kg	–	[110]
Alko alcohol-preferring rats (n=8)	Ethanol	0, 0.25, 0.5 g/kg	–	[110]
Alko non-alcohol-preferring rats (n=8)	Ethanol	0, 0.25, 0.5 g/kg	–	[110]
Rhesus macaque monkeys (n=2; female)	Ketamine	0, 0.1, 0.32 mg/kg	↑ (similarly for both doses)	[109]
Sprague-Dawley rats (n=5)	Pentobarbital	0, 1, 3, 10 mg/kg	–	[104]
Sprague-Dawley rats (n=10)	THC	0, 0.32, 1.0, 3.2, 5.6 mg/kg	↑ (dose-related)	[104]
Rhesus-macaque monkeys (n=2, female)	Morphine	0, 0.1, 0.32 mg/kg	↑ (0.32 mg/kg dose)	[109]

Table 2 (continued)

Subjects (M:F)	Drug	Dose	Effects on delay discounting	Reference
Wistar rats (n=16)	Morphine	0, 0.3, 1, 3, 6 mg/kg	↑ (dose-related; especially at 6 mg/kg; negated by 1.0 mg/kg naloxone)	[58]
Sprague-Dawley rats (n=5)	Morphine	0, 0.3, 1, 3, 10 mg/kg	No systematic effect	[106]
Sprague-Dawley rats (n=10)	Morphine	0, 1, 3.2, 5.6, 10 mg/kg	↑ (dose-related)	[104]
High Alcohol Preferring 2 mice (20:20)	Naltrexone	0, 3, 10 mg/kg	–	[102]
Human studies				
Healthy adults (5:1)	Amphetamine (intranasal and oral)	0, 16, 24, 32 mg	–	[67]
Healthy women (n=46)	Alcohol	0, 0.5, 0.75 g/kg	↑	[65]
Healthy adults (13:14)	Alcohol	40 mg/dl, 80 mg/dl	–	[66]
Healthy marijuana smokers (88:48)	Marijuana cigarettes	0, 2.8 %	–	[61]
Healthy adults (6:6)	Oxycodone	0, 5, 10, 20 mg	–	[62]

Note. Preclinical studies used males only unless otherwise noted. ↑ delay discounting increased, ↓ delay discounting decreased, – no drug effect.

PFC network, as well as negative correlations between activation of these networks and SSRT. Thus, the findings are consistent that drug users, including problem drinkers, smokers, and stimulant users, exhibit less activation in dorsomedial, dorsolateral, and ventrolateral regions of the PFC during response inhibition on the stop-signal task.

Delay Discounting: Preclinical Studies Several studies have examined performance on a delay discounting task in laboratory animals after lesions or inactivation of specific brain areas. These studies of the neuroanatomical correlates of delay discounting have tended to focus on the orbitofrontal cortex (OFC) because of its role in executive function and possible correspondence to ventromedial PFC (vmPFC) in humans. However, several studies have not found any effect of neurotoxic lesions to or inactivation of the OFC on impulsive choice [79, 80]. This failure may be due to the different roles played by the medial and lateral portions of this region. Mar et al. [81] reported that neurotoxic lesions to OFC cortex increased discounting if lesions were confined to the medial area, but lateral area lesions were associated with the opposite effect (but see also [82] using inactivation techniques). A second region of interest is the nucleus accumbens due to its role in reinforcement and reward processes. However, here too the data are inconsistent between lesion and inactivation studies. Lesions of the nucleus accumbens core increase delay discounting [83, 84]; but also see [82] who temporarily inactivated this region. Finally, a single lesion study of the subthalamic nucleus did not identify a role for this structure [85], while Abela and Chudasama [79] did report that rats with ventral hippocampal lesions showed increased impulsive choice.

Delay Discounting: Clinical Studies In humans, several recent studies have examined brain activation during delay discounting tasks in drug users compared with nonusers. These studies typically compare patterns of brain activation when subjects opt for the delayed reward versus decisions for the immediate reward. Consistent with the behavioral studies reviewed above, all but one [86] of the fMRI studies reviewed here found that users discounted more steeply than non-users. As noted above, these group differences on task performance must be taken into account when interpreting the findings. Among problem drinkers, individuals with more severe alcohol use disorders discounted more steeply and showed greater activation of several brain regions, including the anterior insula, inferior frontal gyrus, SMA/rostral anterior cingulate cortex (ACC), DLPFC, and precuneus when selecting delayed options relative to immediate options, suggesting functional anomalies and insufficient neural processing underlying impulsive choice in problem drinkers [87, 88]. In smokers, greater activation was observed in the medial PFC and anterior insular cortex when choosing the delayed option relative

to the immediate option for money and cigarettes [89]. Meade et al. [86] found that among cocaine-using HIV patients, the increase in activity in fronto-parietal regions (including precentral gyrus, ACC, and right DLPFC, VLPFC, and OFC) was less than that in non-users when making hard decisions compared with easy decisions (although they did not differ in discounting rates). Finally, Camchong et al. [90] found that cocaine-dependent individuals discounted more steeply and had higher resting functional connectivity within a network comprising the perigenual ACC and frontal and temporal regions (i.e., DLPFC, middle temporal gyrus, and superior frontal gyrus) compared with controls. In sum, these studies confirm earlier reports of greater discounting in substance abusers, and suggest that steeper discounting is associated with increased prefrontal activation when making decisions for delayed rewards.

Summary of Preclinical and Clinical Studies Taken together, some commonalities in neurobiological findings exist across preclinical and clinical studies. In both preclinical and clinical studies, stop-signal task performance is associated with activity in the prefrontal cortex and SMA. Studies of delay discounting have been less consistent, but there appears to be a role for medial regions of the prefrontal cortex in both humans and laboratory animals. It will be important for future preclinical studies to compare neurobiological correlates of task performance in drug-naïve and non-drug-naïve animals to more directly compare findings across species.

Conclusions and Future Directions

In sum, findings over the last 5 years provide further support for the role of impulsive behavior in drug abuse, both as a determinant and a consequence of use. However, the degree to which impulsivity functions as a predisposing factor to drug abuse and the degree to which it is a result of drug abuse is still not well understood. Regarding the latter, preclinical evidence suggests that delay discounting in drug-naïve animals predicts later drug administration, but longitudinal studies testing this relation in humans are limited. Similarly, there is a growing body of evidence from preclinical studies showing that poor inhibitory control, as measured by the 5-CSRTT, is associated with multiple indices of drug-taking. However, analogs of this task are just now being developed for use in human studies [19, 91, 92], which will enable researchers to assess parallels between preclinical and clinical studies using this task. Regarding impulsive behavior as a result of drug use, there is some evidence that impulsive choice increases after chronic drug administration in animal models [93–97], but as mentioned earlier this is very difficult to study in humans. Evidence showing acute effects of drugs on impulsive behavior

likely provides some indirect evidence regarding chronic effects, but it is not known whether acute effects on either impulsive action or impulsive choice bear any relation to changes in behavior after chronic administration of the drug (as in drug abusers). It seems intuitively logical that if acute behavioral impairments are continued over a long period of time they may lead to chronic changes in behavior, but again, longitudinal studies are needed to directly test this hypothesis. Finally, we are just beginning to understand the neurobiological mechanisms underlying impulsivity and drug abuse. Future translational assessments of the neural mechanisms underlying impulsive behavior and drug abuse will likely provide exciting new information and further our understanding of impulsive behavior as both a cause and consequence of drug abuse.

Acknowledgments This research was supported by National Institute on Drug Abuse Grants DA002812 and DA032015 (HdW) and F32 DA033756 (JW), and the Portland Alcohol Research Center P60 AA10760 (SHM). NIDA and NIAAA had no role in the study design, interpretation, writing the manuscript, or the decision to submit the paper for publication. The authors declare that they have no conflict of interest.

Compliance with Ethics Guidelines

Conflict of Interest Jessica Weaver, Suzanne H. Mitchell and Harriet de Wit declare no conflict of interest.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by the author.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. de Wit H. Impulsivity as a determinant and consequence of drug use: a review of underlying processes. *Addict Biol.* 2009;14(1): 22–31. doi:10.1111/j.1369-1600.2008.00129.x.
2. Grant JE, Chamberlain SR. Impulsive action and impulsive choice across substance and behavioral addictions: Cause or consequence? *Addict Behav.* doi: 10.1016/j.addbeh.2014.04.022
3. Jentsch JD, Ashenurst JR, Cervantes MC, Groman SM, James AS, Pennington ZT. Dissecting impulsivity and its relationships to drug addictions. *Ann N Y Acad Sci.* doi:10.1111/nyas.12388.
4. Perry JL, Carroll ME. The role of impulsive behavior in drug abuse. *Psychopharmacology (Berl).* 2008;200(1):1–26. doi:10.1007/s00213-008-1173-0.
5. Logan GD, Schachar RJ, Tannock R. Impulsivity and inhibitory control. *Psychol Sci.* 1997;8:60–4.
6. Feola TW, de Wit H, Richards JB. Effects of d-amphetamine and alcohol on a measure of behavioral inhibition in rats. *Behav Neurosci.* 2000;114(4):838–48.

7. Eagle DM, Robbins TW. Inhibitory control in rats performing a stop-signal reaction-time task: effects of lesions of the medial striatum and d-amphetamine. *Behav Neurosci*. 2003;117(6):1302–17. doi:10.1037/0735-7044.117.6.1302 2003-10460-017.
8. Ainslie G. Specious reward: a behavioral theory of impulsiveness and impulse control. *Psychol Bull*. 1975;82(4):463–96.
9. Rachlin H, Green L. Commitment, choice and self-control. *J Exp Anal Behav*. 1972;17(1):15–22.
10. Richards JB, Zhang L, Mitchell SH, de Wit H. Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *J Exp Anal Behav*. 1999;71(2):121–43. doi:10.1901/jeab.1999.71-121.
11. Thiebot MH, Le Bihan C, Soubrie P, Simon P. Benzodiazepines reduce the tolerance to reward delay in rats. *Psychopharmacology (Berl)*. 1985;86(1–2):147–52.
12. Perry JL, Larson EB, German JP, Madden GJ, Carroll ME. Impulsivity (delay discounting) as a predictor of acquisition of IV cocaine self-administration in female rats. *Psychopharmacology (Berl)*. 2005;178(2–3):193–201. doi:10.1007/s00213-004-1994-4.
13. Richards JB, Mitchell SH, de Wit H, Seiden LS. Determination of discount functions in rats with an adjusting-amount procedure. *J Exp Anal Behav*. 1997;67(3):353–66. doi:10.1901/jeab.1997.67-353.
14. Evenden JL, Ryan CN. The pharmacology of impulsive behaviour in rats: the effects of drugs on response choice with varying delays of reinforcement. *Psychopharmacology (Berl)*. 1996;128(2):161–70.
15. Joos L, Schmaal L, Goudriaan AE, Fransen E, Van den Brink W, Sabbe BG, et al. Age of onset and neuropsychological functioning in alcohol dependent inpatients. *Alcohol Clin Exp Res*. 2013;37(3):407–16. doi:10.1111/j.1530-0277.2012.01949.x.
16. Lawrence AJ, Luty J, Bogdan NA, Sahakian BJ, Clark L. Impulsivity and response inhibition in alcohol dependence and problem gambling. *Psychopharmacology (Berl)*. 2009;207(1):163–72. doi:10.1007/s00213-009-1645-x.
17. Smith JL, Mattick RP. Evidence of deficits in behavioural inhibition and performance monitoring in young female heavy drinkers. *Drug Alcohol Depend*. 2013;133(2):398–404. doi:10.1016/j.drugalcdep.2013.06.020.
18. Moreno M, Estevez AF, Zaldivar F, Montes JM, Gutierrez-Ferre VE, Esteban L, et al. Impulsivity differences in recreational cannabis users and binge drinkers in a university population. *Drug Alcohol Depend*. 2012;124(3):355–62. doi:10.1016/j.drugalcdep.2012.02.011.
19. Sanchez-Roige S, Baro V, Trick L, Pena-Oliver Y, Stephens DN, Duka T. Exaggerated Waiting Impulsivity Associated with Human Binge Drinking, and High Alcohol Consumption in Mice. *Neuropsychopharmacology*. doi: 10.1038/npp.2014.151.
20. Courtney KE, Arellano R, Barkley-Levenson E, Galvan A, Poldrack RA, Mackillop J, et al. The relationship between measures of impulsivity and alcohol misuse: an integrative structural equation modeling approach. *Alcohol Clin Exp Res*. 2012;36(6):923–31. doi:10.1111/j.1530-0277.2011.01635.x.
21. Crunelle CL, Veltman DJ, van Emmerik-van Oortmerssen K, Booij J, van den Brink W. Impulsivity in adult ADHD patients with and without cocaine dependence. *Drug Alcohol Depend*. 2013;129(1–2):18–24. doi:10.1016/j.drugalcdep.2012.09.006.
22. Grant JE, Chamberlain SR, Schreiber L, Odlaug BL. Neuropsychological deficits associated with cannabis use in young adults. *Drug Alcohol Depend*. 2012;121(1–2):159–62. doi:10.1016/j.drugalcdep.2011.08.015.
23. Vonmoos M, Hulka LM, Preller KH, Jenni D, Schulz C, Baumgartner MR, et al. Differences in self-reported and behavioral measures of impulsivity in recreational and dependent cocaine users. *Drug Alcohol Depend*. 2013;133(1):61–70. doi:10.1016/j.drugalcdep.2013.05.032.
24. Bickel WK, Marsch LA. Toward a behavioral economic understanding of drug dependence: delay discounting processes. *Addiction (Abingdon, England)*. 2001;96(1):73–86. doi:10.1080/09652140020016978.
25. Bickel WK, Miller ML, Yi R, Kowal BP, Lindquist DM, Pitcock JA. Behavioral and neuroeconomics of drug addiction: competing neural systems and temporal discounting processes. *Drug Alcohol Depend*. 2007;90 Suppl 1:S85–91. doi:10.1016/j.drugalcdep.2006.09.016.
26. MacKillop J, Amlung MT, Few LR, Ray LA, Sweet LH, Munafò MR. Delayed reward discounting and addictive behavior: a meta-analysis. *Psychopharmacology (Berl)*. 2011;216(3):305–21. doi:10.1007/s00213-011-2229-0. *This meta-analysis of 46 studies provides strong evidence for greater delay discounting in individuals with addictive disorders.*
27. Murphy C, Mackillop J. Living in the here and now: interrelationships between impulsivity, mindfulness, and alcohol misuse. *Psychopharmacology (Berl)*. 2012;219(2):527–36. doi:10.1007/s00213-011-2573-0.
28. Fernie G, Cole JC, Goudie AJ, Field M. Risk-taking but not response inhibition or delay discounting predict alcohol consumption in social drinkers. *Drug Alcohol Depend*. 2010;112(1–2):54–61. doi:10.1016/j.drugalcdep.2010.05.011.
29. Denhardt AA, Murphy JG. Associations between depression, distress tolerance, delay discounting, and alcohol-related problems in European American and African American college students. *Psychol Addict Behav*. 2011;25(4):595–604. doi:10.1037/a0025807.
30. Yankelevitz RL, Mitchell SH, Zhang Y. Gender differences in factors associated with alcohol drinking: delay discounting and perception of others' drinking. *Drug Alcohol Depend*. 2012;123(1–3):273–6. doi:10.1016/j.drugalcdep.2011.11.012.
31. Amlung M, MacKillop J. Clarifying the Relationship Between Impulsive Delay Discounting and Nicotine Dependence. *Psychol Addict Behav*. doi: 10.1037/a0036726.
32. Friedel JE, Dehart WB, Madden GJ, Odum AL. Impulsivity and cigarette smoking: discounting of monetary and consumable outcomes in current and non-smokers. *Psychopharmacology (Berl)*. doi:10.1007/s00213-014-3597-z.
33. Johnson MW, Bickel WK, Baker F, Moore BA, Badger GJ, Budney AJ. Delay discounting in current and former marijuana-dependent individuals. *Exp Clin Psychopharmacol*. 2010;18(1):99–107. doi:10.1037/a0018333.
34. Mitchell SH, Wilson VB. Differences in delay discounting between smokers and nonsmokers remain when both rewards are delayed. *Psychopharmacology (Berl)*. 2012;219(2):549–62. doi:10.1007/s00213-011-2521-z.
35. Peters J, Bromberg U, Schneider S, Brassens S, Menz M, Banaschewski T, et al. Lower ventral striatal activation during reward anticipation in adolescent smokers. *Am J Psychiatry*. 2011;168(5):540–9. doi:10.1176/appi.ajp.2010.10071024.
36. Poltavski DV, Weatherly JN. Delay and probability discounting of multiple commodities in smokers and never-smokers using multiple-choice tasks. *Behav Pharmacol*. 2013;24(8):659–67. doi:10.1097/FBP.0000000000000100008877-201312000-00004.
37. Moallem NR, Ray LA. Dimensions of impulsivity among heavy drinkers, smokers, and heavy drinking smokers: singular and combined effects. *Addict Behav*. 2012;37(7):871–4. doi:10.1016/j.addbeh.2012.03.002.
38. Johnson MW. An efficient operant choice procedure for assessing delay discounting in humans: initial validation in cocaine-dependent and control individuals. *Exp Clin Psychopharmacol*. 2012;20(3):191–204. doi:10.1037/a0027088.
39. Li X, Zhang F, Zhou Y, Zhang M, Wang X, Shen M. Decision-making deficits are still present in heroin abusers after short-

- long-term abstinence. *Drug Alcohol Depend.* 2013;130(1–3):61–7. doi:10.1016/j.drugalcdep.2012.10.012.
40. Robles E, Huang BE, Simpson PM, McMillan DE. Delay discounting, impulsiveness, and addiction severity in opioid-dependent patients. *J Subst Abuse Treat.* 2011;41(4):354–62. doi:10.1016/j.jsat.2011.05.003.
 41. Albein-Urios N, Martinez-Gonzalez JM, Lozano O, Clark L, Verdejo-Garcia A. Comparison of impulsivity and working memory in cocaine addiction and pathological gambling: Implications for cocaine-induced neurotoxicity. *Drug Alcohol Depend.* 2012;126(1–2):1–6. doi:10.1016/j.drugalcdep.2012.03.008.
 42. Passetti F, Verdejo-Garcia A, Abou-Saleh M. Comparatively preserved impulse control in late-onset opiate users. *Psychopharmacology (Berl).* 2013;230(3):499–505. doi:10.1007/s00213-013-3174-x.
 43. Garcia-Rodriguez O, Secades-Villa R, Weidberg S, Yoon JH. A systematic assessment of delay discounting in relation to cocaine and nicotine dependence. *Behav Process.* 2013;99:100–5. doi:10.1016/j.beproc.2013.07.007.
 44. Verdejo-Garcia A, Del Mar Sanchez-Fernandez M, Alonso-Maroto LM, Fernandez-Calderon F, Perales JC, Lozano O, et al. Impulsivity and executive functions in polysubstance-using rave attenders. *Psychopharmacology (Berl).* 2010;210(3):377–92. doi:10.1007/s00213-010-1833-8.
 45. Robbins TW. The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology (Berl).* 2002;163(3–4):362–80. doi:10.1007/s00213-002-1154-7.
 46. Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ. High impulsivity predicts the switch to compulsive cocaine-taking. *Science.* 2008;320(5881):1352–5. doi:10.1126/science.1158136.
 47. Dalley JW, Fryer TD, Brichard L, Robinson ES, Theobald DE, Laane K, et al. Nucleus accumbens D2/3 receptors predict trait impulsivity and cocaine reinforcement. *Science.* 2007;315(5816):1267–70. doi:10.1126/science.1137073.
 48. Diergaarde L, Pattij T, Poortvliet I, Hogenboom F, de Vries W, Schoffelmeer AN, et al. Impulsive choice and impulsive action predict vulnerability to distinct stages of nicotine seeking in rats. *Biol Psychiatry.* 2008;63(3):301–8. doi:10.1016/j.biopsych.2007.07.011.
 49. Fernie G, Peeters M, Gullo MJ, Christiansen P, Cole JC, Sumnall H, et al. Multiple behavioural impulsivity tasks predict prospective alcohol involvement in adolescents. *Addiction.* 2013;108(11):1916–23. doi:10.1111/add.12283. *This prospective study showed that both stop-task performance and delay discounting reliably predicted alcohol use in adolescents 6 months later.*
 50. Nigg JT, Wong MM, Martel MM, Jester JM, Puttler LI, Glass JM, et al. Poor response inhibition as a predictor of problem drinking and illicit drug use in adolescents at risk for alcoholism and other substance use disorders. *J Am Acad Child Adolesc Psychiatry.* 2006;45(4):468–75. doi:10.1097/01.chi.0000199028.76452.a9.S0890-8567(09)62067-0.
 51. Rubio G, Jimenez M, Rodriguez-Jimenez R, Martinez I, Avila C, Ferre F, et al. The role of behavioral impulsivity in the development of alcohol dependence: a 4-year follow-up study. *Alcohol Clin Exp Res.* 2008;32(9):1681–7. doi:10.1111/j.1530-0277.2008.00746.x.
 52. Anker JJ, Perry JL, Gliddon LA, Carroll ME. Impulsivity predicts the escalation of cocaine self-administration in rats. *Pharmacol Biochem Behav.* 2009;93(3):343–8. doi:10.1016/j.pbb.2009.05.013.
 53. Koffarnus MN, Woods JH. Individual differences in discount rate are associated with demand for self-administered cocaine, but not sucrose. *Addict Biol.* 2013;18(1):8–18. doi:10.1111/j.1369-1600.2011.00361.x.
 54. Diergaarde L, van Mourik Y, Pattij T, Schoffelmeer AN, De Vries TJ. Poor impulse control predicts inelastic demand for nicotine but not alcohol in rats. *Addict Biol.* 2012;17(3):576–87. doi:10.1111/j.1369-1600.2011.00376.x. *This study provides an interesting methodological link between delay discounting and behavioral procedures to assess incentive value of drug reinforcers that can be widely applied in other studies.*
 55. Yates JR, Marusich JA, Gipson CD, Beckmann JS, Bardo MT. High impulsivity in rats predicts amphetamine conditioned place preference. *Pharmacol Biochem Behav.* 2012;100(3):370–6. doi:10.1016/j.pbb.2011.07.012.
 56. Brody GH, Yu T, MacKillop J, Miller GE, Chen E, Obasi EM, et al. Catecholamine levels and delay discounting forecast drug use among African American youths. *Addiction.* 2014;109(7):1112–8. doi:10.1111/add.12516.
 57. Jentsch JD, Taylor JR. Impulsivity resulting from frontostriatal dysfunction in drug abuse: implications for the control of behavior by reward-related stimuli. *Psychopharmacology (Berl).* 1999;146(4):373–90.
 58. Pattij T, Schetters D, Janssen MC, Wiskerke J, Schoffelmeer AN. Acute effects of morphine on distinct forms of impulsive behavior in rats. *Psychopharmacology (Berl).* 2009;205(3):489–502. doi:10.1007/s00213-009-1558-8.
 59. Eagle DM, Lehmann O, Theobald DE, Pena Y, Zakaria R, Ghosh R, et al. Serotonin depletion impairs waiting but not stop-signal reaction time in rats: implications for theories of the role of 5-HT in behavioral inhibition. *Neuropsychopharmacology.* 2009;34(5):1311–21. doi:10.1038/npp.2008.202.
 60. Caswell AJ, Morgan MJ, Duka T. Acute alcohol effects on subtypes of impulsivity and the role of alcohol-outcome expectancies. *Psychopharmacology (Berl).* 2013;229(1):21–30. doi:10.1007/s00213-013-3079-8.
 61. Metrik J, Kahler CW, Reynolds B, McGeary JE, Monti PM, Haney M, et al. Balanced placebo design with marijuana: pharmacological and expectancy effects on impulsivity and risk taking. *Psychopharmacology (Berl).* 2012;223(4):489–99. doi:10.1007/s00213-012-2740-y.
 62. Zaeny JP, de Wit H. The prescription opioid, oxycodone, does not alter behavioral measures of impulsivity in healthy volunteers. *Pharmacol Biochem Behav.* 2009;94(1):108–13. doi:10.1016/j.pbb.2009.07.010.
 63. Wooters TE, Bardo MT. Methylphenidate and fluphenazine, but not amphetamine, differentially affect impulsive choice in spontaneously hypertensive, Wistar-Kyoto and Sprague-Dawley rats. *Brain Res.* 2011;1396:45–53. doi:10.1016/j.brainres.2011.04.040.
 64. Huskinson SL, Anderson KG. Effects of acute and chronic administration of diazepam on delay discounting in Lewis and Fischer 344 rats. *Behav Pharmacol.* 2012;23(4):315–30. doi:10.1097/FBP.0b013e3283564da4.00008877-201208000-00001.
 65. Reed SC, Levin FR, Evans SM. Alcohol increases impulsivity and abuse liability in heavy drinking women. *Exp Clin Psychopharmacol.* 2013;20(6):454–65. doi:10.1037/a0029087.
 66. Bidwell LC, MacKillop J, Murphy JG, Grenga A, Swift RM, McGeary JE. Biphasic effects of alcohol on delay and probability discounting. *Exp Clin Psychopharmacol.* 2013;21(3):214–21. doi:10.1037/a0032284.
 67. Lile JA, Babalonis S, Emurian C, Martin CA, Wermeling DP, Kelly TH. Comparison of the behavioral and cardiovascular effects of intranasal and oral d-amphetamine in healthy human subjects. *J Clin Pharmacol.* 2011;51(6):888–98. doi:10.1177/0091270010375956.
 68. de Wit H, Mitchell SH. Drug effects on delay discounting. In: Madden GJ, Bickel WK, editors. *Impulsivity: the behavioral and neurological science of discounting.* Washington, DC: American Psychological Association; 2010. p. 213–36.
 69. Humby T, Eddy JB, Good MA, Reichelt AC, Wilkinson LS. A novel translational assay of response inhibition and impulsivity:

- effects of prefrontal cortex lesions, drugs used in ADHD, and serotonin 2C receptor antagonism. *Neuropsychopharmacology*. 2013;38(11):2150–9. doi:10.1038/npp.2013.112.
70. Chen X, Scangos KW, Stuphorn V. Supplementary motor area exerts proactive and reactive control of arm movements. *J Neurosci*. 2010;30(44):14657–75. doi:10.1523/JNEUROSCI.2669-10.2010.
 71. Schmidt R, Leventhal DK, Mallet N, Chen F, Berke JD. Canceling actions involves a race between basal ganglia pathways. *Nat Neurosci*. 2013;16(8):1118–24. doi:10.1038/nn.3456.
 72. Ersche KD, Jones PS, Williams GB, Turton AJ, Robbins TW, Bullmore ET. Abnormal brain structure implicated in stimulant drug addiction. *Science*. 2012;335(6068):601–4. doi:10.1126/science.1214463. *This study identified abnormalities in fronto-striatal brain structure in stimulant-dependent individuals and their siblings that were associated with impaired stop task performance.*
 73. Morein-Zamir S, Simon Jones P, Bullmore ET, Robbins TW, Ersche KD. Prefrontal hypoactivity associated with impaired inhibition in stimulant-dependent individuals but evidence for hyperactivation in their unaffected siblings. *Neuropsychopharmacology*. 2013;38(10):1945–53. doi:10.1038/npp.2013.90.
 74. Li CS, Luo X, Yan P, Bergquist K, Sinha R. Altered impulse control in alcohol dependence: neural measures of stop signal performance. *Alcohol Clin Exp Res*. 2009;33(4):740–50. doi:10.1111/j.1530-0277.2008.00891.x.
 75. Courtney KE, Ghahremani DG, Ray LA. Fronto-striatal functional connectivity during response inhibition in alcohol dependence. *Addict Biol*. 2012;18(3):593–604. doi:10.1111/adb.12013.
 76. Galvan A, Poldrack RA, Baker CM, McGlennen KM, London ED. Neural correlates of response inhibition and cigarette smoking in late adolescence. *Neuropsychopharmacology*. 2011;36(5):970–8. doi:10.1038/npp.2010.235.
 77. de Ruiter MB, Oosterlaan J, Veltman DJ, van den Brink W, Goudriaan AE. Similar hypo-responsiveness of the dorsomedial prefrontal cortex in problem gamblers and heavy smokers during an inhibitory control task. *Drug Alcohol Depend*. 2012;121(1–2):81–9. doi:10.1016/j.drugalcdep.2011.08.010.
 78. Elton A, Young J, Smitherman S, Gross RE, Mletzko T, Kilts CD. Neural network activation during a stop-signal task discriminates cocaine-dependent from non-drug-abusing men. *Addict Biol*. 2012;19(3):427–38. doi:10.1111/adb.12011.
 79. Abela AR, Chudasama Y. Dissociable contributions of the ventral hippocampus and orbitofrontal cortex to decision-making with a delayed or uncertain outcome. *Eur J Neurosci*. 2013;37(4):640–7. doi:10.1111/ejn.12071.
 80. Stopper CM, Green EB, Floresco SB. Selective involvement by the medial orbitofrontal cortex in biasing risky, but not impulsive, choice. *Cereb Cortex*. 2014;24(1):154–62. doi:10.1093/cercor/bhs297.
 81. Mar AC, Walker AL, Theobald DE, Eagle DM, Robbins TW. Dissociable effects of lesions to orbitofrontal cortex subregions on impulsive choice in the rat. *J Neurosci*. 2011;31(17):6398–404. doi:10.1523/JNEUROSCI.6620-10.2011.
 82. Moschak TM, Mitchell SH. Partial inactivation of nucleus accumbens core decreases delay discounting in rats without affecting sensitivity to delay or magnitude. *Behav Brain Res*. 2014;268:159–68. doi:10.1016/j.bbr.2014.03.044.
 83. da Costa AS, Body S, Hampson CL, Langley RW, Deakin JF, Anderson IM, et al. Effects of lesions of the nucleus accumbens core on inter-temporal choice: further observations with an adjusting-delay procedure. *Behav Brain Res*. 2009;202(2):272–7. doi:10.1016/j.bbr.2009.04.003.
 84. Galtres T, Kirkpatrick K. The role of the nucleus accumbens core in impulsive choice, timing, and reward processing. *Behav Neurosci*. 2010;124(1):26–43. doi:10.1037/a0018464.
 85. Bezzina G, Cheung TH, Body S, Deakin JF, Anderson IM, Bradshaw CM, et al. Quantitative analysis of the effect of lesions of the subthalamic nucleus on intertemporal choice: further evidence for enhancement of the incentive value of food reinforcers. *Behav Pharmacol*. 2009;20(5–6):437–46. doi:10.1097/FBP.0b013e3283305e4d.
 86. Meade CS, Lowen SB, MacLean RR, Key MD, Lukas SE. fMRI brain activation during a delay discounting task in HIV-positive adults with and without cocaine dependence. *Psychiatry Res*. 2011;192(3):167–75. doi:10.1016/j.psychres.2010.12.011.
 87. Amlung M, Sweet LH, Acker J, Brown CL, MacKillop J. Dissociable brain signatures of choice conflict and immediate reward preferences in alcohol use disorders. *Addict Biol*. 2012;19(4):743–53. doi:10.1111/adb.12017.
 88. Claus ED, Kiehl KA, Hutchison KE. Neural and behavioral mechanisms of impulsive choice in alcohol use disorder. *Alcohol Clin Exp Res*. 2011;35(7):1209–19. doi:10.1111/j.1530-0277.2011.01455.x.
 89. MacKillop J, Amlung MT, Wier LM, David SP, Ray LA, Bickel WK, et al. The neuroeconomics of nicotine dependence: a preliminary functional magnetic resonance imaging study of delay discounting of monetary and cigarette rewards in smokers. *Psychiatry Res*. 2012;202(1):20–9. doi:10.1016/j.psychres.2011.10.003.
 90. Camchong J, MacDonald 3rd AW, Nelson B, Bell C, Mueller BA, Specker S, et al. Frontal hyperconnectivity related to discounting and reversal learning in cocaine subjects. *Biol Psychiatry*. 2011;69(11):1117–23. doi:10.1016/j.biopsych.2011.01.008. *This study showed that functional connectivity abnormalities in cocaine-dependent individuals are associated with increased delay discounting.*
 91. Voon V, Irvine MA, Derbyshire K, Worbe Y, Lange I, Abbott S, et al. Measuring “waiting” impulsivity in substance addictions and binge eating disorder in a novel analogue of rodent serial reaction time task. *Biol Psychiatry*. 2014;75(2):148–55. doi:10.1016/j.biopsych.2013.05.013.
 92. Worbe Y, Savulich G, Voon V, Fernandez-Egea E, Robbins TW. Serotonin depletion induces ‘waiting impulsivity’ on the human four-choice serial reaction time task: cross-species translational significance. *Neuropsychopharmacology*. 2014;39(6):1519–26. doi:10.1038/npp.2013.351.
 93. Dandy KL, Gatch MB. The effects of chronic cocaine exposure on impulsivity in rats. *Behav Pharmacol*. 2009;20(5–6):400–5. doi:10.1097/FBP.0b013e328330ad89.
 94. Gipson CD, Bardo MT. Extended access to amphetamine self-administration increases impulsive choice in a delay discounting task in rats. *Psychopharmacology (Berl)*. 2009;207(3):391–400. doi:10.1007/s00213-009-1667-4.
 95. Harvey-Lewis C, Perdriest J, Franklin KB. The effect of morphine dependence on impulsive choice in rats. *Psychopharmacology (Berl)*. 2012;223(4):477–87. doi:10.1007/s00213-012-2738-5.
 96. Huskinson SL, Krebs CA, Anderson KG. Strain differences in delay discounting between Lewis and Fischer 344 rats at baseline and following acute and chronic administration of d-amphetamine. *Pharmacol Biochem Behav*. 2012;101(3):403–16. doi:10.1016/j.pbb.2012.02.005.
 97. Kolokotroni KZ, Rodgers RJ, Harrison AA. Trait differences in response to chronic nicotine and nicotine withdrawal in rats. *Psychopharmacology (Berl)*. 2014;231(3):567–80. doi:10.1007/s00213-013-3270-y.
 98. Broos N, Schmaal L, Wiskerke J, Kosteljik L, Lam T, Stoop N, et al. The relationship between impulsive choice and impulsive action: a cross-species translational study. *PLoS One*. 2012;7(5):e36781. doi:10.1371/journal.pone.0036781. *This study provides a translational comparison of performance on tasks assessing impulsive behavior in humans and animals. Correlations among*

- performance indices on different tasks indicate the extent to which the performance processes are non-unitary.*
99. Koffarnus MN, Newman AH, Grundt P, Rice KC, Woods JH. Effects of selective dopaminergic compounds on a delay-discounting task. *Behav Pharmacol*. 2011;22(4):300–11. doi:10.1097/FBP.0b013e3283473bcb.
 100. Krebs CA, Anderson KG. Preference reversals and effects of D-amphetamine on delay discounting in rats. *Behav Pharmacol*. 2012;23(3):228–40. doi:10.1097/FBP.0b013e32835342ed00008877-201206000-00002.
 101. Maguire DR, Henson C, France CP. Effects of amphetamine on delay discounting in rats depend upon the manner in which delay is varied. *Neuropharmacology*. doi: 10.1016/j.neuropharm.2014.04.012.
 102. Oberlin BG, Bristow RE, Heighton ME, Grahame NJ. Pharmacologic dissociation between impulsivity and alcohol drinking in high alcohol preferring mice. *Alcohol Clin Exp Res*. 2010;34(8):1363–75. doi:10.1111/j.1530-0277.2010.01220.x.
 103. Slezak JM, Anderson KG. Effects of variable training, signaled and unsignaled delays, and d-amphetamine on delay-discounting functions. *Behav Pharmacol*. 2009;20(5–6):424–36. doi:10.1097/FBP.0b013e3283305ef9.
 104. Tanno T, Maguire DR, Henson C, France CP. Effects of amphetamine and methylphenidate on delay discounting in rats: interactions with order of delay presentation. *Psychopharmacology (Berl)*. 2014;231(1):85–95. doi:10.1007/s00213-013-3209-3.
 105. Slezak JM, Anderson KG. Effects of acute and chronic methylphenidate on delay discounting. *Pharmacol Biochem Behav*. 2011;99(4):545–51. doi:10.1016/j.pbb.2011.05.027.
 106. Slezak JM, Ricaurte GA, Tallarida RJ, Katz JL. Methylphenidate and impulsivity: a comparison of effects of methylphenidate enantiomers on delay discounting in rats. *Psychopharmacology (Berl)*. 2014;231(1):191–8. doi:10.1007/s00213-013-3220-8.
 107. Anderson KG, Diller JW. Effects of acute and repeated nicotine administration on delay discounting in Lewis and Fischer 344 rats. *Behav Pharmacol*. 2010;21(8):754–64. doi:10.1097/FBP.0b013e328340a050.
 108. Kelsey JE, Niraula A. Effects of acute and sub-chronic nicotine on impulsive choice in rats in a probabilistic delay-discounting task. *Psychopharmacology (Berl)*. 2013;227(3):385–92. doi:10.1007/s00213-013-2984-1.
 109. Maguire DR, Li JX, France CP. Impulsivity and drugs of abuse: a juice-reinforced operant procedure for determining within-session delay discounting functions in rhesus monkeys. *J Pharmacol Toxicol Methods*. 2012;66(3):264–9. doi:10.1016/j.vascn.2012.08.168.
 110. Wilhelm CJ, Mitchell SH. Acute ethanol does not always affect delay discounting in rats selected to prefer or avoid ethanol. *Alcohol Alcohol*. 2012;47(5):518–24. doi:10.1093/alcalc/ags059.