



Contractibility of a persistence map preimage

Jacek Cyranka^{1,2} · Konstantin Mischaikow¹ · Charles Weibel¹

Received: 29 October 2018 / Accepted: 14 August 2020 / Published online: 28 August 2020
© The Author(s) 2020

Abstract

This work is motivated by the following question in data-driven study of dynamical systems: given a dynamical system that is observed via time series of persistence diagrams that encode topological features of snapshots of solutions, what conclusions can be drawn about solutions of the original dynamical system? We address this challenge in the context of an N dimensional system of ordinary differential equation defined in \mathbb{R}^N . To each point in \mathbb{R}^N (e.g. an initial condition) we associate a persistence diagram. The main result of this paper is that under this association the preimage of every persistence diagram is contractible. As an application we provide conditions under which multiple time series of persistence diagrams can be used to conclude the existence of a fixed point of the differential equation that generates the time series.

Keywords Topological data analysis · Persistent homology · Dynamical systems · Fixed point theorem

Mathematics Subject Classification 37C25 · 55N31 · 06B35 · 55-08 · 06F30

1 Introduction

Topological data analysis (TDA), especially in the form of persistent homology, is rapidly developing into a widely used tool for the analysis of high dimensional data associated with nonlinear structures (Edelsbrunner and Harer 2010; Zomorodian and

✉ Jacek Cyranka
jcyranka@gmail.com

Konstantin Mischaikow
mishaik@math.rutgers.edu

Charles Weibel
weibel@math.rutgers.edu

¹ Department of Mathematics, Rutgers, The State University of New Jersey, 110 Frelinghusen Rd., Piscataway, NJ 08854-8019, USA

² Institute of Informatics, University of Warsaw, Banacha 2, 02-097 Warsaw, Poland

Carlsson 2005; Oudot 2015). That topological tools can play a role in this subject should not be unexpected, given the central role of nonlinear functional analysis in the study of geometry, analysis, and differential equations, for example. What is perhaps surprising is that, to the best of our knowledge, there have been no systematic attempts to rigorously analyze the dynamics of differential equations using persistent homology.

Persistent homology is often used as a means of data reduction. A typical example takes the form of a complicated scalar function defined over a fixed domain, where the geometry of the sub-(super)-level sets is encoded via homology. Of particular interest to us are settings in which the scalar function arises as a solution to a partial differential equation (PDE); we are interested in tracking the evolution of the function, but experimental data only provides information on the level of digital images of the process. Furthermore, capturing the dynamics of a PDE often requires a long time series of rather large digital images. Thus, rather than storing the full images, one can hope to work with a time series of persistence diagrams. Our aim is to draw conclusions about the dynamics of the original PDE from the time series of the persistence diagrams. This is an extremely ambitious goal and far beyond our capabilities at the moment. A much simpler question is the following: if there is an attracting region in the space of persistence diagrams, under what conditions can we conclude that there is a fixed point for the PDE?

This paper represents a first step towards answering the simpler question. Theorem 4.3 shows that given an ordinary differential equation (ODE) with a global compact attractor $\mathcal{A} \subset \mathbb{R}^N$ and a neighborhood in the space of persistence diagrams that is mapped into itself under the dynamics, then there exists a fixed point for the ODE. In applications one could consider the ODE as arising from a finite difference approximation of the PDE.

The challenge is that to obtain results one must understand the topology of $data_P$, the space of data having a fixed persistence diagram P , a topic for which there are only limited results. That the structure of $data_P$ is complicated follows directly from the fact that persistent homology can provide tremendous data reduction, but in a highly nonlinear fashion. With this in mind, the primary goal of this paper is to show that for a reasonable class of problems the space $data_P$ is a finite set of contractible, simplicial sets. The importance of this result is that it opens the possibility of applying standard algebraic topological tools, e.g., Lefschetz fixed point theorem, Conley index, to dynamics that is observed through the lens of persistent homology.

To state our goal precisely requires the introduction of notation. Throughout this paper \mathcal{S}_N denotes the 1-dimensional simplicial complex composed out of N vertices $[i]$ ($i = 1, \dots, N$) and $N - 1$ edges $[i, i + 1]$ ($i = 1, \dots, N - 1$). It is a simplicial decomposition of closed bounded interval in \mathbb{R} .

We study filtrations of \mathcal{S}_N defined as follows.

Definition 1.1 Let $z = (z_1, \dots, z_N) \in \mathbb{R}^N$. Define $f: \mathbb{R}^N \times \mathcal{S}_N \rightarrow \mathbb{R}$ by

$$f(z, \sigma) := \begin{cases} z_j & \text{if } \sigma = [j], \\ \max \{z_j, z_{j+1}\} & \text{if } \sigma = [j, j + 1]. \end{cases}$$

For $r \in \mathbb{R}$, we set $\mathcal{S}_N(z, r) := \{\sigma \in \mathcal{S}_N : f(z, \sigma) \leq r\}$.

Definition 1.2 Given $z = (z_1, \dots, z_N) \in \mathbb{R}^N$, we can reorder the coordinates of z such that

$$z_{j_1} \leq z_{j_2} \leq \dots \leq z_{j_N}.$$

The *sublevel-set filtration* of \mathcal{S}_N at z ,¹ which we write as $\mathcal{S}_N^F(z)$, is given by

$$\mathcal{S}_N(z, z_{j_1}) \subseteq \mathcal{S}_N(z, z_{j_2}) \subseteq \dots \subseteq \mathcal{S}_N(z, z_{j_N}).$$

Because $\mathcal{S}_N^F(z)$ is a finite filtration of simplicial complexes, completely determined by z , we can use classical results from (Edelsbrunner and Harer 2010; Zomorodian and Carlsson 2005) to compute the persistence diagram of $\mathcal{S}_N^F(z)$. We treat this as a map

$$\text{Dgm}: \mathbb{R}^N \rightarrow \text{Per},$$

where Per denotes the space of all persistence diagrams. Thus the space data_P of all $z \in \mathbb{R}^N$ having persistence diagram P is just $\text{Dgm}^{-1}(P)$. We remark that there are a variety of topologies that can be put on Per such that Dgm becomes a continuous map (Chazal et al. 2016; Cohen-Steiner et al. 2007).

Since \mathcal{S}_N is one-dimensional and contractible, we are only concerned with the persistent homology H_0 , i.e., the persistence diagrams associated with connected components. Therefore for the rest of the paper we restrict our study to consist of the family Per of persistence diagrams of level zero.

Here is the main result of this paper.

Theorem 1.3 *For every persistence diagram P , the space $\text{data}_P \subset \mathbb{R}^N$ is composed of a finite number of mutually disjoint components. Each component is contractible, and is homeomorphic to a finite union of convex, potentially unbounded polytopes.*

The proof of Theorem 1.3 is not particularly difficult, but it is technical. We first describe the connected components of data_P ; see Lemma 2.4. In Sect. 2.2, we introduce the poset Str of cellular strings, which are be used to decompose each component as a finite union of convex polytopes in Sect. 2.3. In Sect. 3, we show that the realization of Str is contractible.

To emphasize that Theorem 1.3 is not a trivial result, we use Fig. 1 to demonstrate that data_P is not a convex subset of \mathbb{R}^N . In particular, consider the vectors $v = (v_1, \dots, v_4)$ and $w = (w_1, \dots, w_4)$ on the left of Fig. 1. It is left to the reader to check that $\text{Dgm}(v) = \text{Dgm}(w)$ and that this persistence diagram is given by the pair of black dots (see right of Fig. 1). Note that the vectors in \mathbb{R}^4 , indicated (on the left) in blue stars and red squares, lie on a straight line from v to w . However, the persistence diagrams indicated (on the right) in blue and red clearly differ from $\text{Dgm}(v)$. Thus, the red and blue vectors do not lie in $\text{data}_{\text{Dgm}(v)}$.

In Sect. 4 we apply Theorem 1.3 to prove the existence of fixed points with given persistence diagrams for a dissipative ordinary differential equation.

¹ Analogous results can be obtain for superlevel set filtrations (see Sect. 5).




Fig. 1 Non-convexity of the preimage $data_P$ under the persistence map Dgm . In the left figure the two vectors, $v = (v_1, \dots, v_4)$ and $w = (w_1, \dots, w_4)$, lie in the preimage of the persistence diagram P , composed out of two black points visible on the right figure. Applying Dgm to convex linear combinations of v and w results in a path in the persistence plane illustrated on the right (the convex path is marked in grey, and two sample vectors on the path are marked using red squares and blue stars) (color figure online)

2 Invariants for a fixed persistence diagram

Fix a persistence diagram P . To describe the structure of the space $data_P$, we introduce two levels of invariants: the critical value sequences, representing the connected components of $data_P$, and (for each of these), a partially ordered set Str indexing a polytope decomposition of the component.

2.1 Components

Fix a persistence diagram P . To describe the (finitely many) connected components of $data_P$, it is useful to introduce notation that records the order in which the relevant local maxima and minima occur.

We say that $z = (z_1, \dots, z_N) \in \mathbb{R}^N$ is a *typical point* if its coordinates are distinct. If z is a typical point and $1 < n < N$, we say that z_n is a *local minimum* (of z) if $z_{n-1} > z_n < z_{n+1}$, and a *local maximum* if $z_{n-1} < z_n > z_{n+1}$; it is a *local extremum* if it is a local minimum or maximum. We say that z_1 and z_N are *boundary extrema*; z_1 is a local minimum (resp., maximum) if $z_1 < z_2$ (resp., $z_1 > z_2$).

Definition 2.1 The *critical value sequence* of a typical point $z = (z_1, \dots, z_N)$ is

$$cv(z) = (z_{n_1}, \dots, z_{n_K}) \in \mathbb{R}^K,$$

where the z_{n_k} are the local extrema of z , excluding boundary extrema that are local maxima, and $n_1 < n_2 < \dots < n_K$.

Example 2.2 Let $z = (1.5, -0.9, 1.1, 2.1, 1.4) \in \mathbb{R}^5$. The local minimum is z_2 and the local maximum is z_4 . The boundary extrema are z_1 and z_5 . Since z_1 is also a local maximum we do not include it in the critical value sequence. Thus $h = cv(z) = (-0.9, 2.1, 1.4)$.

The following notion emphasizes the structure of the critical value sequences.

Definition 2.3 A 010 *critical value sequence* of (odd) length K is a vector $cv = (z_1, \dots, z_K) \in \mathbb{R}^K$ with the property that

$$z_{n_1} < z_{n_2} > z_{n_3} < \dots < z_{n_{K-1}} > z_{n_K}.$$

A 101 *critical value sequence* is defined similarly, with the inequalities reversed.

Since we are using sublevel set filtrations to compute the persistence diagram we focus on 010 critical value sequences.

Lemma 2.4 below shows that the local extrema of z are determined up to order by its persistence diagram, and hence that there are only finitely many critical value sequences for any fixed persistence diagram.

Recall that a persistence diagram is a finite collection of *persistence points* $\{p_i = (p_i^b, p_i^d)\}$, where p_i^b and p_i^d denote birth and death values, respectively. Since \mathcal{S}_N is connected, the persistence diagram of a typical point z has a unique persistence point $p_i = (p_i^b, p_i^d)$ such that $p_i^b = \min_{n=1, \dots, N} z_n$ and $p_i^d = \infty$; without loss of generality, we may relabel p_i as p_1 .

Lemma 2.4 *Let $z \in \mathbb{R}^N$ be a typical point with persistence diagram*

$$\{p_m = (p_m^b, p_m^d) \mid m = 1, \dots, M\}.$$

Then, z has $K = 2M - 1$ local extrema; the local minima of z are precisely $\{p_m^b\}_{m=1}^M$ and the interior local maxima of z are precisely $\{p_m^d\}_{m=2}^M$.

We leave the proof of Lemma 2.4 to the reader, remarking that it still holds when $z \in \mathbb{R}^N$ is not a typical point, except that the persistence diagram may be a multiset (there may be multiple copies of a single persistence point).

Given a point z with persistence diagram P , let $C(z)$ denote the component of $data_P$ containing z .

The following lemma shows that $data_P$ is the disjoint union of the finitely many disjoint components $C(z)$, indexed by the critical value sequences. The proof follows from the observation that the order of the local extrema cannot be changed while preserving the persistence diagram.

Lemma 2.5 *If z and z' are typical points in \mathbb{R}^N then $C(z) = C(z')$ if and only if $cv(z) = cv(z')$.*

Moreover, $C(z)$ is the closure of the set of typical points in $C(z)$.

This proves the first assertion in Theorem 1.3.

Remark 2.6 The components $C(z)$ group vectors into equivalence classes that can be characterized using the notion of *chiral merge tree* as defined in Curry (2018). Corollary 5.5 of Curry (2018) shows that the number of chiral merge trees realizing diagram P is equal to $2^{N-1} \prod_{j=2}^N \mu_B(I_j)$, where B is the barcode realization of P , i.e. set of intervals $I_j = [b_j, d_j]$ having the birth and death values of the j -th persistence point as its endpoints, and $\mu_B(I_j)$ is the number of intervals in B that contain I_j .

2.2 Cellular strings

In this section, we define the poset $\text{Str}(N, M)$ of *cellular strings* associated to M points arising from a vector in \mathbb{R}^N . Thus we fix N and M , where $N \geq 2M - 1$.

Consider a string of symbols $s = s_1 \cdots s_N$ of length N , where each symbol s_n is either 0, 1, or X (we refer to 0 and 1 as *bits*). Any such string can be represented as $s = \gamma_1 \cdots \gamma_J$ where each block γ_j is a substring made up of a single symbol (that is, γ_j is $0 \cdots 0$, $1 \cdots 1$, or $X \cdots X$), and consecutive blocks have different symbols. We refer to $s = \gamma_1 \cdots \gamma_J$ as the *canonical representation* of s .

Definition 2.7 Fix $M < N$. A 010 *cellular string*² is a symbol string s of length N such that, for the canonical representation $s = \gamma_1 \cdots \gamma_J$:

- (i) the symbols that make up γ_j and γ_{j+1} are different;
- (ii) γ_1 and γ_J consist of the symbols 0 or X ;
- (iii) if γ_j consists of the symbol X , then the symbol of γ_{j-1} is different from the symbol of γ_{j+1} ;
- (iv) there are exactly M values of j for which γ_j consists of the symbol 0.

The set $\text{Str}(N, M)$ of cellular strings is a poset, where $s' < s$ if the string s is obtained from s' by replacing some of the bits 0 and 1 in s' by X .

The *dimension* of a cellular string s , $\text{dim}(s)$, is the number of symbols X in s . It follows from (iv) that M of the blocks γ_j have the form $0 \cdots 0$, and $M - 1$ have the form $1 \cdots 1$. Thus, $K = 2M - 1$ of the blocks are bitstrings. If these bitstrings are $\gamma_{j_1}, \dots, \gamma_{j_K}$, then the symbol for γ_{j_k} is 0 if k is odd and 1 if k is even. Since each block has at least one symbol, it follows that any cellular string has dimension at most $L = N - K$.

We write $\text{Str}^{(r)}(N, M)$ for the sub-poset of all cellular strings whose first $r - 1$ symbols are X . Note that $\text{Str}(N, M) = \text{Str}^{(1)}(N, M)$ and $\text{Str}^{(L+1)}(N, M) = \{X \cdots X010 \cdots 10\}$.

Proposition 2.8 *An element of $\text{Str}(N, M)$ is maximal if and only if it is an L -dimensional cellular string, where $L = (N - K)$.*

Proof Let $s = \gamma_1 \cdots \gamma_J \in \text{Str}(N, M)$. By definition, $\text{dim}(s) \leq L$. Conversely, suppose that the symbol X appears in s has less than L times. Then some bitstring γ_j has length ≥ 2 . Let s' be the cellular string obtained by replacing the first symbol of γ_j by X . Then $s < s'$, so s is not maximal. □

Since both N and M are fixed in our analysis, we simplify the notation and write Str for $\text{Str}(N, M)$. Figure 2 illustrates the poset Str when $M = 2$, $K = 3$ and $N = 5$; the right column is $\text{Str}^{(2)}$.

Lemma 2.9 *Every string s' is the greatest lower bound of the set of L -dimensional strings s with $s' < s$.*

It follows that Str has the least upper bound property: if two strings have a lower bound, they have a greatest lower bound.

² A 101 cellular string is defined similarly, interchanging 0 and 1.




Fig. 2 The string poset Str for $M = 2$ and $N = 5$. Two-dimensional, one-dimensional, and zero-dimensional strings are surrounded by rectangles, ellipses, and nothing, respectively. The arrows indicate the partial order. The rightmost column is the sub-poset $\text{Str}^{(2)}$

Proof We proceed by downward induction on the dimension d of s' , the case $d = L$ being clear. Consider the canonical representation, $s' = \gamma_1 \cdots \gamma_J$. If $d < L$, then some bitstring γ_j has length ≥ 2 . Consider the strings $s_1 = \gamma_1 \cdots \gamma_{j-1} X \bar{\gamma} \gamma_{j+1} \cdots \gamma_J$ and $s_2 = \gamma_1 \cdots \gamma_{j-1} \bar{\gamma} X \gamma_{j+1} \cdots \gamma_J$ where $\bar{\gamma}$ is a bitstring consisting of the same symbol as γ_j but of length one less than γ_j . Since this is the form of any cellular string s satisfying $s' < s$ and $\dim s = \dim s' + 1$, the result follows. \square

Let s be an L -dimensional cellular string. Successively replacing an X adjacent to a bit (0 or 1) by that bit yields a chain of strings $s = s_L > s_{L-1} > \cdots > s_1 > s_0$. It follows that every maximal chain in the poset has length L .

Example 2.10 Consider a string $s(n) = \sigma_1 0 X \cdots X 1 \sigma_2$ with a block of n consecutive X 's (where σ_1 and σ_2 are fixed substrings). Let $\text{Str}/s(n)$ denote the sub-poset of Str consisting of all strings $s' \leq s(n)$ which begin in $\sigma_1 0$ and end in $1 \sigma_2$. Then $\text{Str}/s(n)$ is isomorphic to the poset I_n of integer intervals $[i, j]$ with $1 \leq i \leq j \leq n + 1$. (The string corresponding to $[i, j]$ is

$$\sigma_1 0 \cdots 0 X \cdots X 1 \cdots 1 \sigma_2;$$

it has i 0's and the first 1 is in the $(j + 1)^{\text{st}}$ spot.)

If s is a cellular string with k blocks of successive X 's (of lengths n_1, \dots, n_k), the sub-poset Str/s of strings $s' < s$ in Str is isomorphic to the product of posets $\text{Str}/s(n_1), \dots, \text{Str}/s(n_k)$, i.e., to the poset

$$I_{n_1} \times \cdots \times I_{n_k}.$$

2.3 The polytopes

We now turn to identifying the polytopes of Theorem 1.3. Fix a 010 critical value sequence $cv = (z_{n_1}, \dots, z_{n_K})$ as in Definition 2.3. To each d -dimensional cellular string s we assign a d -dimensional polytope $T(s)$ in \mathbb{R}^N ; $T(s)$ will be a product of simplices.

Let $s = \gamma_1 \gamma_2 \dots \gamma_J$ be the canonical representation of a string s , as in Definition 2.7. Let n_j denote the length of the substring γ_j , so $N = \sum n_j$.

- If γ_j is either $0 \dots 0$ or $1 \dots 1$, and γ_j is the k^{th} block from the left involving 0 or 1, we set

$$T(\gamma_j) = \{z_k\}^{n_k} = (z_k, \dots, z_k).$$

- If γ_1 is a block $X \dots X$, then

$$T(\gamma_1) = \{(x_1, \dots, x_{n_1}) \in \mathbb{R}^{n_1} : \infty \geq x_1 \geq \dots \geq x_{n_1} \geq z_1\}.$$

- If γ_J is a block $X \dots X$, then

$$T(\gamma_J) = \{(x_1, \dots, x_{n_J}) \in \mathbb{R}^{n_J} : z_k \leq x_1 \leq \dots \leq x_{n_J} \leq \infty\}.$$

- If γ_j is a block $X \dots X$ (for $1 < j < J$), and γ_{j-1} is the k^{th} block from the left involving 0 or 1, then

$$T(\gamma_j) = \{(x_1, \dots, x_{n_j}) \in \mathbb{R}^{n_j}\} \quad \text{where} \quad \begin{cases} z_k \leq x_1 \leq \dots \leq x_{n_j} \leq z_{k+1} & \text{if } k \text{ is odd;} \\ z_k \geq x_1 \geq \dots \geq x_{n_j} \geq z_{k+1} & \text{if } k \text{ is even.} \end{cases}$$

- We define $T(s) \subset \mathbb{R}^N$ to be the concatenation:

$$T(s) = T(\gamma_1 \gamma_2 \dots \gamma_J) = \prod_{j=1}^J T(\gamma_j).$$

Let P be a persistence diagram and $z \in dgm^{-1}(P)$. The component $C(z)$ of $data_P$ is the union of the $T(s)$, where $s \in Str$ and $T(s)$ is defined using the critical value sequence $cv(z)$. This is clear from Definition 2.1.

Since the critical value sequence is always assumed to be fixed, we will suppress it in the notation.

Example 2.11 Consider the case $K = 3$ and $N = 5$. If $s = 01XX0$, then $(\gamma_1, \dots, \gamma_4) = (0, 1, XX, 0)$. So, $(n_1, n_2, n_3, n_4) = (1, 1, 2, 1)$ and hence

$$\begin{aligned} T(01XX0) &= \{z_1\} \times \{z_2\} \times \{(x_1, x_2) : z_2 \geq x_1 \geq x_2 \geq z_3\} \times \{z_3\} \\ &\cong \Delta^0 \times \Delta^0 \times \Delta^2 \times \Delta^0. \end{aligned}$$

If $s = X01X0$, then $(\gamma_1, \dots, \gamma_4, \gamma_5) = (x, 0, 1, x, 0)$. So, $(n_1, n_2, n_3, n_4, n_5) = (1, 1, 1, 1, 1)$ and hence

$$T(X01X0) = [z_1, \infty) \times \{z_1\} \times \{z_2\} \times [z_3, z_2] \times \{z_3\} \cong [0, \infty) \times \Delta^0 \times \Delta^0 \times \Delta^1 \times \Delta^0$$

Similarly, $T(X0100) = [z_1, \infty) \times \{z_1\} \times \{z_2\} \times \{z_3\} \times \{z_3\} \cong [0, \infty) \times \Delta^0 \times \Delta^0 \times \Delta^0 \times \Delta^0$.

Observe that $X0100 < X01X0$ and $T(X0100) \subset T(X01X0)$.

Let **Poly** denote the poset of polytopes in \mathbb{R}^N under inclusion. By definition, T maps strings in **Str** to polytopes in **Poly**.

Lemma 2.12 $T : \text{Str} \rightarrow \text{Poly}$ is an injective poset morphism, and preserves greatest lower bounds.

Proof Suppose that $s' < s$ and $1 + \dim s' = \dim s$. If $s' = \gamma_1 \cdots \gamma_J$ is the canonical form, then some γ_j has the form $a \cdots a$ (where a is 0 or 1), and s has the form

$$s_1 = \gamma_1 \cdots \bar{\gamma}_j X \cdots \gamma_J \quad \text{or} \quad s_2 = \gamma_1 \cdots X \bar{\gamma}_j \gamma_J,$$

where $\bar{\gamma}_j = a \cdots a$ has one fewer bit than γ_j . It is clear from the definition of T that $T(s_1) \neq T(s_2)$, and $T(s')$ is the intersection of $T(s_1)$ and $T(s_2)$, as desired. \square

2.4 Geometric realization of posets

Let C be a poset (partially ordered set). For any $c \in C$, we write C/c for the sub-poset $\{c' : c' \leq c\}$; C is the union of the C/c . If c_1 and c_2 have a greatest lower bound c_{12} , then $(C/c_1) \cap (C/c_2) = C/c_{12}$.

By definition, the geometric realization BC of any poset C is a simplicial complex whose k -dimensional simplices are indexed by the chains $c_0 < c_1 < \cdots < c_k$ of length k in C . It is the union of the realizations $B(C/c)$ of the sub-posets C/c ; if c_1 and c_2 have a greatest lower bound c_{12} , then $B(C/c_1)$ and $B(C/c_2)$ intersect in $B(C/c_{12})$. See Weibel (2013, IV.3.1) for more details.

Here are some basic facts; see Weibel (2013, IV.3) for a discussion. A poset morphism $f : C \rightarrow C'$ determines a continuous map $BC \rightarrow BC'$, and a natural transformation $\eta : f \Rightarrow f'$ between morphisms gives a homotopy $B\eta : BC \rightarrow BC'$ between f and f' . In addition, realization commutes with products: $B(C_1 \times C_2) \cong (BC_1) \times (BC_2)$. Applying these considerations to the poset **Str**, we see that its realization $B\text{Str}$ is the union of the polytopes $B(\text{Str}/s)$, and if s_{12} is the greatest lower bound of s_1 and s_2 then $B(\text{Str}/s_1) \cap B(\text{Str}/s_2)$ is $B(\text{Str}/s_{12})$.

Let s be a cellular string. We saw in Example 2.10 that the poset Str/s is isomorphic to the product $I_{n_1} \times \cdots \times I_{n_k}$ of the posets I_{n_j} of integer intervals in $[1, n_j + 1]$, corresponding to the blocks of n_j successive X 's in s . It is well known that $B(I_n)$ is homeomorphic to the n -simplex Δ^n . Thus

$$B(\text{Str}/s) \cong \prod B(I_{n_j}) \cong \Delta^{n_1} \times \cdots \times \Delta^{n_k}.$$

By construction, $T(s) = \prod T(\gamma_j)$ also has this form. Hence we have a natural homeomorphism

$$B(\text{Str}/s) \cong \prod B(\text{Str}/s(n_j)) \cong \prod B(I_{n_j}) \cong \prod T(\gamma_j) = T(s).$$

Theorem 2.13 *BStr is homeomorphic to $C(z)$.*

Proof By construction, $C(z) = \bigcup T(s)$, and $B\text{Str} = \bigcup B(\text{Str}/s)$. It suffices to observe that for each s_1, \dots, s_n the restriction of the $B\text{Str}/s_i \cong T(s_i)$ induces a homeomorphism between the intersection of the $B(\text{Str}/s_i)$ and the intersection $T(s_i)$. This holds because the two sides are identified with $B(\text{Str}/s')$ and $T(s')$, where s' is the greatest lower bound of the s_i . □

3 Contractibility

We now define a poset morphism $F_1 : \text{Str} \rightarrow \text{Str}$, and modify it to define poset morphisms $F_\ell : \text{Str}^{(\ell)} \rightarrow \text{Str}^{(\ell)}$ for $\ell > 1$.

Definition 3.1 Let s be an L -dimensional cellular string. We define $F_1(s)$ to be the string obtained from s by transposing the first (i.e., leftmost) X with the bit immediately preceding it. If X is the initial symbol, we set $F_1(s) = s$.

If s is a lower-dimensional cellular string, we define $F_1(s)$ as follows. If s has an initial X with no 00 or 11 preceding it, we do as before: transpose X with the bit immediately preceding it, or do nothing if X is the initial symbol. If s begins with a block of $n + 1$ zeroes, say $s = 00 \cdots 0\sigma_2$, we replace the initial 0 by X , so $F_1(s) = X0 \cdots 0\sigma_2$. Otherwise, the string must have the form $s' = \sigma_1abb\sigma_2$, where a, b are bits, $a \neq b$, σ_1 is an (alternating) bitstring not ending in a , and σ_2 is the remainder of the string. We set

$$F_1(s') = \sigma_1aab\sigma_2.$$

The definition of $F_\ell : \text{Str}^{(\ell)} \rightarrow \text{Str}^{(\ell)}$ mimics that of F_1 . Specifically, if $s = \beta\sigma$, where $\beta = X \cdots X$ is a block of length $\ell - 1$ then $F_\ell(s) = \beta F_1(\sigma)$.

Example 3.2 In Fig. 2, the map F_1 sends strings surrounded by rectangles (resp., ellipses) from one column to strings surrounded by rectangles (resp., ellipses) in the second column to the right, while leaving the last column fixed. Thus $F_1(01100) = 00100$ and $F_1(00100) = X0100$.

Since $\text{Str}^{(2)}$ is the rightmost column, the map F_2 acts on this column, mapping strings surrounded by rectangles (resp., ellipses) to those two rows down. Thus $F_2(XX010) = XX010$, $F_2(X0010) = XX010$, and $F_2(XX010) = XX010$.

Lemma 3.3 $F_1 : \text{Str} \rightarrow \text{Str}$ is a poset morphism, and is the identity on the sub-poset $\text{Str}^{(2)}$.

Furthermore, $F_1^K(\text{Str}) = \text{Str}^{(2)}$.

Proof We proceed by downward induction on $d = \dim(s)$ to show that if $s' < s$ then $F_1(s') \leq F_1(s)$. If s' contains an x with no 00 or 11 preceding it, the same is true for s and the inequality is evident.

Next, suppose that $s' = \sigma_1abb \cdots b\sigma_2$, where σ_1a is an alternating bitstring. If $s = \sigma_1abb \cdots b\sigma'_2$ for some $\sigma_2 \leq \sigma'_2$ then

$$F(s') = \sigma_1aab \cdots b\sigma_2 < F(s) = \sigma_1aab \cdots b\sigma'_2.$$

For $s_1 = \sigma_1aXb \cdots$ and $s_2 = \sigma_1ab \cdots bX\sigma_2$, we also have $F(s') < F(s_1)$ and $F(s') < F(s_2)$. Otherwise, either $s_1 < s$ or $s_2 < s$; in these cases, $F_1(s_1) \leq F_1(s)$ or $F_1(s_2) \leq F_1(s)$, by induction, and hence $F(s') < F(s)$.

Finally, if $s' = 00 \cdots 0\sigma$ then either $s_1 = X0 \cdots 0\sigma \leq s$ or else $s_2 = 00 \cdots 0X\sigma \leq s$. By induction, $F_1(s_1) \leq F_1(s)$ or $F_1(s_2) \leq F_1(s)$, so it suffices to observe that $F_1(s') \leq F_1(s_1), F_1(s_2)$. □

Remark 3.4 The proof of Lemma 3.3 also shows that each F_ℓ is a poset morphism.

We can filter the poset Str by sub-posets Fil_i , where $Fil_0 = \text{Str}^{(2)}, Fil_K = \text{Str}$ and Fil_i is the full poset on the set of strings s with $F_1^i(s) \subset \text{Str}^{(2)}$. In Fig. 2, for example, Fil_1 (resp., Fil_2) is the rightmost 3 columns (resp., 5 columns). Since F_1 maps Fil_i to Fil_{i-1} , the geometric realization of BF_1 restricts to a continuous map from $BFil_i$ to $BFil_{i-1}$. We will prove:

Proposition 3.5 *The inclusions $BFil_{i-1} \subseteq BFil_i$ are homotopy equivalences. Hence $B\text{Str}^{(2)} \subseteq B\text{Str}$ is a homotopy equivalence.*

Proof For $i > 0$, we define poset morphisms $F_{1,i} : Fil_i \rightarrow Fil_{i-1} \subseteq Fil_i$ to be the identity on Fil_{i-1} and F_1 otherwise. The geometric realization of $F_{1,i}$ is a continuous map $BFil_i \rightarrow BFil_{i-1} \subseteq BFil_i$ which is the identity on $BFil_{i-1}$.

We will prove that, on geometric realization, $BF_{1,i}$ is homotopic to the identity on $BFil_i$.

We define a poset morphism $h : Fil_i \rightarrow Fil_i$ as follows. If $s \in Fil_{i-1}$ then $h(s) = s$; if $s \notin Fil_{i-1}$, define $h(s)$ to be the greatest lower bound of s and $F_1(s)$. Thus Bh is a continuous map from $BFil_i$ to itself. For $s \in Fil_i$, the inequalities $s \geq h(s) \leq F_{1,i}(s)$ yield natural transformations $\text{id}_i \leftarrow h \Rightarrow F_1$, and hence homotopies between the maps id_i (the identity map on $BFil_i$), Bh and $BF_{1,i}$. □

Corollary 3.6 *Each $B\text{Str}^{(\ell+1)} \subset B\text{Str}^{(\ell)}$ is a homotopy equivalence. In particular, the inclusion of the point $B\text{Str}^{(L+1)}$ in $B\text{Str}$ is a homotopy equivalence, i.e., $B\text{Str}$ is contractible.*

Remark 3.7 We can describe the map $T(s) \rightarrow T(F_1(s))$ induced by F_1 . For example, suppose that $s = \sigma_1\gamma_{j-1}\gamma_j\sigma_2$, where $\sigma_1 = \gamma_1 \cdots \gamma_{j-1}$ is an alternating bitstring of length ≥ 2 and γ_j is a block $X \cdots X$. Then $T(\gamma_{j-1}) = \{z_{j-1}\}$ and $T(\gamma_j) \subset \mathbb{R}^{n_j}$ is defined by inequalities, either $z_{j-1} \leq x_1 \cdots$ or $z_{j-1} \geq x_1 \cdots$, depending on the parity of j . The map F_1 sends $T(\gamma_{j-1}) \times T(\gamma_j)$ to the subset

$$T(X) \times \{z_{j-1}\} \times T(\gamma'),$$

where $T(X)$ is defined by $z_{j-2} \leq x_1 \leq z_{j-1}$ and $T(\gamma')$ is defined by the equations $z_{j-1} \leq x_2 \cdots$ or $z_{j-2} \geq x_1 \cdots$. In effect, the map sends x_1 to z_{j-1} .

4 Existence of fixed points for flows

As an application of Theorem 1.3, we establish the existence of a fixed point solution of an ordinary differential equation whose trajectories are being observed in the space of persistence diagrams. To be more precise consider a differential equation $\dot{z} = f(z)$, $z \in \mathbb{R}^N$, with the property that it possesses a compact global attractor \mathcal{A} (Raugel 2002). Given an initial condition $z(0) = \bar{z} \in \mathbb{R}^N$, we write $z(t) = \varphi(t, \bar{z})$, $t \in [0, \infty)$ for the solution in forward time. The important consequence of the existence of a compact global attractor is that there exists $R > 0$ such that for any initial condition \bar{z} there exists $t_{\bar{z}} > 0$ such that $\|\varphi(t, \bar{z})\| < R$ for all $t \geq t_{\bar{z}}$. We say that R is a bound for \mathcal{A} . Observing the persistence diagrams along a trajectory results in a curve $\text{Dgm}(\varphi(t, \bar{z})) \in \text{Per}$. In what follows we do not assume that we have knowledge of the nonlinearity of f , or of the actual trajectories $\varphi(t, z)$; we are only given the curves $\text{Dgm}(\varphi(t, \bar{z}))$ of persistence diagrams.

Even if the persistence diagram is constant, we cannot conclude that the underlying differential equation has a fixed point. As an example, consider a differential equation in \mathbb{R}^3 with a periodic solution in which the first coordinate $z_1 = 0$ is constant, and (z_2, z_3) oscillates with the property that $1 \leq z_2 \leq z_3$. The associated curve in Per consists of the constant persistence diagram $P = \{(0, \infty)\}$.

However, Theorem 4.3 provides a scenario under which the observation of sufficiently many trajectories suggests the existence of a fixed point for the unknown ordinary differential equation that generates the dynamics. More general theorems are possible and, as will be discussed in a later paper, these techniques can be lifted to the setting of partial differential equations defined on bounded intervals. The purpose of this example is to emphasize the importance of Theorem 1.3 from the perspective of data analysis. Thus, we focus on a much more modest result. We will show that if a particular type of neighborhood in Per is positively invariant under the dynamics, i.e. if $\text{Dgm}(z)$ is in the neighborhood implies that $\text{Dgm}(\varphi(t, z))$ is in the neighborhood for all $t > 0$, then there exists a fixed point for the differential equation that generates the dynamics. To state and obtain such a result requires the introduction of additional notation.

Definition 4.1 We shall say that a persistence diagram


$$P = \{p_m = (p_m^b, p_m^d) : m = 1, \dots, M\}$$

is *sparse* if each persistence point is unique, i.e. $p_m \neq p_n$ for all $m \neq n$.

Given a sparse persistence diagram we can choose $\mu > 0$ such that $\|p_m - p_n\|_\infty \geq 4\mu$ for all $m \neq n$ and $|p_m^d - p_m^b| \geq 4\mu$ for all m .

Example 4.2 A sparse persistence diagram Q is shown in Fig. 3. We can choose $\mu = 0.25$. A possible critical value sequence associated to Q is $\text{cv}(z) = (3, 4.5, 1, 3.5, 2)$.

Fig. 3 A sparse persistence diagram Q with persistence points $\{(1, \infty), (2, 3.5), (3, 4.5)\}$. The boxes indicate the set N_Q for $\mu = 0.25$



We use μ to define subsets of \mathbb{R}^N and Per . We begin by constructing a subset of \mathbb{R}^N using the set of cellular strings $\text{Str}(N, M)$. Choose a point \hat{z} with persistence diagram P . This gives rise to a fixed critical value sequence $\text{cv}(\hat{z})$ and the associated component $C(\hat{z}) \subset \mathbb{R}^N$ of data_P is given by

$$C(\hat{z}) = \bigcup_{s \in \text{Str}(N, M)} T(s).$$

By Theorem 1.3, $C(\hat{z})$ is a contractible union of polytopes.

Let $B_\mu(C(\hat{z})) \subset \mathbb{R}^N$ be the set of points that lie within a distance μ of $C(\hat{z})$ using the sup-norm. The bound on the choice of μ guarantees that if $s', s'' \in \text{Str}(N, M)$ are of maximal dimension and there does not exist $s \in \text{Str}(N, M)$ such that $s < s'$ and $s < s''$, then $B_\mu(T(s'))$ and $B_\mu(T(s''))$ are disjoint. Therefore $B_\mu(C(\hat{z}))$ is contractible.

We now turn to the subset of Per . For each $m = 1, \dots, M$ set

$$P_m := \left\{ p = (p^b, p^d) : \|p - p_m\|_1 \leq \mu \right\}$$

and

$$D := \left\{ p = (p^b, p^d) : p^b \in [p_1^b - \mu, R] \text{ and } 0 \leq p^d - p^b \leq \mu \right\}$$

for some $R > \sup\{p_m^b\} + \mu$. See Fig. 3. Define $N_P \subset \text{Per}$ to be the set of persistence diagrams generated by elements of \mathbb{R}^N with the property that for each $m = 1, \dots, M$ there exists a unique persistence point in P_m and any other persistence points lie in D .

These constructions allow us to prove the following theorem concerning the existence of fixed points of the unknown, underlying dynamical system φ .

Theorem 4.3 Consider a dynamical system generated by an ordinary differential equation that has a global compact attractor \mathcal{A} with a bound R , and whose trajectories

are represented by $\varphi(t, z)$. Let P be a sparse persistence diagram and let $N_P \subset \text{Per}$ be defined as above. Assume that if $\text{Dgm}(z) \in N_P$, then $\text{Dgm}(\varphi(t, z)) \in N_P$ for all $t \geq 0$.

Then, for each component of $\text{Dgm}^{-1}(N_P) \subset \mathbb{R}^N$ there exists a vector \hat{z} such that $\text{Dgm}(\hat{z}) \in N_P$ and $\varphi(t, \hat{z}) = \hat{z}$ for all $t \in \mathbb{R}$, i.e. \hat{z} is a fixed point for the dynamical system.

Proof We begin with the observation that if $z \in B_\mu(C(\hat{z}))$ and there exists $t_1 > 0$ such that $\varphi(t_1, z) \notin B_\mu(C(\hat{z}))$, then there exists $t_0 \in (0, t_1]$ such that $\text{Dgm}(\varphi(t_0, z)) \notin N_P$. This follows from the stability theorem of persistent homology using the bottleneck distance (Cohen-Steiner et al. 2007). This contradicts the hypothesis, therefore, that $B_\mu(C(\hat{z}))$ is a contractible, positively invariant region under the dynamics. By McCord and Mischaikow (1996, Proposition 3.1) the Conley index of the maximal invariant set is that of a hyperbolic attracting fixed point. By McCord (1988, Corollary 5.8) (which utilizes the well known Lefschetz fixed point theorem), the maximal invariant set in $B_\mu(C(\hat{z}))$ contains a fixed point. \square

5 Conclusion and future work

Recall from Remark 2.6 that Curry (2018) provides a count of the contractible components of the preimage of a persistence map. However, to the best of our knowledge, this paper provides the first detailed analysis of the homotopy type of these of a components. Although we have presented the results in the context of sublevel set filtrations, the same arguments can be applied in the setting of superlevel set filtrations. The only significant change is that one needs to use 101 cellular strings; see Definitions 2.3 and 2.7.

Theorem 4.3, and the use of persistence diagrams to obtain results about the dynamics of an ODE, may appear somewhat artificial. However, consider a PDE, such as a reaction diffusion equation, defined on an interval. A finite spatial sampling of the solution at a time point gives rise to a vector. We can think of this vector as arising from two different procedures: (i) numerical, e.g. the values of an ODE derived from a Galerkin approximation to the PDE, or (ii) experimental, e.g. a pixelated image of the solution. Theorem 4.3 is applicable in both cases, and one expects that for fine enough discretization or resolution that the results of Theorem 4.3 will be applicable to the PDE. The example involving images brings us much closer to current treatments of complex spatio-temporal dynamics (Kramar et al. 2016; Levanger et al. 2019). Hence, the natural next step in our research is to obtain an analogous result about existence of fixed points for one-dimensional PDEs whose trajectories are observed in the persistence space.

Finally, the obvious open question as a result of this paper is: given a d -dimensional simplicial complex \mathcal{S} with a function f , similar in form to that of Definition 1.1, can one determine the homology of components of the pre-image of a persistence diagram?

Acknowledgements The work of JC and KM was partially supported by Grants NSF-DMS-1125174, 1248071, 1521771 and a DARPA contract HR0011-16-2-0033. CW was supported by NSF grants 1702233 and 2001417. In addition KM was partially supported by DARPA contract FA8750-17-C-

0054, NIH Grant R01 GM126555-01, and NSF Grants 1934924, 1839294, 1622401, and the NSF HDR TRIPODS award CCF-1934924. In addition JC was partially supported by NAWA Polish Returns Grant PPN/PPO/2018/1/00029.

The first two authors are grateful to an anonymous reviewer of a dramatically different version of this paper for suggesting the relation of our efforts to K -theory, which greatly simplified the proof of Theorem 1.3.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Chazal, F., de Silva, V., Glisse, M., Oudot, S.: The Structure and Stability of Persistence Modules. Springer Briefs in Mathematics. Springer International Publishing AG Switzerland (2016)
- Cohen-Steiner, D., Edelsbrunner, H., Harer, J.: Stability of persistence diagrams. *Discret. Comput. Geom.* **37**(1), 103–120 (2007)
- Curry, J.: The fiber of the persistence map for functions on the interval. *J. Appl. Comput. Topol.* **2**, 301–321 (2018)
- Edelsbrunner, H., Harer, J.: *Computational Topology*. American Mathematical Society, Providence (2010)
- Kramar, M., Levanger, R., Tithof, J., Suri, B., Xu, M., Paul, M., Schatz, M.F., Mischaikow, K.: Analysis of Kolmogorov flow and Rayleigh–Benard convection using persistent homology. *Phys. D-Nonlinear Phenom.* **334**, 82–98 (2016)
- Levanger, R., Xu, M., Cyranka, J., Schatz, M.F., Mischaikow, K., Paul, M.R.: Correlations between the leading Lyapunov vector and pattern defects for chaotic Rayleigh–Benard convection. *Chaos: Interdiscip. J. Nonlinear Sci.* **29**(5), 053103 (2019)
- McCord, C.: Mappings and homological properties in the Conley index theory. *Ergod. Theory Dynam. Syst.* **8***(Charles Conley Memorial Issue), 175–198 (1988)
- McCord, C., Mischaikow, K.: On the global dynamics of attractors for scalar delay equations. *J. Am. Math. Soc.* **9**(4), 1095–1133 (1996)
- Oudot, S.Y.: *Persistence Theory: From Quiver Representations to Data Analysis*. Mathematical Surveys and Monographs, vol. 209. American Mathematical Society, Providence (2015)
- Raugel, G.: Global attractors in partial differential equations. In: Fiedler, B. (ed.) *Handbook of dynamical systems*, vol. 2, pp. 885–982. North-Holland, Amsterdam (2002)
- Weibel, C.A.: *The K -book*. Graduate Studies in Mathematics, vol. 145. American Mathematical Society, Providence (2013). An introduction to algebraic K -theory
- Zomorodian, A., Carlsson, G.: Computing persistent homology. *Discret. Comput. Geom.* **33**(2), 249–274 (2005)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.