

---

---

ERRATA

---

---

**Erratum to: “The Multiple de la Vallé-Poussin Problem on Convex Domains in the Kernel of the Convolution Operator” [Doklady Mathematics 90 (2), 581–583 (2014)]**

**K. R. Zimens<sup>a</sup> and Corresponding Member of the RAS V. V. Napalkov<sup>b</sup>**

(Submitted November 5, 2014; accepted for publication November 5, 2014)

DOI: 10.1134/S1064562414070370

Theorem 3 should read as follows:

**Theorem 3.** *Let  $\mu_k$  be zeros of multiplicity  $s_k$  of a function  $\psi \in H(D)$  such that  $\mu_{k+1} > \mu_k$  and  $\mu_k \in \mathbb{R}_+$ , and let  $\lambda_k$  be zeros of a function  $\varphi(z)$  lying on a ray perpendicular to the tangent line to the domain  $D$  at the limit point of  $\mu_k$  and such that  $\lambda_k \nearrow \infty$  and  $|\arg \lambda_k| < \frac{\pi}{2} - \varepsilon$ ,  $\varepsilon > 0$ .*

*Then  $N_\varphi$  is a sequentially sufficient set in the kernel of the operator  $M_\psi$  in  $P_D$ .*

*K.R. Zimens  
and Corresponding Member of the RAS V.V. Napalkov*

---

<sup>a</sup> *Ufa State Technical University of Aviation,  
Ufa, Bashkortostan, Russia  
e-mail: 15i19@mail.ru, karinazabirova@gmail.com*

<sup>b</sup> *Institute of Mathematics and Computer Center,  
Ufa Scientific Center, Russian Academy of Sciences,  
ul. Chernyshevskogo 112, Ufa, 450077 Bashkortostan, Russia  
e-mail: shaig@anrb.ru*