



Weak decays of doubly heavy baryons: SU(3) analysis

Wei Wang, Zhi-Peng Xing^a, Ji Xu^b

INPAC, Shanghai Key Laboratory for Particle Physics and Cosmology, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China

Received: 24 July 2017 / Accepted: 3 November 2017 / Published online: 23 November 2017

© The Author(s) 2017. This article is an open access publication

Abstract Motivated by the recent LHCb observation of doubly charmed baryon Ξ_{cc}^{++} in the $\Lambda_c^+ K^- \pi^+ \pi^+$ final state, we analyze the weak decays of doubly heavy baryons Ξ_{cc} , Ω_{cc} , Ξ_{bc} , Ω_{bc} , Ξ_{bb} and Ω_{bb} under the flavor SU(3) symmetry. The decay amplitudes for various semileptonic and nonleptonic decays are parametrized in terms of a few SU(3) irreducible amplitudes. We find a number of relations or sum rules between decay widths and CP asymmetries, which can be examined in future measurements at experimental facilities like LHC, Belle II and CEPC. Moreover, once a few decay branching fractions have been measured in the future, some of these relations may provide hints for exploration of new decay modes.

1 Introduction

The existence of doubly heavy baryons is predicted in the quark model, but the experimental search for doubly heavy baryons has been a while [1–6]. Recently, the LHCb collaboration has observed the doubly charmed baryon Ξ_{cc}^{++} with the mass given as [7]

$$m_{\Xi_{cc}^{++}} = (3621.40 \pm 0.72 \pm 0.27 \pm 0.14) \text{ MeV}. \quad (1)$$

Without no doubt, this observation will make a great impact on the hadron spectroscopy and it will also trigger more interests in this research field [8,9]. Moreover after this observation, we also anticipate more experimental investigations of decays of doubly heavy baryons. Thus theoretical studies on weak decays of doubly heavy baryons will be of great importance and are strongly required [10–21].

QCD as the fundamental theory for strong interactions shows two distinct facets. At high energy, the interaction strength is weak that allows the use of perturbation theory. At low energy, quarks and gluons are confined into hadrons.

The large coupling constant prohibits a direct application of perturbative expansions. For a high-energy process with generic hard scattering, one often uses the factorization to separate the high-energy and low-energy degrees of freedoms. The factorization approach has been widely applied to heavy meson decays [22–30], in which the long-distance contributions are parametrized in terms of the low-energy inputs, mostly the light-cone distribution amplitudes. For heavy baryon decays, the factorization analysis is much more involved due to the lack of knowledge on low-energy inputs and the complicated hard-scattering kernels, and see Refs. [31–36] for some recent discussions.

In heavy quark decays, the flavor SU(3) symmetry is a useful tool [37–71]. There are a few advantages to adopt the SU(3) symmetry. First once the branching fractions for a few decay channels have been measured, the flavor SU(3) symmetry offers an opportunity to obtain the knowledge on the related channels. Secondly, the investigation of a few related decay channels can allow one to examine the CKM parameters with the help of SU(3) symmetry. Thirdly, when enough data is available, one may use the data to extract the SU(3) irreducible amplitudes. These amplitudes are expected to calculable in different factorization approaches, and can then be used to examine the factorization schemes themselves. Thus in this paper we will use the flavor SU(3) symmetry and analyze various decays of doubly heavy baryons.

The rest of this paper is organized as follows. In Sect. 2, we will collect the representations for the particle multiplets in the SU(3) symmetry. In Sect. 3, we will analyze the semileptonic decays of the doubly heavy baryons. The nonleptonic decays of doubly charmed baryons, doubly bottom baryons and the baryons with b , c quarks are investigated in Sects. 4, 5 and 6, respectively. The last section contains a brief summary.

2 Particle multiplets

In this section, we will collect the representations for the multiplets of the flavor SU(3) group.

^a e-mail: zpxing@sjtu.edu.cn

^b e-mail: xuj1991@sjtu.edu.cn

Table 1 Quantum numbers for the ground state of doubly heavy baryons. The light quark q corresponds to u, d quark. The J^P denotes the total spin and parity of the baryons. The label S_h^π corresponds to the spin of the heavy quark system

Baryon	Quark content	S_h^π	J^P	Baryon	Quark content	S_h^π	J^P
Ξ_{cc}	{cc}q	1 ⁺	1/2 ⁺	Ξ_{bb}	{bb}q	1 ⁺	1/2 ⁺
Ξ_{cc}^*	{cc}q	1 ⁺	3/2 ⁺	Ξ_{bb}^*	{bb}q	1 ⁺	3/2 ⁺
Ω_{cc}	{cc}s	1 ⁺	1/2 ⁺	Ω_{bb}	{bb}s	1 ⁺	1/2 ⁺
Ω_{cc}^*	{cc}s	1 ⁺	3/2 ⁺	Ω_{bb}^*	{bb}s	1 ⁺	3/2 ⁺
Ξ'_{bc}	{bc}q	0 ⁺	1/2 ⁺	Ω'_{bc}	{bc}s	0 ⁺	1/2 ⁺
Ξ_{bc}	{bc}q	1 ⁺	1/2 ⁺	Ω_{bc}	{bc}s	1 ⁺	1/2 ⁺
Ξ_{bc}^*	{bc}q	1 ⁺	3/2 ⁺	Ω_{bc}^*	{bc}s	1 ⁺	3/2 ⁺




Fig. 1 Anti-triplets (a) and sextets (b) of charmed baryons with one charm quark and two light quarks

Quantum numbers of the doubly heavy baryons are derived from the quark model [72] and are given in Table 1. These baryons can form an SU(3) triplet which are expressed as:

$$T_{cc} = \begin{pmatrix} \Xi_{cc}^{++}(ccu) \\ \Xi_{cc}^+(ccd) \\ \Omega_{cc}^+(ccs) \end{pmatrix}, \quad T_{bc} = \begin{pmatrix} \Xi_{bc}^+(bcu) \\ \Xi_{bc}^0(bcd) \\ \Omega_{bc}^0(bcs) \end{pmatrix},$$

$$T_{bb} = \begin{pmatrix} \Xi_{bb}^0(bbu) \\ \Xi_{bb}^-(bbd) \\ \Omega_{bb}^-(bbs) \end{pmatrix}. \quad (2)$$

The singly charmed baryons can form an antitriplet or sextet as shown in Fig. 1. In the antitriplet case, we have the matrix expression:

$$T_{\bar{c}3} = \begin{pmatrix} 0 & \Lambda_c^+ & \Xi_c^+ \\ -\Lambda_c^+ & 0 & \Xi_c^0 \\ -\Xi_c^+ & -\Xi_c^0 & 0 \end{pmatrix}. \quad (3)$$

For the sextet, we have the multiplet:

$$T_{c6} = \begin{pmatrix} \Sigma_c^{++} & \frac{1}{\sqrt{2}}\Sigma_c^+ & \frac{1}{\sqrt{2}}\Xi_c'^+ \\ \frac{1}{\sqrt{2}}\Sigma_c^+ & \Sigma_c^0 & \frac{1}{\sqrt{2}}\Xi_c'^0 \\ \frac{1}{\sqrt{2}}\Xi_c'^+ & \frac{1}{\sqrt{2}}\Xi_c'^0 & \Omega_c^0 \end{pmatrix}. \quad (4)$$

This is similar for baryons with a bottom quark.

The light baryons form an SU(3) octet and a decuplet. The octet has the expression:

$$T_8 = \begin{pmatrix} \frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda^0 & \Sigma^+ & p \\ \Sigma^- & -\frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda^0 & n \\ \Xi^- & \Xi^0 & -\sqrt{\frac{2}{3}}\Lambda^0 \end{pmatrix}, \quad (5)$$

while the light baryon decuplet is given as

$$(T_{10})^{111} = \Delta^{++}, \quad (T_{10})^{112} = (T_{10})^{121} = (T_{10})^{211} = \frac{1}{\sqrt{3}}\Delta^+,$$

$$(T_{10})^{222} = \Delta^-, \quad (T_{10})^{122} = (T_{10})^{212} = (T_{10})^{221} = \frac{1}{\sqrt{3}}\Delta^0,$$

$$(T_{10})^{113} = (T_{10})^{131} = (T_{10})^{311} = \frac{1}{\sqrt{3}}\Sigma'^+, \quad (T_{10})^{223} = (T_{10})^{232} = (T_{10})^{322} = \frac{1}{\sqrt{3}}\Sigma'^-,$$

$$(T_{10})^{123} = (T_{10})^{132} = (T_{10})^{213} = (T_{10})^{231} = (T_{10})^{312} = (T_{10})^{321} = \frac{1}{\sqrt{6}}\Sigma^0,$$

$$(T_{10})^{133} = (T_{10})^{313} = (T_{10})^{331} = \frac{1}{\sqrt{3}}\Xi'^0, \quad (T_{10})^{233} = (T_{10})^{332} = \frac{1}{\sqrt{3}}\Xi'^-,$$

$$(T_{10})^{333} = \Omega^-. \quad (6)$$

In the meson sector, the light pseudo-scalar meson is an octet, which can be written as:

$$M_8 = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -2\frac{\eta}{\sqrt{6}} \end{pmatrix}, \quad (7)$$

and we will not consider the flavor singlet η_1 in this paper. The following analysis is also applicable to the vector meson octet and other light mesons. The charmed meson forms an SU(3) antitriplet:

$$D_i = (D^0, D^+, D_s^+), \quad (8)$$

and the anticharmed meson forms an SU(3) triplet:

$$\overline{D}^i = (\overline{D}^0, D^-, D_s^-). \quad (9)$$

The above two SU(3) triplets are also applicable to the bottom mesons.

3 Semi-leptonic decays

3.1 Ξ_{cc} and Ω_{cc} decays

The $c \rightarrow q\bar{\ell}\nu$ transition is induced by the effective Hamiltonian:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[V_{cq}^* \bar{q} \gamma^\mu (1 - \gamma_5) c \bar{\nu} \gamma_\mu (1 - \gamma_5) l \right] + h.c., \quad (10)$$

where $q = d, s$ and the V_{cd} and V_{cs} are CKM matrix elements. The heavy-to-light quark operators will form an SU(3) triplet, denoted H_3 with the components $(H_3)^1 = 0$, $(H_3)^2 = V_{cd}^*$, $(H_3)^3 = V_{cs}^*$. At the hadron level, the effective Hamiltonian for decays of Ξ_{cc} and Ω_{cc} into a singly charmed baryon is constructed as:

$$\mathcal{H}_{\text{eff}} = a_1 (T_{cc})^i (H_3)^j (\overline{T}_{c3})_{[ij]} \bar{\nu}_l l + a_2 (T_{cc})^i (H_3)^j (\overline{T}_{c6})_{\{ij\}} \bar{\nu}_l l. \quad (11)$$

Here the a_1 and a_2 are SU(3) irreducible nonperturbative amplitudes. The Feynman diagrams for these decays are given in Fig. 2.

The decay amplitudes for different channels can be deduced from the Hamiltonian in Eq. (11), and given in Table 2. From these amplitudes, we can find the relations for decay widths in the SU(3) symmetry limit:

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Lambda_c^+ l^+ \nu) = \Gamma(\Omega_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu) \\ = \frac{|V_{cd}|^2}{|V_{cs}|^2} \Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu), \quad (12)$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu) = \Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu), \quad (13)$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Sigma_c^+ l^+ \nu) = \Gamma(\Omega_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu) \\ = \frac{1}{2} \Gamma(\Xi_{cc}^+ \rightarrow \Sigma_c^0 l^+ \nu) \\ = \frac{|V_{cd}|^2}{|V_{cs}|^2} \Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu), \quad (14)$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu) = \Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu) \\ = \frac{1}{2} \Gamma(\Omega_{cc}^+ \rightarrow \Omega_c^0 l^+ \nu). \quad (15)$$

Recently, inspired by the LHCb observation of Ξ_{cc} , the weak decays of doubly heavy baryons have been studied in Ref. [21], where the authors first derived the hadronic form factors for these transitions in the light-front approach




Fig. 2 The Feynman diagrams for the semileptonic decays of Ξ_{cc} and Ω_{cc}

Table 2 SU(3) amplitudes for doubly charmed baryons Ξ_{cc} and Ω_{cc} decays into a singly charmed baryon

Channel	Amplitude	Channel	Amplitude
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+ l^+ \nu$	$\frac{a_2 V_{cd}^*}{\sqrt{2}}$	$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ l^+ \nu$	$a_1 V_{cd}^*$
$\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu$	$\frac{a_2 V_{cs}^*}{\sqrt{2}}$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ l^+ \nu$	$a_1 V_{cs}^*$
$\Xi_{cc}^+ \rightarrow \Sigma_c^0 l^+ \nu$	$a_2 V_{cd}^*$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu$	$a_1 V_{cs}^*$
$\Xi_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu$	$\frac{a_2 V_{cs}^*}{\sqrt{2}}$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu$	$-a_1 V_{cd}^*$
$\Omega_{cc}^+ \rightarrow \Xi_c^0 l^+ \nu$	$\frac{a_2 V_{cd}^*}{\sqrt{2}}$		
$\Omega_{cc}^+ \rightarrow \Omega_c^0 l^+ \nu$	$a_2 V_{cs}^*$		

and then applied the results to predict the partial widths for the semi-leptonic and nonleptonic decays of doubly heavy baryons. The SU(3) symmetry can be confronted with these results. We should note that the same comparison in semileptonic Ξ_{bb} , Ω_{bb} and Ξ_{bc} , Ω_{bc} decays and in nonleptonic decays of doubly heavy baryons can also be made. Compared to these explicit model calculations, we found that the SU(3) symmetry works well for the bottom quark decays, while the symmetry breaking effects are sizable for the charm quark decays, largely due to the phase-space differences.

3.2 Semileptonic Ξ_{bb} and Ω_{bb} decays

The b quark decay is controlled by the Hamiltonian

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[V_{q'b\bar{q}'} \bar{q}' \gamma^\mu (1 - \gamma_5) b \bar{l} \gamma_\mu (1 - \gamma_5) l \right] + h.c., \quad (16)$$

with $q' = u, c$. The $b \rightarrow c$ transition is an SU(3) singlet, while the $b \rightarrow u$ transition forms an SU(3) triplet H'_3 with $(H'_3)^1 = 1$ and $(H'_3)^2, (H'_3)^3 = 0$. The Feynman diagrams can be obtained from Fig. 2 by replacing the c quark by b quark, and the final d/s quarks replaced by the c/u quark. The hadron-




Fig. 3 The Feynman diagrams for the semileptonic decays of Ξ_{bb} and Ω_{bb}

Table 3 Similar to Table 2 but for the doubly bottom baryons

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^- \rightarrow \Lambda_b^0 l^- \bar{\nu}$	$-a_4 V_{ub}$	$\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ l^- \bar{\nu}$	$a_3 V_{cb}$
$\Omega_{bb}^- \rightarrow \Xi_b^0 l^- \bar{\nu}$	$-a_4 V_{ub}$	$\Xi_{bb}^- \rightarrow \Xi_{bc}^0 l^- \bar{\nu}$	$a_3 V_{cb}$
$\Xi_{bb}^0 \rightarrow \Sigma_b^+ l^- \bar{\nu}$	$a_5 V_{ub}$	$\Omega_{bb}^- \rightarrow \Omega_{bc}^0 l^- \bar{\nu}$	$a_3 V_{cb}$
$\Xi_{bb}^- \rightarrow \Sigma_b^0 l^- \bar{\nu}$	$\frac{a_5 V_{ub}}{\sqrt{2}}$		
$\Omega_{bb}^- \rightarrow \Xi_b^0 l^- \bar{\nu}$	$\frac{a_5 V_{ub}}{\sqrt{2}}$		

level Hamiltonian for the semileptonic Ξ_{bb} and Ω_{bb} decays is constructed as

$$H_{\text{eff}} = a_3(T_{bb})^i(\bar{T}_{bc})_i \bar{l} \nu_l + a_4(T_{bb})^i(H'_3)^j(\bar{T}_{b\bar{3}})_{[ij]} \bar{l} \nu_l + a_5(T_{bb})^i(H'_3)^j(\bar{T}_{b6})_{\{ij\}} \bar{l} \nu_l \quad (17)$$

The Feynman diagrams for these decays are given in Fig. 3. The decay amplitudes can be deduced from this Hamiltonian, and the results are given in Table 3. It leads to the relations for the decay widths:

$$\Gamma(\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ l^- \bar{\nu}) = \Gamma(\Xi_{bb}^- \rightarrow \Xi_{bc}^0 l^- \bar{\nu}) = \Gamma(\Omega_{bb}^- \rightarrow \Omega_{bc}^0 l^- \bar{\nu}), \quad (18)$$

$$\Gamma(\Xi_{bb}^- \rightarrow \Lambda_b^0 l^- \bar{\nu}) = \Gamma(\Omega_{bb}^- \rightarrow \Xi_b^0 l^- \bar{\nu}), \quad (19)$$

$$\Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^+ l^- \bar{\nu}) = 2\Gamma(\Xi_{bb}^- \rightarrow \Sigma_b^0 l^- \bar{\nu}) = 2\Gamma(\Omega_{bb}^- \rightarrow \Xi_b^0 l^- \bar{\nu}). \quad (20)$$

3.3 Semileptonic Ξ_{bc} and Ω_{bc} decays

The effective Hamiltonian for the semileptonic Ξ_{bc} and Ω_{bc} decays is given by

$$H_{\text{eff}} = a_6(T_{bc})^i(\bar{T}_{cc})_i \bar{l} \nu_l + a_7(T_{bc})^i(H'_3)^j(\bar{T}_{c\bar{3}})_{[ij]} \bar{l} \nu_l + a_8(T_{bc})^i(H'_3)^j(\bar{T}_{c6})_{\{ij\}} \bar{l} \nu_l$$

Table 4 Similar to Table 2 but for the doubly heavy bcq baryons

Channel	Amplitude	Channel	Amplitude
$\Xi_{bc}^+ \rightarrow \Lambda_b^0 l^+ \nu$	$a_9 V_{cd}^*$	$\Xi_{bc}^+ \rightarrow \Xi_{cc}^{++} l^- \bar{\nu}$	$a_6 V_{cb}$
$\Xi_{bc}^+ \rightarrow \Xi_b^0 l^+ \nu$	$a_9 V_{cs}^*$	$\Xi_{bc}^0 \rightarrow \Xi_{cc}^+ l^- \bar{\nu}$	$a_6 V_{cb}$
$\Xi_{bc}^0 \rightarrow \Xi_b^- l^+ \nu$	$a_9 V_{cs}^*$	$\Omega_{bc}^0 \rightarrow \Omega_{cc}^+ l^- \bar{\nu}$	$a_6 V_{cb}$
$\Omega_{bc}^0 \rightarrow \Xi_b^- l^+ \nu$	$-a_9 V_{cd}^*$	$\Xi_{bc}^0 \rightarrow \Lambda_c^+ l^- \bar{\nu}$	$-a_7 V_{ub}$
$\Xi_{bc}^+ \rightarrow \Sigma_b^0 l^+ \nu$	$\frac{a_{10} V_{cd}^*}{\sqrt{2}}$	$\Omega_{bc}^0 \rightarrow \Xi_c^+ l^- \bar{\nu}$	$-a_7 V_{ub}$
$\Xi_{bc}^+ \rightarrow \Xi_b^0 l^+ \nu$	$\frac{a_{10} V_{cs}^*}{\sqrt{2}}$	$\Xi_{bc}^+ \rightarrow \Sigma_c^{++} l^- \bar{\nu}$	$a_8 V_{ub}$
$\Xi_{bc}^0 \rightarrow \Sigma_b^- l^+ \nu$	$a_{10} V_{cd}^*$	$\Xi_{bc}^0 \rightarrow \Sigma_c^+ l^- \bar{\nu}$	$\frac{a_8 V_{ub}}{\sqrt{2}}$
$\Xi_{bc}^0 \rightarrow \Xi_b' l^+ \nu$	$\frac{a_{10} V_{cs}^*}{\sqrt{2}}$	$\Omega_{bc}^0 \rightarrow \Xi_c' l^- \bar{\nu}$	$\frac{a_8 V_{ub}}{\sqrt{2}}$
$\Omega_{bc}^0 \rightarrow \Xi_b' l^+ \nu$	$\frac{a_{10} V_{cd}^*}{\sqrt{2}}$		
$\Omega_{bc}^0 \rightarrow \Omega_b^- l^+ \nu$	$a_{10} V_{cs}^*$		

$$+ a_9(T_{bc})^i(H_3)^j(\bar{T}_{b\bar{3}})_{[ij]} \bar{l} \nu_l + a_{10}(T_{bc})^i(H_3)^j(\bar{T}_{b6})_{\{ij\}} \bar{l} \nu_l. \quad (21)$$

In this equation, we have included both charm quark and bottom quark decays. The decay amplitudes for different channels are obtained by expanding the above Hamiltonian and are collected in Table 4.

Apparently, the Ξ_{bc} and Ω_{bc} decay amplitudes can be obtained by the ones for T_{cc} and T_{bb} decays. For the charm quark decays, one would derive the results with the replacement, $T_{cc} \rightarrow T_{bc}$, $T_c \rightarrow T_b$. The replacement in bottom quark decays is $T_{bb} \rightarrow T_{bc}$, $T_b \rightarrow T_c$. Thus we have the following relations for the decay widths:

$$\Gamma(\Xi_{bc}^+ \rightarrow \Lambda_b^0 l^+ \nu) = \Gamma(\Omega_{bc}^0 \rightarrow \Xi_b^- l^+ \nu) = \frac{|V_{cd}|^2}{|V_{cs}|^2} \Gamma(\Xi_{bc}^+ \rightarrow \Xi_b^0 l^+ \nu), \quad (22)$$

$$\Gamma(\Xi_{bc}^+ \rightarrow \Xi_b^0 l^+ \nu) = \Gamma(\Xi_{bc}^0 \rightarrow \Xi_b^- l^+ \nu), \quad (23)$$

$$\begin{aligned} \Gamma(\Xi_{bc}^+ \rightarrow \Sigma_b^0 l^+ \nu) &= \Gamma(\Omega_{bc}^0 \rightarrow \Xi_b' l^+ \nu) \\ &= \frac{1}{2} \Gamma(\Xi_{bc}^0 \rightarrow \Sigma_b^- l^+ \nu) \\ &= \frac{|V_{cd}|^2}{|V_{cs}|^2} \Gamma(\Xi_{bc}^+ \rightarrow \Xi_b' l^+ \nu), \end{aligned} \quad (24)$$

$$\begin{aligned} \Gamma(\Xi_{bc}^+ \rightarrow \Xi_b' l^+ \nu) &= \Gamma(\Xi_{bc}^0 \rightarrow \Xi_b' l^+ \nu) \\ &= \frac{1}{2} \Gamma(\Omega_{bc}^0 \rightarrow \Omega_b^- l^+ \nu), \end{aligned} \quad (25)$$

$$\begin{aligned} \Gamma(\Xi_{bc}^+ \rightarrow \Xi_{cc}^{++} l^- \bar{\nu}) &= \Gamma(\Xi_{bc}^0 \rightarrow \Xi_{cc}^+ l^- \bar{\nu}) \\ &= \Gamma(\Omega_{bc}^0 \rightarrow \Omega_{cc}^+ l^- \bar{\nu}), \end{aligned} \quad (26)$$

$$\Gamma(\Xi_{bc}^0 \rightarrow \Lambda_c^+ l^- \bar{\nu}) = \Gamma(\Omega_{bc}^0 \rightarrow \Xi_c^+ l^- \bar{\nu}), \quad (27)$$

$$\begin{aligned}\Gamma(\Xi_{bc}^+ \rightarrow \Sigma_c^{++} l^- \bar{\nu}) &= 2\Gamma(\Xi_{bc}^0 \rightarrow \Sigma_c^+ l^- \bar{\nu}) \\ &= 2\Gamma(\Omega_{bc}^0 \rightarrow \Xi_c^+ l^- \bar{\nu}).\end{aligned}\quad (28)$$

4 Nonleptonic Ξ_{cc} and Ω_{cc} decays

Usually the charm quark decays into light quarks are categorized into three groups: Cabibbo-allowed, singly Cabibbo-suppressed, and doubly Cabibbo-suppressed:

$$c \rightarrow s\bar{d}u, \quad c \rightarrow u\bar{d}d/\bar{s}s, \quad c \rightarrow d\bar{s}u. \quad (29)$$

The tree operators transform under the flavor SU(3) symmetry as $\mathbf{3} \otimes \bar{\mathbf{3}} \otimes \mathbf{3} = \mathbf{3} \oplus \mathbf{3} \oplus \bar{\mathbf{6}} \oplus \mathbf{15}$. So the Hamiltonian can be decomposed in terms of a vector (H_3), a traceless tensor antisymmetric in upper indices, $H_{\bar{6}}$, and a traceless tensor symmetric in upper indices, H_{15} . As we will show in the following, the vector representation H_3 will vanish as an approximation.

For the $c \rightarrow s\bar{d}u$ transition, we have

$$(H_{\bar{6}})_2^{31} = -(H_{\bar{6}})_2^{13} = 1, \quad (H_{15})_2^{31} = (H_{15})_2^{13} = 1, \quad (30)$$

while, for the $c \rightarrow d\bar{s}u$ transition, which is doubly Cabibbo-suppressed, we have

$$(H_{\bar{6}})_3^{21} = -(H_{\bar{6}})_3^{12} = \sin^2 \theta_C, \quad (H_{15})_3^{21} = (H_{15})_3^{12} = \sin^2 \theta_C. \quad (31)$$

For the transition $c \rightarrow u\bar{d}d$, we have

$$\begin{aligned}(H_3)^1 &= 1, \quad (H_{\bar{6}})_2^{21} = -(H_{\bar{6}})_2^{12} = (H_{\bar{6}})_3^{13} \\ &= -(H_{\bar{6}})_3^{31} = \frac{1}{2}, \\ \frac{1}{3}(H_{15})_2^{21} &= \frac{1}{3}(H_{15})_2^{12} = -\frac{1}{2}(H_{15})_1^{11} = -(H_{15})_3^{13} \\ &= -(H_{15})_3^{31} = \frac{1}{4},\end{aligned}\quad (32)$$

with all other remaining entries zero. The overall factor is $V_{cd}^* V_{ud} \simeq -\sin(\theta_C)$. Meanwhile, for the transition $c \rightarrow \bar{u}s$, we have

$$\begin{aligned}(H_3)^1 &= 1, \quad (H_{\bar{6}})_3^{31} = -(H_{\bar{6}})_3^{13} \\ &= (H_{\bar{6}})_2^{12} = -(H_{\bar{6}})_2^{21} = \frac{1}{2}, \\ \frac{1}{3}(H_{15})_3^{31} &= \frac{1}{3}(H_{15})_3^{13} = -\frac{1}{2}(H_{15})_1^{11} \\ &= -(H_{15})_2^{12} = -(H_{15})_2^{21} = \frac{1}{4},\end{aligned}\quad (33)$$

with all other remaining entries zero. The overall factor is $V_{cs}^* V_{us} \simeq \sin(\theta_C)$. With both the $c \rightarrow u\bar{d}d$ and the $c \rightarrow \bar{u}s$, the singly Cabibbo-suppressed channel has the following effective Hamiltonian:

$$\begin{aligned}(H_{\bar{6}})_3^{31} &= -(H_{\bar{6}})_3^{13} = (H_{\bar{6}})_2^{12} = -(H_{\bar{6}})_2^{21} = \sin(\theta_C), \\ (H_{15})_3^{31} &= (H_{15})_3^{13} = -(H_{15})_2^{12} \\ &= -(H_{15})_2^{21} = \sin(\theta_C).\end{aligned}\quad (34)$$

4.1 Decays into a charmed baryon and a light meson

With the above expressions, one may derive the effective Hamiltonian for decays involving the antitriplet heavy baryons as

$$\begin{aligned}\mathcal{H}_{\text{eff}} = b_3(T_{cc})^i &(\bar{T}_{c\bar{3}})_{[ij]} M_l^k (H_{\bar{6}})_k^{jl} \\ &+ b_4(T_{cc})^i (\bar{T}_{c\bar{3}})_{[jl]} M_i^k (H_{\bar{6}})_k^{jl} \\ &+ b_5(T_{cc})^i (\bar{T}_{c\bar{3}})_{[jk]} M_l^k (H_{\bar{6}})_i^{jl} \\ &+ b_6(T_{cc})^i (\bar{T}_{c\bar{3}})_{[ij]} M_l^k (H_{15})_k^{jl} \\ &+ b_7(T_{cc})^i (\bar{T}_{c\bar{3}})_{[jk]} M_l^k (H_{15})_i^{jl}.\end{aligned}\quad (35)$$

For the sextet baryon, we have the Hamiltonian

$$\begin{aligned}\mathcal{H}_{\text{eff}} = b_{10}(T_{cc})^i &(\bar{T}_{c6})_{\{ij\}} M_l^k (H_{15})_k^{jl} \\ &+ b_{11}(T_{cc})^i (\bar{T}_{c6})_{\{jl\}} M_i^k (H_{15})_k^{jl} \\ &+ b_{12}(T_{cc})^i (\bar{T}_{c6})_{\{jk\}} M_l^k (H_{15})_i^{jl} \\ &+ b_{13}(T_{cc})^i (\bar{T}_{c6})_{\{ij\}} M_l^k (H_{\bar{6}})_l^{jk} \\ &+ b_{14}(T_{cc})^i (\bar{T}_{c6})_{\{jk\}} M_l^k (H_{\bar{6}})_i^{jl}.\end{aligned}\quad (36)$$

The Feynman diagrams for these decays are given in Fig. 4.




Fig. 4 The Feynman diagrams for Ξ_{cc} and Ω_{cc} decays into a charmed baryon and a light meson

Table 5 Doubly charmed baryons decays into a cqq (antitriplet) and a light meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$	$b_3 - 2b_4 + b_6$	$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+$	$(b_3 - 2b_4 + b_6)(-\sin(\theta_C))$
$\Xi_c^+ \rightarrow \Lambda_c^+ \bar{K}^0$	$b_3 - b_5 - b_6 + b_7$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+$	$(b_3 - 2b_4 + b_6)\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Xi_c^+ \pi^0$	$\frac{2b_4 - b_5 - b_7}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Lambda_c^+ \pi^0$	$\frac{(b_3 - 2b_4 - b_6 + 2b_7)\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Xi_c^+ \eta$	$\frac{-2b_4 + b_5 - 3b_7}{\sqrt{6}}$	$\Xi_{cc}^+ \rightarrow \Lambda_c^+ \eta$	$\frac{(-3b_3 + 2b_4 + 2b_5 + 3b_6)\sin(\theta_C)}{\sqrt{6}}$
$\Xi_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$b_3 - b_5 + b_6 - b_7$	$\Xi_{cc}^+ \rightarrow \Xi_c^+ K^0$	$(2b_4 - b_5 + b_7)(-\sin(\theta_C))$
$\Omega_{cc}^+ \rightarrow \Xi_c^+ \bar{K}^0$	$b_3 - 2b_4 - b_6$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+$	$(b_3 - b_5 + b_6 - b_7)\sin(\theta_C)$
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^+$	$(b_3 - 2b_4 + b_6)\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Lambda_c^+ \bar{K}^0$	$(2b_4 - b_5 + b_7)\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^0$	$(b_3 - 2b_4 - b_6)\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi_c^+ \pi^0$	$\frac{(b_3 - b_5 - b_6 - b_7)\sin(\theta_C)}{\sqrt{2}}$
$\Omega_{cc}^+ \rightarrow \Lambda_c^+ \pi^0$	$-\sqrt{2}b_7 \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi_c^+ \eta$	$\frac{(-3b_3 + 4b_4 + b_5 + 3b_6 - 3b_7)\sin(\theta_C)}{\sqrt{6}}$
$\Omega_{cc}^+ \rightarrow \Lambda_c^+ \eta$	$\sqrt{\frac{2}{3}}(2b_4 - b_5)\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$(b_3 - b_5 + b_6 - b_7)\sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow \Xi_c^+ K^0$	$(b_3 - b_5 - b_6 + b_7)\sin^2(\theta_C)$		
$\Omega_{cc}^+ \rightarrow \Xi_c^0 K^+$	$(b_3 - b_5 + b_6 - b_7)(-\sin^2(\theta_C))$		

Table 6 Doubly charmed baryons decays into a cqq (sextet) and a light meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \bar{K}^0$	$b_{10} - b_{13}$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \pi^0$	$\frac{(b_{10} - b_{13})\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^{++} \rightarrow \Xi_c' \pi^+$	$\frac{b_{10} + 2b_{11} + b_{13}}{\sqrt{2}}$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \eta$	$-\sqrt{\frac{3}{2}}(b_{10} - b_{13})\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Sigma_c^{++} K^-$	$b_{12} - b_{14}$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^+ \pi^+$	$-\frac{(b_{10} + 2b_{11} + b_{13})\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Sigma_c^+ \bar{K}^0$	$\frac{b_{10} + b_{12} - b_{13} - b_{14}}{\sqrt{2}}$	$\Xi_{cc}^{++} \rightarrow \Xi_c' K^+$	$\frac{(b_{10} + 2b_{11} + b_{13})\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Xi_c' \pi^0$	$\frac{1}{2}(-2b_{11} + b_{12} + b_{14})$	$\Xi_{cc}^+ \rightarrow \Sigma_c^{++} \pi^-$	$(b_{14} - b_{12})\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Xi_c' \eta$	$\frac{2b_{11} - b_{12} + 3b_{14}}{2\sqrt{3}}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^+ \pi^0$	$\frac{1}{2}(b_{10} + 2b_{11} - b_{13} - 2b_{14})\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\frac{b_{10} + b_{12} + b_{13} + b_{14}}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^+ \eta$	$-\frac{(3b_{10} + 2b_{11} + 2b_{12} - 3b_{13})\sin(\theta_C)}{2\sqrt{3}}$
$\Xi_{cc}^+ \rightarrow \Omega_c^0 K^+$	$b_{12} + b_{14}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^0 \pi^+$	$(b_{10} + b_{12} + b_{13} + b_{14})(-\sin(\theta_C))$
$\Omega_{cc}^+ \rightarrow \Xi_c' \bar{K}^0$	$\frac{b_{10} + 2b_{11} - b_{13}}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Xi_c^+ K^0$	$\frac{(2b_{11} - b_{12} + b_{14})\sin(\theta_C)}{\sqrt{2}}$
$\Omega_{cc}^+ \rightarrow \Omega_c^0 \pi^+$	$b_{10} + b_{13}$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+$	$\frac{(b_{10} - b_{12} + b_{13} - b_{14})\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} K^0$	$(b_{10} - b_{13})\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++} K^-$	$(b_{12} - b_{14})\sin(\theta_C)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+ K^+$	$\frac{(b_{10} + 2b_{11} + b_{13})\sin^2(\theta_C)}{\sqrt{2}}$	$\Omega_{cc}^+ \rightarrow \Sigma_c^+ \bar{K}^0$	$-\frac{(2b_{11} - b_{12} + b_{14})\sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Sigma_c^+ K^0$	$\frac{(b_{10} + 2b_{11} - b_{13})\sin^2(\theta_C)}{\sqrt{2}}$	$\Omega_{cc}^+ \rightarrow \Xi_c' \pi^0$	$\frac{1}{2}(b_{10} + b_{12} - b_{13} + b_{14})\sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Sigma_c^0 K^+$	$(b_{10} + b_{13})\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi_c' \eta$	$-\frac{(3b_{10} + 4b_{11} + b_{12} - 3b_{13} - 3b_{14})\sin(\theta_C)}{2\sqrt{3}}$
$\Omega_{cc}^+ \rightarrow \Sigma_c^{++} \pi^-$	$(b_{12} - b_{14})\sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$-\frac{(b_{10} - b_{12} + b_{13} - b_{14})\sin(\theta_C)}{\sqrt{2}}$
$\Omega_{cc}^+ \rightarrow \Sigma_c^+ \pi^0$	$b_{14} \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Omega_c^0 K^+$	$(b_{10} + b_{12} + b_{13} + b_{14})\sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow \Sigma_c^+ \eta$	$\frac{(b_{12} - 2b_{11})\sin^2(\theta_C)}{\sqrt{3}}$		
$\Omega_{cc}^+ \rightarrow \Sigma_c^0 \pi^+$	$(b_{12} + b_{14})\sin^2(\theta_C)$		
$\Omega_{cc}^+ \rightarrow \Xi_c' K^0$	$\frac{(b_{10} + b_{12} - b_{13} - b_{14})\sin^2(\theta_C)}{\sqrt{2}}$		
$\Omega_{cc}^+ \rightarrow \Xi_c^0 K^+$	$\frac{(b_{10} + b_{12} + b_{13} + b_{14})\sin^2(\theta_C)}{\sqrt{2}}$		

Expanding the above equations, we will obtain the decay amplitudes given in Table 5 for the antitriplet baryon and Table 6 for the sextet. Thus we have the following relations for decay widths:

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+) = \Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+), \quad (37)$$

$$\Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^+ K^0) = \Gamma(\Omega_{cc}^+ \rightarrow \Lambda_c^+ \bar{K}^0), \quad (38)$$

$$\Gamma(\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+) = \Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+). \quad (39)$$

For the decays into the sextet, we have

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \pi^0) = \frac{1}{3} \Gamma(\Xi_{cc}^{++} \rightarrow \Sigma_c^{++} \eta), \quad (40)$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Sigma_c^+ \pi^+) = \Gamma(\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+), \quad (41)$$

$$\Gamma(\Xi_{cc}^+ \rightarrow \Sigma_c^{++} \pi^-) = \Gamma(\Omega_{cc}^+ \rightarrow \Sigma_c^{++} K^-), \quad (42)$$

$$\Gamma(\Xi_{cc}^+ \rightarrow \Sigma_c^0 \pi^+) = \Gamma(\Omega_{cc}^+ \rightarrow \Sigma_c^0 K^+), \quad (43)$$

$$\Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^0 K^0) = \Gamma(\Omega_{cc}^+ \rightarrow \Sigma_c^+ \bar{K}^0), \quad (44)$$

$$\Gamma(\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+) = \Gamma(\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+). \quad (45)$$




Fig. 5 The Feynman diagrams for Ξ_{cc} and Ω_{cc} decays into a light baryon and a charmed meson

Table 7 Doubly charmed baryons decays into a light baryon in the octet and a charmed meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{cc}^{++} \rightarrow \Sigma^+ D^+$	$-c_7 - c_9$	$\Xi_{cc}^{++} \rightarrow \Sigma^+ D_s^+$	$(-c_7 - c_9) \sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Lambda^0 D^+$	$\frac{-c_6 - c_7 + 3c_8 + 3c_9}{\sqrt{6}}$	$\Xi_{cc}^+ \rightarrow p D^+$	$(-c_7 - c_9) \sin(\theta_C)$
$\Xi_{cc}^+ \rightarrow \Sigma^+ D^0$	$-c_6 - c_8$	$\Xi_{cc}^+ \rightarrow \Lambda^0 D_s^+$	$\frac{(2c_6 - c_7 + 3c_9) \sin(\theta_C)}{\sqrt{6}}$
$\Xi_{cc}^+ \rightarrow \Sigma^0 D^+$	$\frac{c_6 + c_7 + c_8 + c_9}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Sigma^0 D_s^+$	$\frac{(c_7 + 2c_8 + c_9) \sin(\theta_C)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Xi^0 D_s^+$	$-c_6 + c_8$	$\Xi_{cc}^+ \rightarrow p D^0$	$(-c_6 - c_8) \sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow \Xi^0 D^+$	$-c_7 + c_9$	$\Xi_{cc}^+ \rightarrow n D^+$	$(-c_6 - c_7 + c_8 + c_9) \sin(\theta_C)$
$\Xi_{cc}^{++} \rightarrow p D_s^+$	$(c_7 + c_9) \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Lambda^0 D^+$	$\frac{-(c_6 - 2c_7 - 3c_8) \sin(\theta_C)}{\sqrt{6}}$
$\Xi_{cc}^+ \rightarrow n D_s^+$	$(c_7 - c_9) \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Sigma^+ D^0$	$(-c_6 - c_8) \sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow \Lambda^0 D_s^+$	$\sqrt{\frac{2}{3}} (-c_6 - c_7) \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Sigma^0 D^+$	$\frac{(c_6 + c_8 + 2c_9) \sin(\theta_C)}{\sqrt{2}}$
$\Omega_{cc}^+ \rightarrow \Sigma^0 D_s^+$	$-\sqrt{2} (c_8 + c_9) \sin^2(\theta_C)$	$\Omega_{cc}^+ \rightarrow \Xi^0 D_s^+$	$(-c_6 - c_7 + c_8 + c_9) \sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow p D^0$	$(c_6 + c_8) \sin^2(\theta_C)$		
$\Omega_{cc}^+ \rightarrow n D^+$	$(c_6 - c_8) \sin^2(\theta_C)$		

4.2 Decays into a light octet baryon and a charmed meson

The effective Hamiltonian for the decays of T_{cc} into a light octet baryon and a charmed meson is given as

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & c_4(T_{cc})^l \bar{D}^m \epsilon_{ijk}(T_8)_l^k (H_6)_m^{ij} \\ & + c_5(T_{cc})^l \bar{D}^m \epsilon_{ijk}(T_8)_m^k (H_6)_l^{ij} \\ & + c_6(T_{cc})^l \bar{D}^i \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm} \\ & + c_7(T_{cc})^i \bar{D}^l \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm} \\ & + c_8(T_{cc})^l \bar{D}^i \epsilon_{ijk}(T_8)_m^k (H_{15})_l^{jm} \\ & + c_9(T_{cc})^i \bar{D}^l \epsilon_{ijk}(T_8)_m^k (H_{15})_l^{jm}. \end{aligned} \quad (46)$$

In the above Hamiltonian we find the following relations:

$$(T_{cc})^l \bar{D}^m \epsilon_{ijk}(T_8)_l^k (H_6)_m^{ij} = -2(T_{cc})^i \bar{D}^l \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm},$$

$$(T_{cc})^l \bar{D}^m \epsilon_{ijk}(T_8)_m^k (H_6)_l^{ij} = -2(T_{cc})^l \bar{D}^i \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm}. \quad (47)$$

Thus two of the reduced matrix elements are not independent. In the following, we will eliminate the c_4 and c_5 and use the effective Hamiltonian:

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & c_6(T_{cc})^l \bar{D}^i \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm} \\ & + c_7(T_{cc})^i \bar{D}^l \epsilon_{ijk}(T_8)_m^k (H_6)_l^{jm} \\ & + c_8(T_{cc})^l \bar{D}^i \epsilon_{ijk}(T_8)_m^k (H_{15})_l^{jm} \\ & + c_9(T_{cc})^i \bar{D}^l \epsilon_{ijk}(T_8)_m^k (H_{15})_l^{jm}. \end{aligned} \quad (48)$$

The Feynman diagrams for these decays are given in Fig. 5. Expanding the above equations, we will obtain the decay amplitudes given in Table 7, which leads to the relations for decay widths:

$$\Gamma(\Xi_{cc}^{++} \rightarrow p D^+) = \Gamma(\Xi_{cc}^{++} \rightarrow \Sigma^+ D_s^+),$$

$$\Gamma(\Omega_{cc}^+ \rightarrow \Sigma^+ D^0) = \Gamma(\Xi_{cc}^+ \rightarrow p D^0). \quad (49)$$

4.3 Decays into a light decuplet baryon and a charmed meson

The effective Hamiltonian for a light decuplet in the final state is given as

$$\mathcal{H}_{\text{eff}} = d_4(T_{cc})^l \bar{D}^m (T_{10})_{ijl} (H_{15})_m^{ij} + d_5(T_{cc})^l \bar{D}^m (T_{10})_{ijm} (H_{15})_l^{ij}. \quad (50)$$

The Feynman diagrams for these decays are the same as in Fig. 5. The corresponding decay amplitudes are given in Table 8 and it leads to the relations for decay widths:

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Delta^+ D^+) = \Gamma(\Xi_{cc}^{++} \rightarrow \Sigma'^+ D_s^+), \quad (51)$$

Table 8 Doubly charmed baryons decays into a light baryon in the decuplet and a charmed meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{cc}^{++} \rightarrow \Sigma'^+ D^+$	$\frac{2d_4}{\sqrt{3}}$	$\Xi_{cc}^{++} \rightarrow \Delta^+ D^+$	$-\frac{2d_4 \sin(\theta_C)}{\sqrt{3}}$
$\Xi_{cc}^+ \rightarrow \Sigma'^+ D^0$	$\frac{2d_5}{\sqrt{3}}$	$\Xi_{cc}^{++} \rightarrow \Sigma'^+ D_s^+$	$\frac{2d_4 \sin(\theta_C)}{\sqrt{3}}$
$\Xi_{cc}^+ \rightarrow \Sigma'^0 D^+$	$\sqrt{\frac{2}{3}}(d_4 + d_5)$	$\Xi_{cc}^+ \rightarrow \Delta^+ D^0$	$-\frac{2d_5 \sin(\theta_C)}{\sqrt{3}}$
$\Xi_{cc}^+ \rightarrow \Xi'^0 D_s^+$	$\frac{2d_5}{\sqrt{3}}$	$\Xi_{cc}^+ \rightarrow \Delta^0 D^+$	$-\frac{2(d_4 + d_5) \sin(\theta_C)}{\sqrt{3}}$
$\Omega_{cc}^+ \rightarrow \Xi'^0 D^+$	$\frac{2d_4}{\sqrt{3}}$	$\Xi_{cc}^+ \rightarrow \Sigma'^0 D_s^+$	$\sqrt{\frac{2}{3}}(d_4 - d_5) \sin(\theta_C)$
$\Xi_{cc}^{++} \rightarrow \Delta^+ D_s^+$	$\frac{2d_4 \sin^2(\theta_C)}{\sqrt{3}}$	$\Omega_{cc}^+ \rightarrow \Sigma'^+ D^0$	$\frac{2d_5 \sin(\theta_C)}{\sqrt{3}}$
$\Xi_{cc}^+ \rightarrow \Delta^0 D_s^+$	$\frac{2d_4 \sin^2(\theta_C)}{\sqrt{3}}$	$\Omega_{cc}^+ \rightarrow \Sigma'^0 D^+$	$\sqrt{\frac{2}{3}}(d_5 - d_4) \sin(\theta_C)$
$\Omega_{cc}^+ \rightarrow \Delta^+ D^0$	$\frac{2d_5 \sin^2(\theta_C)}{\sqrt{3}}$	$\Omega_{cc}^+ \rightarrow \Xi'^0 D_s^+$	$\frac{2(d_4 + d_5) \sin(\theta_C)}{\sqrt{3}}$
$\Omega_{cc}^+ \rightarrow \Delta^0 D^+$	$\frac{2d_5 \sin^2(\theta_C)}{\sqrt{3}}$		
$\Omega_{cc}^+ \rightarrow \Sigma'^0 D_s^+$	$\sqrt{\frac{2}{3}}(d_4 + d_5) \sin^2(\theta_C)$		

$$\Gamma(\Xi_{cc}^+ \rightarrow \Delta^0 D^+) = \Gamma(\Omega_{cc}^+ \rightarrow \Xi'^0 D_s^+), \quad (52)$$

$$\Gamma(\Xi_{cc}^+ \rightarrow \Sigma'^+ D^0) = \Gamma(\Xi_{cc}^+ \rightarrow \Xi'^0 D_s^+), \quad (53)$$

$$\Gamma(\Omega_{cc}^+ \rightarrow \Delta^+ D^0) = \Gamma(\Omega_{cc}^+ \rightarrow \Delta^0 D^+), \quad (54)$$

$$\Gamma(\Omega_{cc}^+ \rightarrow \Sigma'^0 D^+) = \Gamma(\Xi_{cc}^+ \rightarrow \Sigma'^0 D_s^+). \quad (55)$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Delta^+ D_s^+) = \Gamma(\Xi_{cc}^{++} \rightarrow \Delta^0 D_s^+),$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Sigma'^+ D^+) = \Gamma(\Omega_{cc}^{++} \rightarrow \Xi'^0 D^+),$$

$$\Gamma(\Xi_{cc}^{++} \rightarrow \Delta^+ D^0) = \Gamma(\Omega_{cc}^{++} \rightarrow \Sigma'^+ D^0), \quad (55)$$

In addition from the decay amplitudes, one can see that there are relations between the widths between Cabibbo-allowed, singly Cabibbo-suppressed and doubly Cabibbo-

Table 9 Relations for Cabibbo-allowed, singly Cabibbo-suppressed and doubly Cabibbo-suppressed processes in Tables 5 and 7

Channel 1	Channel 2	r	Channel 1	Channel 2	r
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+$	$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^+$	$-\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^+ K^0$	$\Omega_{cc}^+ \rightarrow \Lambda_c^+ \bar{K}^0$	-1
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$	$-\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+$	$\csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ \pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+$	-1	$\Xi_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$	$\sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 K^+$	$-\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+$	$\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	1
$\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^+ K^+$	$\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0 K^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 K^+$	$-\csc(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^0$	$\Omega_{cc}^+ \rightarrow \Xi_c^+ \bar{K}^0$	$\sin^2(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 \pi^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0 K^+$	$-\csc(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Lambda_c^+ \bar{K}^0$	$\Omega_{cc}^+ \rightarrow \Xi_c^+ K^0$	$\csc^2(\theta_c)$			
$\Xi_{cc}^{++} \rightarrow p D_s^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^+ D^+$	$-\sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow p D^0$	$\Xi_{cc}^+ \rightarrow \Sigma^+ D^0$	$\sin(\theta_c)$
$\Xi_{cc}^+ \rightarrow n D_s^+$	$\Omega_{cc}^+ \rightarrow \Xi^0 D^+$	$-\sin^2(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Sigma^+ D^0$	$\Xi_{cc}^+ \rightarrow \Sigma^+ D^0$	$\sin(\theta_c)$
$\Omega_{cc}^+ \rightarrow p D^0$	$\Xi_{cc}^+ \rightarrow \Sigma^+ D^0$	$-\sin^2(\theta_c)$	$\Xi_{cc}^{++} \rightarrow p D_s^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^+ D_s^+$	$-\sin(\theta_c)$
$\Omega_{cc}^+ \rightarrow n D^+$	$\Xi_{cc}^+ \rightarrow \Xi^0 D_s^+$	$-\sin^2(\theta_c)$	$\Xi_{cc}^{++} \rightarrow p D_s^+$	$\Xi_{cc}^{++} \rightarrow p D^+$	$-\sin(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma^+ D_s^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^+ D^+$	$\sin(\theta_c)$	$\Omega_{cc}^+ \rightarrow p D^0$	$\Xi_{cc}^+ \rightarrow p D^0$	$-\sin(\theta_c)$
$\Xi_{cc}^{++} \rightarrow p D^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^+ D^+$	$\sin(\theta_c)$	$\Omega_{cc}^+ \rightarrow p D^0$	$\Omega_{cc}^+ \rightarrow \Sigma^+ D^0$	$-\sin(\theta_c)$

Table 10 Relations for Cabibbo-allowed, singly Cabibbo-suppressed and doubly Cabibbo-suppressed processes in Table 6

Channel 1	Channel 2	r	Channel 1	Channel 2	r
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\pi^0$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\eta$	$-\frac{1}{\sqrt{3}}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\pi^0$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}K^0$	$\frac{\csc(\theta_c)}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	$\csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\pi^0$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\bar{K}^0$	$\frac{\sin(\theta_c)}{\sqrt{2}}$	$\Xi_{cc}^+ \rightarrow \Sigma_c^+K^0$	$\Omega_{cc}^+ \rightarrow \Xi_c^{'+}\bar{K}^0$	$\sin^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\eta$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}K^0$	$-\sqrt{\frac{3}{2}} \csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma_c^{'}\bar{K}^0$	$\Omega_{cc}^+ \rightarrow \Xi_c^{'}K^0$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\eta$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\bar{K}^0$	$-\sqrt{\frac{3}{2}} \sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^{'}K^0$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{'}\bar{K}^0$	-1
$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}K^0$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^{++}\bar{K}^0$	$\sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma_c^0\pi^+$	$\Xi_{cc}^+ \rightarrow \Xi_c^0\pi^+$	$-\sqrt{2} \sin(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+\pi^+$	$\Xi_{cc}^{++} \rightarrow \Sigma_c^+K^+$	$-\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma_c^0\pi^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0K^+$	$-\sqrt{2} \csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+\pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}\pi^+$	$-\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma_c^0\pi^+$	$\Omega_{cc}^+ \rightarrow \Omega_c^0K^+$	-1
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+\pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}K^+$	-1	$\Xi_{cc}^+ \rightarrow \Sigma_c^0K^+$	$\Omega_{cc}^+ \rightarrow \Omega_c^0\pi^+$	$\sin^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+K^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}\pi^+$	$\sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0\pi^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0K^+$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma_c^+K^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}K^+$	$\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0\pi^+$	$\Omega_{cc}^+ \rightarrow \Omega_c^0K^+$	$\frac{\csc(\theta_c)}{\sqrt{2}}$
$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}\pi^+$	$\Xi_{cc}^{++} \rightarrow \Xi_c^{'}K^+$	$\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi_c^0K^+$	$\Omega_{cc}^+ \rightarrow \Xi_c^0\pi^+$	-1
$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	$-\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Omega_c^0K^+$	$\Omega_{cc}^+ \rightarrow \Sigma_c^0\pi^+$	$\csc^2(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$-\csc(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	$\sin(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Omega_{cc}^+ \rightarrow \Sigma_c^{++}K^-$	-1	$\Omega_{cc}^+ \rightarrow \Xi_c^0K^+$	$\Omega_{cc}^+ \rightarrow \Omega_c^0K^+$	$\frac{\sin(\theta_c)}{\sqrt{2}}$

Table 11 Decay width relations for Cabibbo-allowed, singly Cabibbo-suppressed and doubly Cabibbo-suppressed processes in Table 8

Channel 1	Channel 2	r	Channel 1	Channel 2	r
$\Xi_{cc}^{++} \rightarrow \Delta^+D^+$	$\Xi_{cc}^{++} \rightarrow \Delta^+D_s^+$	$-\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Delta^0D^+$	$\Xi_{cc}^+ \rightarrow \Sigma^0D^+$	$-\sqrt{2} \sin(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D^+$	$-\frac{1}{2} \sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Delta^0D^+$	$\Omega_{cc}^+ \rightarrow \Sigma^0D_s^+$	$-\sqrt{2} \csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D_s^+$	$-\frac{1}{2}$	$\Xi_{cc}^+ \rightarrow \Delta^0D^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D_s^+$	-1
$\Xi_{cc}^{++} \rightarrow \Delta^+D^+$	$\Xi_{cc}^+ \rightarrow \Delta^0D_s^+$	$-\frac{1}{2} \csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Delta^0D_s^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D^+$	$\sin^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D^+$	$-\frac{1}{2} \sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\Xi_{cc}^+ \rightarrow \Xi^0D_s^+$	1
$\Xi_{cc}^{++} \rightarrow \Delta^+D_s^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D^+$	$\frac{1}{2} \sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\Omega_{cc}^+ \rightarrow \Delta^+D^0$	$2 \csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D_s^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D_s^+$	$\frac{\sin(\theta_c)}{2}$	$\Xi_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\Omega_{cc}^+ \rightarrow \Delta^0D^+$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D_s^+$	$\Xi_{cc}^+ \rightarrow \Delta^0D_s^+$	$\frac{1}{2}$	$\Xi_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\Omega_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\csc(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Delta^+D_s^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D^+$	$\frac{1}{2} \sin^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma^{'0}D^+$	$\Omega_{cc}^+ \rightarrow \Sigma^{'0}D_s^+$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D^+$	$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D_s^+$	$\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma^{'0}D^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D_s^+$	$\frac{\csc(\theta_c)}{\sqrt{2}}$
$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D^+$	$\Xi_{cc}^+ \rightarrow \Delta^0D_s^+$	$\csc^2(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Sigma^{'0}D_s^+$	$\Omega_{cc}^+ \rightarrow \Sigma^{'0}D^+$	-1
$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D^+$	1	$\Xi_{cc}^+ \rightarrow \Xi^0D_s^+$	$\Omega_{cc}^+ \rightarrow \Delta^+D^0$	$2 \csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D_s^+$	$\Xi_{cc}^+ \rightarrow \Delta^0D_s^+$	$\csc(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi^0D_s^+$	$\Omega_{cc}^+ \rightarrow \Delta^0D^+$	$\csc^2(\theta_c)$
$\Xi_{cc}^{++} \rightarrow \Sigma^{'+}D_s^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D^+$	$\sin(\theta_c)$	$\Xi_{cc}^+ \rightarrow \Xi^0D_s^+$	$\Omega_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\csc(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Delta^+D^0$	$\Xi_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$-\frac{1}{2} \sin(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Delta^+D^0$	$\Omega_{cc}^+ \rightarrow \Delta^0D^+$	$\frac{1}{2}$
$\Xi_{cc}^+ \rightarrow \Delta^+D^0$	$\Xi_{cc}^+ \rightarrow \Xi^0D_s^+$	$-\frac{1}{2} \sin(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Delta^+D^0$	$\Omega_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\frac{\sin(\theta_c)}{2}$
$\Xi_{cc}^+ \rightarrow \Delta^+D^0$	$\Omega_{cc}^+ \rightarrow \Delta^+D^0$	$-\csc(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Delta^0D^+$	$\Omega_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$\sin(\theta_c)$
$\Xi_{cc}^+ \rightarrow \Delta^+D^0$	$\Omega_{cc}^+ \rightarrow \Delta^0D^+$	$-\frac{1}{2} \csc(\theta_c)$	$\Omega_{cc}^+ \rightarrow \Sigma^{'0}D_s^+$	$\Omega_{cc}^+ \rightarrow \Xi^0D_s^+$	$\frac{\sin(\theta_c)}{\sqrt{2}}$
$\Xi_{cc}^+ \rightarrow \Delta^+D^0$	$\Omega_{cc}^+ \rightarrow \Sigma^{'+}D^0$	$-\frac{1}{2}$			

suppressed decay modes:

$$r^2 = \frac{\Gamma(\text{channel 1})}{\Gamma(\text{channel 2})}. \quad (56)$$

These relations are given in Tables 9, 10 and 11, respectively.

5 Nonleptonic Ξ_{bb} and Ω_{bb} decays

For the bottom quark decay, there are generically four kinds of quark-level transitions:

$$b \rightarrow c\bar{c}d/s, b \rightarrow c\bar{u}d/s, b \rightarrow u\bar{c}d/s, b \rightarrow q_1\bar{q}_2q_3, \quad (57)$$

with $q_{1,2,3}$ being the light quarks. Each of them will induce more than one types of decay modes at hadron level, which will be analyzed in order in the following.

5.1 $b \rightarrow c\bar{c}d/s$

5.1.1 Decays into J/ψ plus a bottom baryon

Such decays have the same topology as semileptonic $b \rightarrow s\ell^+\ell^-$ decays. The transition operator $b \rightarrow c\bar{c}d/s$ can form an SU(3) triplet, leading to the effective Hamiltonian:

$$H_{\text{eff}} = a_1(T_{bb})^i(H_3)^j(\bar{T}_{b\bar{3}})_{[ij]} J/\psi + a_2(T_{bb})^i(H_3)^j(\bar{T}_{b6})_{\{ij\}} J/\psi, \quad (58)$$

with $(H_3)_2 = V_{cd}^*$ and $(H_3)_3 = V_{cs}^*$. Feynman diagrams for these decays are given in Fig. 6.

The decay amplitudes are given in Table 12. Thus we have the following relations for decay widths:

$$\Gamma(\Xi_{bb}^0 \rightarrow \Lambda_b^0 J/\psi) = \Gamma(\Omega_{bb}^- \rightarrow \Xi_b^- J/\psi), \quad (59)$$

$$\Gamma(\Xi_{bb}^0 \rightarrow \Xi_b^0 J/\psi) = \Gamma(\Xi_{bb}^- \rightarrow \Xi_b^- J/\psi), \quad (60)$$

$$\begin{aligned} \Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^0 J/\psi) &= \Gamma(\Omega_{bb}^- \rightarrow \Xi_b'^- J/\psi) \\ &= \frac{1}{2}\Gamma(\Xi_{bb}^- \rightarrow \Sigma_b^- J/\psi), \end{aligned} \quad (61)$$




Fig. 6 The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into J/ψ and a bottom baryon

Table 12 Doubly bottom baryons decays into a J/ψ and a light baryon

Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 J/\psi$	$a_1 V_{cd}^*$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 J/\psi$	$a_1 V_{cs}^*$
$\Xi_{bb}^- \rightarrow \Xi_b^- J/\psi$	$a_1 V_{cs}^*$
$\Omega_{bb}^- \rightarrow \Xi_b^- J/\psi$	$-a_1 V_{cd}^*$
$\Xi_{bb}^0 \rightarrow \Sigma_b^0 J/\psi$	$\frac{a_2 V_{cd}^*}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow \Xi_b' J/\psi$	$\frac{a_2 V_{cs}^*}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Sigma_b^- J/\psi$	$a_2 V_{cd}^*$
$\Xi_{bb}^- \rightarrow \Xi_b'^- J/\psi$	$\frac{a_2 V_{cs}^*}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b'^- J/\psi$	$\frac{a_2 V_{cd}^*}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Omega_b^- J/\psi$	$a_2 V_{cs}^*$




Fig. 7 The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a doubly heavy baryon and an anticharmed meson

$$\begin{aligned} \Gamma(\Xi_{bb}^0 \rightarrow \Xi_b' J/\psi) &= \Gamma(\Xi_{bb}^- \rightarrow \Xi_b'^- J/\psi) \\ &= \frac{1}{2}\Gamma(\Omega_{bb}^- \rightarrow \Omega_b^- J/\psi). \end{aligned} \quad (62)$$

5.1.2 Decays into a doubly heavy baryon bcq plus a anticharmed meson

The $b \rightarrow c\bar{c}d/s$ transition can lead to another type of effective Hamiltonian:

$$H_{\text{eff}} = a_3(T_{bb})^i(H_3)^j(\bar{T}_{bc})_i D_j + a_4(T_{bb})^i(H_3)^j(\bar{T}_{bc})_j D_i, \quad (63)$$

which corresponds to the decays into doubly heavy baryon bcq plus a anticharmed meson. The Feynman diagrams for these decays are given in Fig. 7. The decay amplitudes are given in Table 13. Thus we obtain the following relations for the decay widths:

$$\begin{aligned} \Gamma(\Xi_{bb}^0 \rightarrow \Xi_{bc}^0 \bar{D}^0) &= \Gamma(\Omega_{bb}^- \rightarrow \Xi_{bc}^0 D_s^-), \\ \Gamma(\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 \bar{D}^0) &= \Gamma(\Xi_{bb}^- \rightarrow \Omega_{bc}^0 D^-), \\ \Gamma(\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ D^-) &= \Gamma(\Omega_{bb}^- \rightarrow \Omega_{bc}^0 D^-), \\ \Gamma(\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ D_s^-) &= \Gamma(\Xi_{bb}^- \rightarrow \Xi_{bc}^0 D_s^-). \end{aligned} \quad (64)$$

Table 13 Doubly bottom baryons decays into a bcq and an anticharmed meson

Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ D^-$	$a_3 V_{cd}^*$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ D_s^-$	$a_3 V_{cs}^*$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^0 \bar{D}^0$	$a_4 V_{cd}^*$
$\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 \bar{D}^0$	$a_4 V_{cs}^*$
$\Xi_{bb}^- \rightarrow \Xi_{bc}^0 D^-$	$(a_3 + a_4) V_{cd}^*$
$\Xi_{bb}^- \rightarrow \Xi_{bc}^0 D_s^-$	$a_3 V_{cs}^*$
$\Xi_{bb}^- \rightarrow \Omega_{bc}^0 D^-$	$a_4 V_{cs}^*$
$\Omega_{bb}^- \rightarrow \Xi_{bc}^0 D_s^-$	$a_4 V_{cd}^*$
$\Omega_{bb}^- \rightarrow \Omega_{bc}^0 D^-$	$a_3 V_{cd}^*$
$\Omega_{bb}^- \rightarrow \Omega_{bc}^0 D_s^-$	$(a_3 + a_4) V_{cs}^*$


5.2 $b \rightarrow c\bar{u}d/s$ transition

5.2.1 Decays into a doubly heavy baryon bcq plus a light meson

The operator to produce a charm quark from the b -quark decay, $\bar{c}b\bar{q}u$, is given by

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* [C_1 O_1^{\bar{c}u} + C_2 O_2^{\bar{c}u}] + \text{h.c..} \quad (65)$$

The light quarks in this effective Hamiltonian form an octet with the non-zero entry

**Fig. 8** The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a doubly heavy baryon and a light meson**Table 14** Doubly bottom baryons decays into a bcq and a light meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ \pi^-$	$(a_5 + a_7) V_{ud}^*$	$\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 \eta$	$\frac{(a_6 - 2a_7)V_{us}^*}{\sqrt{6}}$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^+ K^-$	$(a_5 + a_7) V_{us}^*$	$\Xi_{bb}^- \rightarrow \Xi_{bc}^0 \pi^-$	$(a_5 + a_6) V_{ud}^*$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^0 \pi^0$	$\frac{(a_6 - a_7)V_{ud}^*}{\sqrt{2}}$	$\Xi_{bb}^- \rightarrow \Xi_{bc}^0 K^-$	$a_5 V_{us}^*$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^0 \bar{K}^0$	$a_7 V_{us}^*$	$\Xi_{bb}^- \rightarrow \Omega_{bc}^0 \pi^-$	$a_6 V_{us}^*$
$\Xi_{bb}^0 \rightarrow \Xi_{bc}^0 \eta$	$\frac{(a_6 + a_7)V_{ud}^*}{\sqrt{6}}$	$\Omega_{bb}^- \rightarrow \Xi_{bc}^0 K^-$	$a_6 V_{ud}^*$
$\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 \pi^0$	$\frac{a_6 V_{us}^*}{\sqrt{2}}$	$\Omega_{bb}^- \rightarrow \Omega_{bc}^0 \pi^-$	$a_5 V_{ud}^*$
$\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 K^0$	$a_7 V_{ud}^*$	$\Omega_{bb}^- \rightarrow \Omega_{bc}^0 K^-$	$(a_5 + a_6) V_{us}^*$

$$(H_8)_1^2 = V_{ud}^*, \quad (66)$$

for the $b \rightarrow c\bar{u}d$ transition, and $(H_8)_1^3 = V_{us}^*$ for the $b \rightarrow c\bar{s}s$ transition. The hadron-level effective Hamiltonian is then given as

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & a_5 (T_{bb})^i (\bar{T}_{bc})_i M_j^k (H_8)_k^j \\ & + a_6 (T_{bb})^i (\bar{T}_{bc})_j M_i^k (H_8)_k^j \\ & + a_7 (T_{bb})^i (\bar{T}_{bc})_k M_j^k (H_8)_i^j. \end{aligned} \quad (67)$$

The Feynman diagrams for these decays are given in Fig. 8. The decay amplitudes are given in Table 14, which leads to:

$$\Gamma(\Xi_{bb}^0 \rightarrow \Omega_{bc}^0 \pi^0) = \frac{1}{2} \Gamma(\Xi_{bb}^- \rightarrow \Omega_{bc}^0 \pi^-). \quad (68)$$

5.2.2 Decays into a bottom baryon bqq plus a charmed meson

The effective Hamiltonian from the operator $\bar{c}b\bar{q}u$ gives

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & a_8 (T_{bb})^i (\bar{T}_{b\bar{3}})_{[ij]} \bar{D}^k (H_8)_k^j \\ & + a_9 (T_{bb})^i (\bar{T}_{b\bar{3}})_{[jk]} \bar{D}^k (H_8)_i^j \\ & + a_{10} (T_{bb})^i (\bar{T}_{b6})_{\{ij\}} \bar{D}^k (H_8)_k^j \\ & + a_{11} (T_{bb})^i (\bar{T}_{b6})_{\{jk\}} \bar{D}^k (H_8)_i^j. \end{aligned} \quad (69)$$

The Feynman diagrams for these decays are given in Fig. 9. Results are given in Table 15, thus we have the relations for







Fig. 9 The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a bottom baryon and a charmed meson

Table 15 Doubly bottom baryons decays into a bqq and a charmed meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 D^0$	$(a_8 - a_9) V_{ud}^*$	$\Xi_{bb}^0 \rightarrow \Sigma_b^0 D^0$	$\frac{(a_{10} + a_{11}) V_{ud}^*}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 D^0$	$(a_8 - a_9) V_{us}^*$	$\Xi_{bb}^0 \rightarrow \Sigma_b^- D^+$	$a_{11} V_{ud}^*$
$\Xi_{bb}^0 \rightarrow \Xi_b^- D^+$	$-a_9 V_{us}^*$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 D^0$	$\frac{(a_{10} + a_{11}) V_{us}^*}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow \Xi_b^- D_s^+$	$a_9 V_{ud}^*$	$\Xi_{bb}^0 \rightarrow \Xi_b'^- D^+$	$\frac{a_{11} V_{us}^*}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^- D^0$	$a_8 V_{us}^*$	$\Xi_{bb}^0 \rightarrow \Xi_b'^- D_s^+$	$\frac{a_{11} V_{ud}^*}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- D^0$	$-a_8 V_{ud}^*$	$\Xi_{bb}^0 \rightarrow \Omega_b^- D_s^+$	$a_{11} V_{us}^*$
		$\Xi_{bb}^- \rightarrow \Sigma_b^- D^0$	$a_{10} V_{ud}^*$
		$\Xi_{bb}^- \rightarrow \Xi_b'^- D^0$	$\frac{a_{10} V_{us}^*}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Xi_b'^- D^0$	$\frac{a_{10} V_{ud}^*}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Omega_b^- D^0$	$a_{10} V_{us}^*$

decay amplitudes:

$$\begin{aligned} \Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^- D^+) &= 2\Gamma(\Xi_{bb}^0 \rightarrow \Xi_b'^- D_s^+), \\ \Gamma(\Xi_{bb}^0 \rightarrow \Xi_b'^- D^+) &= \frac{1}{2}\Gamma(\Xi_{bb}^0 \rightarrow \Omega_b^- D_s^+). \\ \Gamma(\Xi_{bb}^- \rightarrow \Sigma_b^- D^0) &= 2\Gamma(\Omega_{bb}^- \rightarrow \Xi_b'^- D^0), \\ \Gamma(\Xi_{bb}^- \rightarrow \Xi_b'^- D^0) &= \frac{1}{2}\Gamma(\Omega_{bb}^- \rightarrow \Omega_b^- D^0). \end{aligned} \quad (70)$$

5.3 $b \rightarrow u\bar{c}d/s$: decays into a bottom baryon bqq plus an anticharmed meson.

For the anticharm production, the operator having the quark contents $(\bar{u}b)(\bar{q}c)$ is given by

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{ub} V_{cq}^* [C_1 O_1^{\bar{u}c} + C_2 O_2^{\bar{u}c}] + \text{h.c..} \quad (71)$$

The two light antiquarks form the $\bar{\mathbf{3}}$ and $\mathbf{6}$ representations. The antisymmetric tensor $H''_{\bar{\mathbf{3}}}$ and the symmetric tensor H_6 have non-zero components

$$(H''_{\bar{\mathbf{3}}})^{13} = -(H''_{\bar{\mathbf{3}}})^{31} = V_{cs}^*, \quad (H_6)^{13} = (H_6)^{31} = V_{cs}^*, \quad (72)$$

Fig. 10 The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a bottom baryon and an anticharmed meson

for the $b \rightarrow u\bar{c}s$ transition. For the transition $b \rightarrow u\bar{c}d$ one requires the interchange of $2 \leftrightarrow 3$ in the subscripts, and V_{cs} replaced by V_{cd} .

The effective Hamiltonian is constructed as

$$\begin{aligned} \mathcal{H}_{\text{eff}} = & b_1(T_{bb})^i (\bar{T}_{b\bar{3}})_{\{ij\}} D_k (H''_{\bar{\mathbf{3}}})^{jk} \\ & + b_2(T_{bb})^k (\bar{T}_{b\bar{3}})_{\{ij\}} D_k (H''_{\bar{\mathbf{3}}})^{ij} \\ & + b_3(T_{bb})^i (\bar{T}_{b\bar{3}})_{\{ij\}} D_k (H''_6)^{jk} \\ & + b_4(T_{bb})^i (\bar{T}_{b6})_{\{ij\}} D_k (H''_6)^{jk} \\ & + b_5(T_{bb})^k (\bar{T}_{b6})_{\{ij\}} D_k (H''_6)^{ij} \\ & + b_6(T_{bb})^i (\bar{T}_{b6})_{\{ij\}} D_k (H''_{\bar{\mathbf{3}}})^{jk}. \end{aligned} \quad (73)$$

The Feynman diagrams for these decays are given in Fig. 10. The decay amplitudes for different channels are given in Table 16.

Thus we have the relations for decay amplitudes:

$$\begin{aligned} \Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^+ D^-) &= 2\Gamma(\Omega_{bb}^- \rightarrow \Xi_b^0 D^-), \\ \Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^+ D_s^-) &= 2\Gamma(\Xi_{bb}^- \rightarrow \Sigma_b^0 D_s^-), \\ \Gamma(\Xi_{bb}^- \rightarrow \Sigma_b^- \bar{D}^0) &= 2\Gamma(\Omega_{bb}^- \rightarrow \Xi_b'^- \bar{D}^0), \\ \Gamma(\Xi_{bb}^- \rightarrow \Xi_b'^- \bar{D}^0) &= \frac{1}{2}\Gamma(\Omega_{bb}^- \rightarrow \Omega_b^- \bar{D}^0). \end{aligned}$$

As one can see, the Ξ_{bb} can decay into both $\Xi_b D^0$ and $\Xi_b \bar{D}^0$. The D^0 and \bar{D}^0 can form the CP eigenstate D_+ and D_- . Thus using the Ξ_{bb} decays into the $\Xi_b D_{\pm}$, one may construct the interference between the $b \rightarrow c\bar{s}u$ and $b \rightarrow u\bar{c}s$. The CKM angle γ can then be extracted from measuring decay widths of these channels, as in the case of $B \rightarrow DK$ [73–78], $B \rightarrow DK_{0,2}^*$ [79, 80] and others. This is also similar for the $\Omega_{bb} \rightarrow \Omega^- D_{\pm}$ decays and the following $\Xi_{bc} \rightarrow \Xi_c D_{\pm}$ and $\Omega_{bc} \rightarrow \Omega^0 D_{\pm}$ modes.

5.4 Charmless $b \rightarrow q_1 \bar{q}_2 q_3$ decays

5.4.1 Decays into a bottom baryon and a light meson

The charmless $b \rightarrow q$ ($q = d, s$) transition is controlled by the weak Hamiltonian \mathcal{H}_{eff} :

Table 16 Doubly bottom baryons decays into a $b\bar{q}q$ and an anticharmed meson

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 \bar{D}^0$	$(-b_1 + 2b_2 + b_3) V_{cd}^*$	$\Xi_{bb}^0 \rightarrow \Sigma_b^+ D^-$	$(b_4 + b_6) V_{cd}^*$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 \bar{D}^0$	$(-b_1 + 2b_2 + b_3) V_{cs}^*$	$\Xi_{bb}^0 \rightarrow \Sigma_b^+ D_s^-$	$(b_4 + b_6) V_{cs}^*$
$\Xi_{bb}^- \rightarrow \Lambda_b^0 D^-$	$-(b_1 - 2b_2 + b_3) V_{cd}^*$	$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \bar{D}^0$	$\frac{(b_4 + 2b_5 - b_6)V_{cd}^*}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Lambda_b^0 D_s^-$	$-(b_1 + b_3) V_{cs}^*$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 \bar{D}^0$	$\frac{(b_4 + 2b_5 - b_6)V_{cs}^*}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^0 D^-$	$2b_2 V_{cs}^*$	$\Xi_{bb}^- \rightarrow \Sigma_b^0 D^-$	$\frac{(b_4 + 2b_5 + b_6)V_{cd}^*}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^- \bar{D}^0$	$(b_3 - b_1) V_{cs}^*$	$\Xi_{bb}^- \rightarrow \Sigma_b^0 D_s^-$	$\frac{(b_4 + b_6)V_{cs}^*}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Lambda_b^0 D_s^-$	$2b_2 V_{cd}^*$	$\Xi_{bb}^- \rightarrow \Sigma_b^- \bar{D}^0$	$(b_4 - b_6) V_{cd}^*$
$\Omega_{bb}^- \rightarrow \Xi_b^0 D^-$	$-(b_1 + b_3) V_{cd}^*$	$\Xi_{bb}^- \rightarrow \Xi_b^0 D^-$	$\sqrt{2}b_5 V_{cs}^*$
$\Omega_{bb}^- \rightarrow \Xi_b^0 D_s^-$	$-(b_1 - 2b_2 + b_3) V_{cs}^*$	$\Xi_{bb}^- \rightarrow \Xi_b'^- \bar{D}^0$	$\frac{(b_4 - b_6)V_{cs}^*}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- \bar{D}^0$	$(b_1 - b_3) V_{cd}^*$	$\Omega_{bb}^- \rightarrow \Sigma_b^0 D_s^-$	$\sqrt{2}b_5 V_{cd}^*$
		$\Omega_{bb}^- \rightarrow \Xi_b^0 D^-$	$\frac{(b_4 + b_6)V_{cd}^*}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Xi_b^0 D_s^-$	$\frac{(b_4 + 2b_5 + b_6)V_{cs}^*}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Xi_b'^- \bar{D}^0$	$\frac{(b_4 - b_6)V_{cd}^*}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Omega_b^- \bar{D}^0$	$(b_4 - b_6) V_{cs}^*$

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left\{ V_{ub} V_{uq}^* [C_1 O_1^{\bar{u}u} + C_2 O_2^{\bar{u}u}] - V_{tb} V_{tq}^* \left[\sum_{i=3}^{10} C_i O_i \right] \right\} + \text{h.c.}, \quad (74)$$

where O_i is a four-quark operator or a moment type operator. At the hadron level, penguin operators behave as the $\mathbf{3}$ representation while tree operators can be decomposed in terms of a vector H_3 , a traceless tensor antisymmetric in upper indices, H_6 , and a traceless tensor symmetric in upper indices, H_{15} . For the $\Delta S = 0(b \rightarrow d)$ decays, the non-zero components of the effective Hamiltonian are [37, 45, 58]

$$\begin{aligned} (H_3)^2 &= 1, \quad (H_6)_1^{12} = -(H_6)_1^{21} \\ &= (H_6)_3^{23} = -(H_6)_3^{32} = 1, \\ 2(H_{15})_1^{12} &= 2(H_{15})_1^{21} = -3(H_{15})_2^{22} \\ &= -6(H_{15})_3^{23} = -6(H_{15})_3^{32} = 6, \end{aligned} \quad (75)$$

and all other remaining entries are zero. For the $\Delta S = 1(b \rightarrow s)$ decays the non-zero entries in the H_3 , H_6 , H_{15} are obtained from Eq. (75) with the exchange $2 \leftrightarrow 3$.

The effective hadron-level Hamiltonian for decays into the bottom antitriplet is constructed as

$$\begin{aligned} \mathcal{H}_{\text{eff}} = c_1(T_{bb})^i (\bar{T}_{b\bar{3}})_{[ij]} M_l^j (H_3)^l \\ + c_2(T_{bb})^i (\bar{T}_{b\bar{3}})_{[jl]} M_i^j (H_3)^l \\ + c_3(T_{bb})^i (\bar{T}_{b\bar{3}})_{[ij]} M_l^k (H_6)_k^{jl} \\ + c_4(T_{bb})^i (\bar{T}_{b\bar{3}})_{[jl]} M_i^k (H_6)_k^{jl} \\ + c_5(T_{bb})^i (\bar{T}_{b\bar{3}})_{[jk]} M_l^k (H_6)_i^{jl} \\ + c_6(T_{bb})^i (\bar{T}_{b\bar{3}})_{[ij]} M_l^k (H_{15})_k^{jl} \end{aligned}$$

$$+ c_7(T_{bb})^i (\bar{T}_{b\bar{3}})_{[jk]} M_l^k (H_{15})_i^{jl}, \quad (76)$$

while, for the sextet baryon, we have

$$\begin{aligned} \mathcal{H}_{\text{eff}} = c_8(T_{bb})^i (\bar{T}_{b6})_{\{ij\}} M_l^j (H_3)^l \\ + c_9(T_{bb})^i (\bar{T}_{b6})_{\{jl\}} M_i^j (H_3)^l \\ + c_{10}(T_{bb})^i (\bar{T}_{b6})_{\{ij\}} M_l^k (H_{15})_k^{jl} \\ + c_{11}(T_{bb})^i (\bar{T}_{b6})_{\{jl\}} M_i^k (H_{15})_k^{jl} \\ + c_{12}(T_{bb})^i (\bar{T}_{b6})_{\{jk\}} M_l^k (H_{15})_i^{jl} \\ + c_{13}(T_{bb})^i (\bar{T}_{b6})_{\{ij\}} M_l^k (H_6)_k^{jl} \\ + c_{14}(T_{bb})^i (\bar{T}_{b6})_{\{jk\}} M_l^k (H_6)_i^{jl}. \end{aligned} \quad (77)$$

The Feynman diagrams for these decays are given in Fig. 11. The decay amplitudes for different channels are given in Tables 17 and 18 for the $b \rightarrow d$ transition and the $b \rightarrow s$ transition, respectively. Thus, it leads to the relations for the decay widths,

$$\Gamma(\Xi_{bb}^0 \rightarrow \Sigma_b^- \pi^+) = 2\Gamma(\Xi_{bb}^0 \rightarrow \Xi_b'^- K^+), \quad (78)$$

$$\Gamma(\Xi_{bb}^0 \rightarrow \Xi_b'^- \pi^+) = \frac{1}{2}\Gamma(\Xi_{bb}^0 \rightarrow \Omega_b^- K^+). \quad (79)$$

5.4.2 Decays into a bottom meson and a light baryon octet

The effective Hamiltonian is given as

$$\begin{aligned} \mathcal{H}_{\text{eff}} = d_1(T_{bb})^i \bar{B}^j \epsilon_{ijk} (T_8)_l^k (H_3)^l \\ + d_2(T_{bb})^i \bar{B}'^l \epsilon_{ijk} (T_8)_l^k (H_3)^j \\ + d_3(T_{bb})^l \bar{B}'^j \epsilon_{ijk} (T_8)_l^k (H_3)^i \end{aligned}$$




Fig. 11 The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a bottom baryon and a light meson

Table 17 Doubly bottom baryons decays into a bqq and a light meson induced by the charmless $b \rightarrow d$ transition

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 \pi^0$	$-\frac{c_1 - c_2 + c_3 - 2c_4 - 5c_6 + 6c_7}{\sqrt{2}}$	$\Xi_{bb}^0 \rightarrow \Sigma_b^+ \pi^-$	$c_8 + 3c_{10} + 3c_{12} + c_{13} + c_{14}$
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 \eta$	$\frac{c_1 + c_2 - 3c_3 + 2c_4 + 2c_5 + 3c_6}{\sqrt{6}}$	$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \pi^0$	$\frac{1}{2}(-c_8 + c_9 + 5c_{10} + 6c_{11} - c_{13} - 2c_{14})$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 K^0$	$c_1 - c_3 + c_5 - c_6 + 3c_7$	$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \eta$	$\frac{c_8 + c_9 + 3c_{10} + 6c_{11} + 6c_{12} - 3c_{13}}{2\sqrt{3}}$
$\Xi_{bb}^0 \rightarrow \Xi_b^- K^+$	$-c_2 + 2c_4 - c_5 + 3c_7$	$\Xi_{bb}^0 \rightarrow \Sigma_b^- \pi^+$	$c_9 - 2c_{11} + 3c_{12} - c_{14}$
$\Xi_{bb}^- \rightarrow \Lambda_b^0 \pi^-$	$-c_1 + c_2 - c_3 + 2c_4 - 3c_6 + 2c_7$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 K^0$	$\frac{c_8 - c_{10} + 3c_{12} - c_{13} + c_{14}}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^- K^0$	$c_1 - c_2 - c_3 + 2c_4 - c_6 - 2c_7$	$\Xi_{bb}^0 \rightarrow \Xi_b^- K^+$	$\frac{c_9 - 2c_{11} + 3c_{12} - c_{14}}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Lambda_b^0 K^-$	$c_2 + 2c_4 - c_5 + c_7$	$\Xi_{bb}^- \rightarrow \Sigma_b^0 \pi^-$	$\frac{c_8 + c_9 + 3c_{10} + 6c_{11} - 2c_{12} + c_{13}}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- \pi^-$	$-c_1 - c_3 + c_5 - 3c_6 + c_7$	$\Xi_{bb}^- \rightarrow \Sigma_b^- \pi^0$	$-\frac{c_8 + c_9 - 5c_{10} - 2c_{11} - 2c_{12} + c_{13}}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- \pi^0$	$\frac{c_1 + c_3 - c_5 - 5c_6 - c_7}{\sqrt{2}}$	$\Xi_{bb}^- \rightarrow \Sigma_b^- \eta$	$\frac{c_8 + c_9 + 3c_{10} - 2c_{11} - 2c_{12} - 3c_{13}}{\sqrt{6}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- \eta$	$-\frac{c_1 - 2c_2 - 3c_3 + 4c_4 + c_5 + 3c_6 - 3c_7}{\sqrt{6}}$	$\Xi_{bb}^- \rightarrow \Xi_b^- K^0$	$\frac{c_8 + c_9 - c_{10} - 2c_{11} - 2c_{12} - c_{13}}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Sigma_b^0 K^-$	$\frac{c_9 + 6c_{11} - c_{12} + c_{14}}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Sigma_b^- \bar{K}^0$	$c_9 - 2c_{11} - c_{12} + c_{14}$
		$\Omega_{bb}^- \rightarrow \Xi_b^0 \pi^-$	$\frac{c_8 + 3c_{10} - c_{12} + c_{13} - c_{14}}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Xi_b^- \pi^0$	$\frac{1}{2}(-c_8 + 5c_{10} + c_{12} - c_{13} + c_{14})$
		$\Omega_{bb}^- \rightarrow \Xi_b^- \eta$	$\frac{c_8 - 2c_9 + 3c_{10} + 4c_{11} + c_{12} - 3c_{13} - 3c_{14}}{2\sqrt{3}}$
		$\Omega_{bb}^- \rightarrow \Omega_b^- K^0$	$c_8 - c_{10} - c_{12} - c_{13} - c_{14}$


$$\begin{aligned}
 & + d_4 (T_{bb})^l \bar{B}^n \epsilon_{ijk} (T_8)_l^k (H_6)_n^{ij} \\
 & + d_5 (T_{bb})^l \bar{B}^n \epsilon_{ijk} (T_8)_n^k (H_6)_l^{ij} \\
 & + d_6 (T_{bb})^l \bar{B}^i \epsilon_{ijk} (T_8)_n^k (H_6)_l^{jn} \\
 & + d_7 (T_{bb})^i \bar{B}^l \epsilon_{ijk} (T_8)_n^k (H_6)_l^{jn} \\
 & + d_8 (T_{bb})^l \bar{B}^i \epsilon_{ijk} (T_8)_n^k (H_{15})_l^{jn} \\
 & + d_9 (T_{bb})^i \bar{B}^l \epsilon_{ijk} (T_8)_n^k (H_{15})_l^{jn}.
 \end{aligned} \tag{80}$$

Similarly, we find the reduced matrix elements d_4, d_7 and d_5, d_6 are not independent. So we use the following effective Hamiltonian:

$$\begin{aligned}
 \mathcal{H}_{\text{eff}} = & d_1 (T_{bb})^i \bar{B}^j \epsilon_{ijk} (T_8)_l^k (H_3)^l \\
 & + d_2 (T_{bb})^i \bar{B}^l \epsilon_{ijk} (T_8)_l^k (H_3)^j \\
 & + d_3 (T_{bb})^l \bar{B}^j \epsilon_{ijk} (T_8)_l^k (H_3)^i
 \end{aligned}$$

Table 18 Doubly bottom baryons decays into a bqq and a light meson induced by the charmless $b \rightarrow s$ transition

Channel	Amplitude	Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 \bar{K}^0$	$c_1 - c_3 + c_5 - c_6 + 3c_7$	$\Xi_{bb}^0 \rightarrow \Sigma_b^+ K^-$	$c_8 + 3c_{10} + 3c_{12} + c_{13} + c_{14}$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 \pi^0$	$\frac{c_2 - 2c_3 + 2c_4 + c_5 + 4c_6 - 3c_7}{\sqrt{2}}$	$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \bar{K}^0$	$\frac{c_8 - c_{10} + 3c_{12} - c_{13} + c_{14}}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow \Xi_b^0 \eta$	$\frac{-2c_1 + c_2 + 2c_4 - c_5 + 6c_6 - 9c_7}{\sqrt{6}}$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 \pi^0$	$\frac{1}{2} (c_9 + 4c_{10} + 6c_{11} + 3c_{12} - 2c_{13} - c_{14})$
$\Xi_{bb}^0 \rightarrow \Xi_b^- \pi^+$	$c_2 - 2c_4 + c_5 - 3c_7$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 \eta$	$\frac{-2c_8 + c_9 + 6c_{10} + 6c_{11} - 3c_{12} - 3c_{14}}{2\sqrt{3}}$
$\Xi_{bb}^- \rightarrow \Lambda_b^0 K^-$	$-c_1 - c_3 + c_5 - 3c_6 + c_7$	$\Xi_{bb}^0 \rightarrow \Xi_b^- \pi^+$	$\frac{c_9 - 2c_{11} + 3c_{12} - c_{14}}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^0 \pi^-$	$c_2 + 2c_4 - c_5 + c_7$	$\Xi_{bb}^0 \rightarrow \Omega_b^- K^+$	$c_9 - 2c_{11} + 3c_{12} - c_{14}$
$\Xi_{bb}^- \rightarrow \Xi_b^- \pi^0$	$\frac{-c_2 + 2c_3 - 2c_4 - c_5 - 4c_6 + c_7}{\sqrt{2}}$	$\Xi_{bb}^- \rightarrow \Sigma_b^0 K^-$	$\frac{c_8 + 3c_{10} - c_{12} + c_{13} - c_{14}}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi_b^- \eta$	$\frac{-2c_1 + c_2 - 2c_4 + c_5 + 6c_6 + 3c_7}{\sqrt{6}}$	$\Xi_{bb}^- \rightarrow \Sigma_b^- \bar{K}^0$	$c_8 - c_{10} - c_{12} - c_{13} - c_{14}$
$\Omega_{bb}^- \rightarrow \Xi_b^0 K^-$	$-c_1 + c_2 - c_3 + 2c_4 - 3c_6 + 2c_7$	$\Xi_{bb}^- \rightarrow \Xi_b^0 \pi^-$	$\frac{c_9 + 6c_{11} - c_{12} + c_{14}}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi_b^- \bar{K}^0$	$-c_1 + c_2 + c_3 - 2c_4 + c_6 + 2c_7$	$\Xi_{bb}^- \rightarrow \Xi_b' \pi^0$	$\frac{1}{2} (-c_9 + 4c_{10} + 2c_{11} + c_{12} - 2c_{13} - c_{14})$
		$\Xi_{bb}^- \rightarrow \Xi_b' \eta$	$\frac{-2c_8 + c_9 + 6c_{10} - 2c_{11} + c_{12} + 3c_{14}}{2\sqrt{3}}$
		$\Xi_{bb}^- \rightarrow \Omega_b^- K^0$	$c_9 - 2c_{11} - c_{12} + c_{14}$
		$\Omega_{bb}^- \rightarrow \Xi_b^0 K^-$	$\frac{c_8 + c_9 + 3c_{10} + 6c_{11} - 2c_{12} + c_{13}}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Xi_b^- \bar{K}^0$	$\frac{c_8 + c_9 - c_{10} - 2c_{11} - 2c_{12} - c_{13}}{\sqrt{2}}$
		$\Omega_{bb}^- \rightarrow \Omega_b^- \pi^0$	$\sqrt{2} (2c_{10} - c_{13})$
		$\Omega_{bb}^- \rightarrow \Omega_b^- \eta$	$-\sqrt{\frac{2}{3}} (c_8 + c_9 - 3c_{10} - 2c_{11} - 2c_{12})$

**Fig. 12** The Feynman diagrams for Ξ_{bb} and Ω_{bb} decays into a bottom meson and a light baryon

$$\begin{aligned}
& + d_6(T_{bb})^l \bar{B}^i \epsilon_{ijk} (T_8)_n^k (H_6)_l^{jn} \\
& + d_7(T_{bb})^i \bar{B}^l \epsilon_{ijk} (T_8)_n^k (H_6)_l^{jn} \\
& + d_8(T_{bb})^l \bar{B}^i \epsilon_{ijk} (T_8)_n^k (H_{15})_l^{jn} \\
& + d_9(T_{bb})^i \bar{B}^l \epsilon_{ijk} (T_8)_n^k (H_{15})_l^{jn}.
\end{aligned} \quad (81)$$

The Feynman diagrams for these decays are given in Fig. 12. The decay amplitudes for different channels are given in Tables 19 and 20 for the $b \rightarrow d$ transition and the $b \rightarrow s$ transition, respectively.

5.4.3 Decays into a bottom meson and a light baryon decuplet

The effective Hamiltonian is given as

$$\mathcal{H}_{\text{eff}} = f_3(T_{bb})^l \bar{B}^j (T_{10})_{ijl} (H_3)^i$$

$$\begin{aligned}
& + f_4(T_{bb})^l \bar{B}^n (T_{10})_{ijl} (H_{15})_n^{ij} \\
& + f_5(T_{bb})^l \bar{B}^n (T_{10})_{ijn} (H_{15})_l^{ij}.
\end{aligned} \quad (82)$$

The Feynman diagrams for these decays are the same as in Fig. 12. The decay amplitudes for different channels are given in Tables 19 and 20 for the $b \rightarrow d$ transition and the $b \rightarrow s$ transition, respectively.

We summarize the relations for decay widths for Ξ_{bb} and Ω_{bb} decay into a bottom meson and a light baryon,

$$\Gamma(\Xi_{bb}^0 \rightarrow \Delta^0 \bar{B}^0) = 2\Gamma(\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}_s^0), \quad (83)$$

$$\begin{aligned}
& \Gamma(\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0) \\
& = 3\Gamma(\Xi_{bb}^- \rightarrow \Sigma^- \bar{B}_s^0) = 3\Gamma(\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0) \\
& = 3\Gamma(\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0),
\end{aligned} \quad (84)$$

Table 19 Doubly bottom baryons decays into a bottom meson and a light baryon induced by the charmless $b \rightarrow d$ transition

Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda^0 \bar{B}_s^0$	$\frac{-d_1+2d_2-d_3-2d_6+d_7-3d_9}{\sqrt{6}}$
$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}_s^0$	$\frac{d_1+d_3-d_7-6d_8-d_9}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow p \bar{B}^-$	$d_2 - d_3 - d_6 - d_7 + 3d_8 + 3d_9$
$\Xi_{bb}^0 \rightarrow n \bar{B}^0$	$d_1 + d_2 - d_6 - 3d_8 - 2d_9$
$\Xi_{bb}^- \rightarrow \Sigma^- \bar{B}_s^0$	$d_1 + d_3 - d_7 + 2d_8 - d_9$
$\Xi_{bb}^- \rightarrow n \bar{B}^-$	$-d_1 - d_3 - d_7 - 2d_8 - 3d_9$
$\Omega_{bb}^- \rightarrow \Lambda^0 \bar{B}^-$	$\frac{d_1-d_2+2d_3-d_6+2d_7+3d_8}{\sqrt{6}}$
$\Omega_{bb}^- \rightarrow \Sigma^- \bar{B}^0$	$-d_1 - d_2 - d_6 - d_8 + 2d_9$
$\Omega_{bb}^- \rightarrow \Sigma^0 \bar{B}^-$	$\frac{-d_1+d_2+d_6+d_8+6d_9}{\sqrt{2}}$
$\Omega_{bb}^- \rightarrow \Xi^- \bar{B}_s^0$	$-d_2 + d_3 - d_6 - d_7 + d_8 + d_9$
$\Xi_{bb}^0 \rightarrow \Delta^+ \bar{B}^-$	$\frac{f_3+6(f_4+f_5)}{\sqrt{3}}$
$\Xi_{bb}^0 \rightarrow \Delta^0 \bar{B}^0$	$\frac{f_3-2f_4+6f_5}{\sqrt{3}}$
$\Xi_{bb}^0 \rightarrow \Sigma'^0 \bar{B}_s^0$	$\frac{f_3-2f_4+6f_5}{\sqrt{6}}$
$\Xi_{bb}^- \rightarrow \Delta^0 \bar{B}^-$	$\frac{f_3+6f_4-2f_5}{\sqrt{3}}$
$\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0$	$f_3 - 2(f_4 + f_5)$
$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Sigma'^0 \bar{B}^-$	$\frac{f_3+6f_4-2f_5}{\sqrt{6}}$
$\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$

$$\Gamma(\Xi_{bb}^- \rightarrow \Delta^0 \bar{B}^-) = 2\Gamma(\Omega_{bb}^- \rightarrow \Sigma'^0 \bar{B}^-). \\ \Gamma(\Xi_{bb}^- \rightarrow \Delta^0 \bar{B}^-) = 2\Gamma(\Omega_{bb}^- \rightarrow \Sigma'^0 \bar{B}^-).$$

$$\Gamma(\Xi_{bb}^0 \rightarrow \Sigma'^0 \bar{B}^0) = \frac{1}{2}\Gamma(\Xi_{bb}^0 \rightarrow \Xi'^0 \bar{B}_s^0), \quad (85)$$

$$\Gamma(\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}^0) = \Gamma(\Xi_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0) \\ = \Gamma(\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}^0) = \frac{1}{3}\Gamma(\Omega_{bb}^- \rightarrow \Omega^- \bar{B}_s^0). \\ \Gamma(\Xi_{bb}^- \rightarrow \Xi'^0 \bar{B}^-) = \frac{1}{2}\Gamma(\Omega^- \bar{B}_s^0 \rightarrow \Xi'^0 \bar{B}^-). \quad (86)$$

5.4.4 U-spin for Ξ_{bb} and Ω_{bb} decays

For Ξ_{bb} , Ω_{bb} decays induced by the $b \rightarrow q_1 q_2 q_3$, there are two amplitudes with different CKM factors. We consider the connected decays with the decay amplitudes

$$A(\Delta S = 0) = r(V_{ub} V_{ud}^* A_{\Xi_{bb}, \Omega_{bb}}^T + V_{tb} V_{td}^* A_{\Xi_{bb}, \Omega_{bb}}^P), \\ A(\Delta S = 1) = V_{ub} V_{us}^* A_{\Xi_{bb}, \Omega_{bb}}^T + V_{tb} V_{ts}^* A_{\Xi_{bb}, \Omega_{bb}}^P. \quad (87)$$

As pointed out in Refs. [40, 43, 44], there exists a relation for the CP violating quantity $\Delta = \Gamma - \bar{\Gamma}$. The relation about

Table 20 Doubly bottom baryons decays into a bottom meson and a light baryon induced by the charmless $b \rightarrow s$ transition

Channel	Amplitude
$\Xi_{bb}^0 \rightarrow \Lambda^0 \bar{B}^0$	$\frac{-2d_1-d_2-d_3+d_6+d_7+9d_8+3d_9}{\sqrt{6}}$
$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}^0$	$\frac{d_2-d_3-d_6+d_7+3d_8-d_9}{\sqrt{2}}$
$\Xi_{bb}^0 \rightarrow \Sigma^+ \bar{B}^-$	$-d_2 + d_3 + d_6 + d_7 - 3d_8 - 3d_9$
$\Xi_{bb}^0 \rightarrow \Xi^- \bar{B}_s^0$	$-d_1 - d_2 + d_6 + 3d_8 + 2d_9$
$\Xi_{bb}^- \rightarrow \Lambda^0 \bar{B}^-$	$\frac{2d_1+d_2+d_3+d_6+d_7+3d_8+9d_9}{\sqrt{6}}$
$\Xi_{bb}^- \rightarrow \Sigma^+ \bar{B}^0$	$d_2 - d_3 + d_6 + d_7 - d_8 - d_9$
$\Xi_{bb}^- \rightarrow \Sigma^0 \bar{B}^-$	$\frac{d_2-d_3+d_6-d_7-d_8+3d_9}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Xi^- \bar{B}_s^0$	$d_1 + d_2 + d_6 + d_8 - 2d_9$
$\Omega_{bb}^- \rightarrow \Xi^0 \bar{B}^-$	$-d_1 - d_3 + d_7 - 2d_8 + d_9$
$\Omega_{bb}^- \rightarrow \Sigma^0 \bar{B}^-$	$d_1 + d_3 + d_7 + 2d_8 + 3d_9$
$\Xi_{bb}^0 \rightarrow \Sigma'^+ \bar{B}^-$	$\frac{f_3+6(f_4+f_5)}{\sqrt{3}}$
$\Xi_{bb}^0 \rightarrow \Sigma'^0 \bar{B}^0$	$\frac{f_3-2f_4+6f_5}{\sqrt{6}}$
$\Xi_{bb}^0 \rightarrow \Xi'^0 \bar{B}_s^0$	$\frac{f_3-2f_4+6f_5}{\sqrt{3}}$
$\Xi_{bb}^- \rightarrow \Sigma'^0 \bar{B}^-$	$\frac{f_3+6f_4-2f_5}{\sqrt{6}}$
$\Xi_{bb}^- \rightarrow \Sigma'-' \bar{B}^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$
$\Xi_{bb}^- \rightarrow \Xi'-' \bar{B}_s^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Xi'^0 \bar{B}^-$	$\frac{f_3+6f_4-2f_5}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Xi'-' \bar{B}^0$	$\frac{f_3-2(f_4+f_5)}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Omega \bar{B}_s^0$	$f_3 - 2(f_4 + f_5)$

decay widths $\Gamma(\Delta S = i)$ and CP asymmetry $A_{CP}(\Delta S = i)$ is

$$\frac{A_{CP}(\Delta S = 0)}{A_{CP}(\Delta S = 1)} = -r^2 \frac{\Gamma(\Delta S = 1)}{\Gamma(\Delta S = 0)}. \quad (88)$$

In Tables 21 and 22, we collect the Ξ_{bb} , Ω_{bb} decay pairs related by U-spin. The CP asymmetries and decay widths for these pairs satisfy relation in Eq. (88). The experimental data in the future is important to test flavor SU(3) symmetry and also the CKM mechanism for CP violation.

6 Nonleptonic Ξ_{bc} and Ω_{bc} decays

The decays of Ξ_{bc} and Ω_{bc} can proceed via the b quark decay or the c quark decay, which are induced by the following quark transitions:

$$c \rightarrow s \bar{d} u, \quad c \rightarrow u \bar{d} d / \bar{s} s, \quad c \rightarrow d \bar{s} u, \\ b \rightarrow \bar{c} c d / s, \quad b \rightarrow c \bar{u} d / s, \quad b \rightarrow u \bar{c} d / s, \quad b \rightarrow q_1 \bar{q}_2 q_3. \quad (89)$$

As we have shown in the semileptonic case, for the charm quark decays, one can obtain the decay amplitudes from those

Table 21 U-spin relations for Ξ_{bb} , Ω_{bb} decays into a bottom baryon and a light meson. Results in the “channel 1” are for $b \rightarrow d$ processes and the ones in the “channel 2” are for $b \rightarrow s$ processes. r denotes the ratio of the two amplitudes

Channel 1	Channel 2	r	Channel 1	Channel 2	r
$\Xi_{bb}^0 \rightarrow \Lambda_b^0 \bar{K}^0$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 K^0$	1	$\Xi_{bb}^0 \rightarrow \Xi_b^- \pi^+$	$\Xi_{bb}^0 \rightarrow \Xi_b'^- K^+$	1
$\Xi_{bb}^0 \rightarrow \Xi_b^- \pi^+$	$\Xi_{bb}^0 \rightarrow \Xi_b^- K^+$	-1	$\Xi_{bb}^0 \rightarrow \Omega_b^- K^+$	$\Xi_{bb}^0 \rightarrow \Sigma_b^- \pi^+$	1
$\Xi_{bb}^- \rightarrow \Lambda_b^0 K^-$	$\Omega_{bb}^- \rightarrow \Xi_b^0 \pi^-$	1	$\Xi_{bb}^0 \rightarrow \Omega_b^- K^+$	$\Xi_{bb}^0 \rightarrow \Xi_b'^- K^+$	$\sqrt{2}$
$\Xi_{bb}^- \rightarrow \Xi_b^0 \pi^-$	$\Omega_{bb}^- \rightarrow \Lambda_b^0 K^-$	1	$\Xi_{bb}^- \rightarrow \Sigma_b^0 K^-$	$\Omega_{bb}^- \rightarrow \Xi_b^0 \pi^-$	1
$\Omega_{bb}^- \rightarrow \Xi_b^0 K^-$	$\Xi_{bb}^- \rightarrow \Lambda_b^0 \pi^-$	1	$\Xi_{bb}^- \rightarrow \Xi_b^0 \pi^-$	$\Omega_{bb}^- \rightarrow \Sigma_b^0 K^-$	1
$\Omega_{bb}^- \rightarrow \Xi_b^- \bar{K}^0$	$\Xi_{bb}^- \rightarrow \Xi_b^- K^0$	-1	$\Xi_{bb}^- \rightarrow \Sigma_b^- \bar{K}^0$	$\Omega_{bb}^- \rightarrow \Omega_b^- K^0$	1
$\Xi_{bb}^0 \rightarrow \Sigma_b^+ K^-$	$\Xi_{bb}^0 \rightarrow \Sigma_b^+ \pi^-$	1	$\Xi_{bb}^- \rightarrow \Omega_b^- K^0$	$\Omega_{bb}^- \rightarrow \Sigma_b^- \bar{K}^0$	1
$\Xi_{bb}^0 \rightarrow \Sigma_b^0 \bar{K}^0$	$\Xi_{bb}^0 \rightarrow \Xi_b^0 K^0$	1	$\Omega_{bb}^- \rightarrow \Xi_b^0 K^-$	$\Xi_{bb}^- \rightarrow \Sigma_b^0 \pi^-$	1
$\Xi_{bb}^0 \rightarrow \Xi_b'^- \pi^+$	$\Xi_{bb}^0 \rightarrow \Sigma_b^- \pi^+$	$\frac{1}{\sqrt{2}}$	$\Omega_{bb}^- \rightarrow \Xi_b'^- \bar{K}^0$	$\Xi_{bb}^- \rightarrow \Xi_b'^- K^0$	1

Table 22 U-spin relations for Ξ_{bb} , Ω_{bb} decays into a bottom meson and a light baryon. Results in the “channel 1” are for $b \rightarrow d$ processes and the ones in the “channel 2” are for $b \rightarrow s$ processes. r denotes the ratio of the two amplitudes

Channel 1	Channel 2	r	Channel 1	Channel 2	r
$\Xi_{bb}^0 \rightarrow p B^-$	$\Xi_{bb}^0 \rightarrow \Sigma^+ B^-$	-1	$\Xi_{bb}^- \rightarrow \Delta^0 B^-$	$\Omega_{bb}^- \rightarrow \Xi^0 B^-$	1
$\Xi_{bb}^0 \rightarrow n \bar{B}^0$	$\Xi_{bb}^0 \rightarrow \Xi^0 \bar{B}_s^0$	-1	$\Xi_{bb}^- \rightarrow \Sigma^- \bar{B}_s^0$	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	1
$\Xi_{bb}^- \rightarrow \Sigma^- \bar{B}_s^0$	$\Omega_{bb}^- \rightarrow \Xi^- \bar{B}^0$	-1	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	$\Xi_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	1
$\Xi_{bb}^- \rightarrow n B^-$	$\Omega_{bb}^- \rightarrow \Xi^0 B^-$	-1	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}^0$	1
$\Omega_{bb}^- \rightarrow \Sigma^- \bar{B}^0$	$\Xi_{bb}^- \rightarrow \Xi^- \bar{B}_s^0$	-1	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	$\Omega_{bb}^- \rightarrow \Omega^- \bar{B}_s^0$	$\frac{1}{\sqrt{3}}$
$\Omega_{bb}^- \rightarrow \Xi^- \bar{B}_s^0$	$\Xi_{bb}^- \rightarrow \Sigma^- \bar{B}^0$	-1	$\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}_s^0$	1
$\Xi_{bb}^0 \rightarrow \Delta^0 \bar{B}^0$	$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}^0$	$\sqrt{2}$	$\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\Xi_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	1
$\Xi_{bb}^0 \rightarrow \Delta^0 \bar{B}^0$	$\Xi_{bb}^0 \rightarrow \Xi^0 \bar{B}_s^0$	1	$\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}^0$	1
$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}_s^0$	$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}^0$	1	$\Omega_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\Omega_{bb}^- \rightarrow \Omega^- \bar{B}_s^0$	$\frac{1}{\sqrt{3}}$
$\Xi_{bb}^0 \rightarrow \Sigma^0 \bar{B}_s^0$	$\Xi_{bb}^0 \rightarrow \Xi^0 \bar{B}_s^0$	$\frac{1}{\sqrt{2}}$	$\Omega_{bb}^- \rightarrow \Sigma^0 B^-$	$\Xi_{bb}^- \rightarrow \Sigma^0 B^-$	1
$\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0$	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	$\sqrt{3}$	$\Omega_{bb}^- \rightarrow \Sigma^0 B^-$	$\Omega_{bb}^- \rightarrow \Xi^0 B^-$	$\frac{1}{\sqrt{2}}$
$\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0$	$\Xi_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\sqrt{3}$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\Xi_{bb}^- \rightarrow \Sigma'^- \bar{B}^0$	1
$\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}^0$	$\sqrt{3}$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\Xi_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	1
$\Xi_{bb}^- \rightarrow \Delta^- \bar{B}^0$	$\Omega_{bb}^- \rightarrow \Omega^- \bar{B}_s^0$	1	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}^0$	1
$\Xi_{bb}^- \rightarrow \Delta^0 B^-$	$\Xi_{bb}^- \rightarrow \Sigma^0 B^-$	$\sqrt{2}$	$\Omega_{bb}^- \rightarrow \Xi'^- \bar{B}_s^0$	$\Omega_{bb}^- \rightarrow \Omega^- \bar{B}_s^0$	$\frac{1}{\sqrt{3}}$

for the Ξ_{cc} and Ω_{cc} decays with the replacement of $T_{cc} \rightarrow T_{bc}$, $T_c \rightarrow T_b$ and $D \rightarrow B$. For the bottom quark decay, one can obtain them from those for Ξ_{bb} and Ω_{bb} decays with $T_{bb} \rightarrow T_{bc}$, $T_b \rightarrow T_c$ and $B \rightarrow D$. Thus it is not necessary to repeat the tedious calculations here.

7 Conclusions

Quite recently, the LHCb collaboration has observed the doubly charmed baryon Ξ_{cc}^{++} in the final state $\Lambda_c K^- \pi^+ \pi^+$. Such an important observation will undoubtedly promote the research on the hadron spectroscopy and also on weak decays of doubly heavy baryons.

In this paper, we have analyzed the weak decays of doubly heavy baryons Ξ_{cc} , Ω_{cc} , $\Xi_{bc}^{(\prime)}$, $\Omega_{bc}^{(\prime)}$, Ξ_{bb} and Ω_{bb} under the flavor SU(3) symmetry. The decay amplitudes for various

semileptonic and nonleptonic decays have been parametrized in terms of a few SU(3) irreducible amplitudes. We have found a number of relations or sum rules between decay widths and CP asymmetries, which can be examined in future measurements at experimental facilities like LHC, Belle II and CEPC. Moreover, once a few decay branching fractions have been measured in the future, some of these relations may provide hints for the exploration of new decay modes.

Acknowledgements The authors are grateful to Jibo He, Xiao-Hui Hu, Cai-Dian Lü, Fu-Sheng Yu, Zhen-Xing Zhao for useful discussions and valuable comments. We also thank Prof. Xiao-Gang He and Prof. De-Shan Yang for discussions of the SU(3) decomposition. This work is supported in part by National Natural Science Foundation of China under Grant nos. 11575110, 11655002, 11735010, Natural Science Foundation of Shanghai under Grant nos. 15DZ2272100 and 15ZR1423100, Shanghai Key Laboratory for Particle Physics and Cosmology, and by Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Funded by SCOAP³.

References

- M. Mattson et al., [SELEX Collaboration]. *Phys. Rev. Lett.* **89**, 112001 (2002). <https://doi.org/10.1103/PhysRevLett.89.112001>. [arXiv:hep-ex/0208014](https://arxiv.org/abs/hep-ex/0208014)
- A. Ocherashvili et al., [SELEX Collaboration]. *Phys. Lett. B* **628**, 18 (2005). <https://doi.org/10.1016/j.physletb.2005.09.043>. [arXiv:hep-ex/0406033](https://arxiv.org/abs/hep-ex/0406033)
- Y. Kato et al. [Belle Collaboration], *Phys. Rev. D* **89**(5), 052003 (2014) <https://doi.org/10.1103/PhysRevD.89.052003>. [arXiv:1312.1026](https://arxiv.org/abs/1312.1026) [hep-ex]
- R. Aaij et al., [LHCb Collaboration]. *JHEP* **1312**, 090 (2013). [https://doi.org/10.1007/JHEP12\(2013\)090](https://doi.org/10.1007/JHEP12(2013)090). [arXiv:1310.2538](https://arxiv.org/abs/1310.2538) [hep-ex]
- B. Aubert et al., [BaBar Collaboration]. *Phys. Rev. D* **74**, 011103 (2006). <https://doi.org/10.1103/PhysRevD.74.011103>. [arXiv:hep-ex/0605075](https://arxiv.org/abs/hep-ex/0605075)
- S.P. Ratti, *Nucl. Phys. Proc. Suppl.* **115**, 33 (2003). [https://doi.org/10.1016/S0920-5632\(02\)01948-5](https://doi.org/10.1016/S0920-5632(02)01948-5)
- R. Aaij et al. [LHCb Collaboration], *Phys. Rev. Lett.* **119**, no. 11, 112001 (2017) <https://doi.org/10.1103/PhysRevLett.119.112001>. [arXiv:1707.01621](https://arxiv.org/abs/1707.01621) [hep-ex]
- H.X. Chen, Q. Mao, W. Chen, X. Liu, S.L. Zhu, *Phys. Rev. D* **96**(3), 031501 (2017). <https://doi.org/10.1103/PhysRevD.96.031501>. [arXiv:1707.01779](https://arxiv.org/abs/1707.01779) [hep-ph]
- L. Meng, N. Li, S.l. Zhu. [arXiv:1707.03598](https://arxiv.org/abs/1707.03598) [hep-ph]
- X.H. Guo, H.Y. Jin, X.Q. Li, *Phys. Rev. D* **58**, 114007 (1998). <https://doi.org/10.1103/PhysRevD.58.114007>. [arXiv:hep-ph/9805301](https://arxiv.org/abs/hep-ph/9805301)
- M.A. Sanchez-Lozano, *Nucl. Phys. B* **440**, 251 (1995). [https://doi.org/10.1016/0550-3213\(95\)00064-Y](https://doi.org/10.1016/0550-3213(95)00064-Y). [arXiv:hep-ph/9502359](https://arxiv.org/abs/hep-ph/9502359)
- A. Faessler, T. Gutsche, M.A. Ivanov, J.G. Korner, V.E. Lyubovitskij, *Phys. Lett. B* **518**, 55 (2001). [https://doi.org/10.1016/S0370-2693\(01\)01024-3](https://doi.org/10.1016/S0370-2693(01)01024-3). [arXiv:hep-ph/0107205](https://arxiv.org/abs/hep-ph/0107205)
- D. Ebert, R.N. Faustov, V.O. Galkin, A.P. Martynenko, *Phys. Rev. D* **70**, 014018 (2004) Erratum: [*Phys. Rev. D* **77**, 079903 (2008)]. <https://doi.org/10.1103/PhysRevD.70.014018>. <https://doi.org/10.1103/PhysRevD.77.079903>. [arXiv:hep-ph/0404280](https://arxiv.org/abs/hep-ph/0404280)
- C. Albertus, E. Hernandez, J. Nieves, J.M. Verde-Velasco, *Eur. Phys. J. A* **31**, 691 (2007). <https://doi.org/10.1140/epja/i2006-10242-2>. [arXiv:hep-ph/0610131](https://arxiv.org/abs/hep-ph/0610131)
- E. Hernandez, J. Nieves, J.M. Verde-Velasco, *Phys. Lett. B* **663**, 234 (2008). <https://doi.org/10.1016/j.physletb.2008.03.072>. [arXiv:0710.1186](https://arxiv.org/abs/0710.1186) [hep-ph]
- J. M. Flynn, J. Nieves, *Phys. Rev. D* **76**, 017502 (2007) Erratum: [*Phys. Rev. D* **77**, 099901 (2008)] <https://doi.org/10.1103/PhysRevD.76.017502>. <https://doi.org/10.1103/PhysRevD.77.099901>. [arXiv:0706.2805](https://arxiv.org/abs/0706.2805) [hep-ph]
- C. Albertus, E. Hernandez, J. Nieves, *Phys. Lett. B* **683**, 21 (2010). <https://doi.org/10.1016/j.physletb.2009.11.048>. [arXiv:0911.0889](https://arxiv.org/abs/0911.0889) [hep-ph]
- A. Faessler, T. Gutsche, M.A. Ivanov, J.G. Korner, V.E. Lyubovitskij, *Phys. Rev. D* **80**, 034025 (2009). <https://doi.org/10.1103/PhysRevD.80.034025>. [arXiv:0907.0563](https://arxiv.org/abs/0907.0563) [hep-ph]
- R.H. Li, C.D. Lü, W. Wang, F.S. Yu, Z.T. Zou, *Phys. Lett. B* **767**, 232 (2017). <https://doi.org/10.1016/j.physletb.2017.02.003>. [arXiv:1701.03284](https://arxiv.org/abs/1701.03284) [hep-ph]
- F.S. Yu, H.Y. Jiang, R.H. Li, C.D. Lü, W. Wang, Z.X. Zhao. [arXiv:1703.09086](https://arxiv.org/abs/1703.09086) [hep-ph]
- W. Wang, F.S. Yu, Z.X. Zhao. [arXiv:1707.02834](https://arxiv.org/abs/1707.02834) [hep-ph]
- M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, *Phys. Rev. Lett.* **83**, 1914 (1999). <https://doi.org/10.1103/PhysRevLett.83.1914>. [arXiv:hep-ph/9905312](https://arxiv.org/abs/hep-ph/9905312)
- C.W. Bauer, S. Fleming, M.E. Luke, *Phys. Rev. D* **63**, 014006 (2000). <https://doi.org/10.1103/PhysRevD.63.014006>. [arXiv:hep-ph/0005275](https://arxiv.org/abs/hep-ph/0005275)
- C.W. Bauer, S. Fleming, D. Pirjol, I.W. Stewart, *Phys. Rev. D* **63**, 114020 (2001). <https://doi.org/10.1103/PhysRevD.63.114020>. [arXiv:hep-ph/0011336](https://arxiv.org/abs/hep-ph/0011336)
- M. Beneke, T. Feldmann, *Nucl. Phys. B* **685**, 249 (2004). <https://doi.org/10.1016/j.nuclphysb.2004.02.033>. [arXiv:hep-ph/0311335](https://arxiv.org/abs/hep-ph/0311335)
- Y.Y. Keum, H.N. Li, A.I. Sanda, *Phys. Lett. B* **504**, 6 (2001). [https://doi.org/10.1016/S0370-2693\(01\)00247-7](https://doi.org/10.1016/S0370-2693(01)00247-7). [arXiv:hep-ph/0004004](https://arxiv.org/abs/hep-ph/0004004)
- Y.Y. Keum, H.N. Li, A.I. Sanda, *Phys. Rev. D* **63**, 054008 (2001). <https://doi.org/10.1103/PhysRevD.63.054008>. [arXiv:hep-ph/0004173](https://arxiv.org/abs/hep-ph/0004173)
- C.D. Lu, K. Ukai, M.Z. Yang, *Phys. Rev. D* **63**, 074009 (2001). <https://doi.org/10.1103/PhysRevD.63.074009>. [arXiv:hep-ph/0004213](https://arxiv.org/abs/hep-ph/0004213)
- C.D. Lu, M.Z. Yang, *Eur. Phys. J. C* **23**, 275 (2002). <https://doi.org/10.1007/s100520100878>. [arXiv:hep-ph/0011238](https://arxiv.org/abs/hep-ph/0011238)
- T. Kurimoto, H.N. Li, A.I. Sanda, *Phys. Rev. D* **65**, 014007 (2002). <https://doi.org/10.1103/PhysRevD.65.014007>. [arXiv:hep-ph/0105003](https://arxiv.org/abs/hep-ph/0105003)
- W. Wang, *Phys. Lett. B* **708**, 119 (2012). <https://doi.org/10.1016/j.physletb.2012.01.036>. [arXiv:1112.0237](https://arxiv.org/abs/1112.0237) [hep-ph]
- T. Feldmann, M.W.Y. Yip, *Phys. Rev. D* **85**, 014035 (2012) Erratum: [*Phys. Rev. D* **86**, 079901 (2012)]. <https://doi.org/10.1103/PhysRevD.85.014035>. <https://doi.org/10.1103/PhysRevD.86.079901>. [arXiv:1111.1844](https://arxiv.org/abs/1111.1844) [hep-ph]
- T. Mannel, Y.M. Wang, *JHEP* **1112**, 067 (2011). [https://doi.org/10.1007/JHEP12\(2011\)067](https://doi.org/10.1007/JHEP12(2011)067). [arXiv:1111.1849](https://arxiv.org/abs/1111.1849) [hep-ph]
- A. Ali, C. Hamrock, A.Y. Parkhomenko, W. Wang, *Eur. Phys. J. C* **73**(2), 2302 (2013). <https://doi.org/10.1140/epjc/s10052-013-2302-4>. [arXiv:1212.3280](https://arxiv.org/abs/1212.3280) [hep-ph]
- W. Detmold, C.-J.D. Lin, S. Meinel, M. Wingate, *Phys. Rev. D* **87**(7), 074502 (2013). <https://doi.org/10.1103/PhysRevD.87.074502>. [arXiv:1212.4827](https://arxiv.org/abs/1212.4827) [hep-lat]
- G. Bell, T. Feldmann, Y.M. Wang, M.W.Y. Yip, *JHEP* **1311**, 191 (2013). [https://doi.org/10.1007/JHEP11\(2013\)191](https://doi.org/10.1007/JHEP11(2013)191). [arXiv:1308.6114](https://arxiv.org/abs/1308.6114) [hep-ph]
- M.J. Savage, M.B. Wise, *Phys. Rev. D* **39**, 3346 (1989) Erratum: [*Phys. Rev. D* **40**, 3127 (1989)]. <https://doi.org/10.1103/PhysRevD.39.3346>. [arXiv:hep-ph/9404283](https://arxiv.org/abs/hep-ph/9404283)
- B. Grinstein, R.F. Lebed, *Phys. Rev. D* **53**, 6344 (1996). <https://doi.org/10.1103/PhysRevD.53.6344>. [arXiv:hep-ph/9602218](https://arxiv.org/abs/hep-ph/9602218)
- X.G. He, *Eur. Phys. J. C* **9**, 443 (1999). <https://doi.org/10.1007/s100529900064>. [arXiv:hep-ph/9810397](https://arxiv.org/abs/hep-ph/9810397)
- N.G. Deshpande, X.G. He, J.Q. Shi, *Phys. Rev. D* **62**, 034018 (2000). <https://doi.org/10.1103/PhysRevD.62.034018>. [arXiv:hep-ph/0002260](https://arxiv.org/abs/hep-ph/0002260)
- X.G. He, J.Y. Leou, C.Y. Wu, *Phys. Rev. D* **62**, 114015 (2000). <https://doi.org/10.1103/PhysRevD.62.114015>. [arXiv:hep-ph/0007080](https://arxiv.org/abs/hep-ph/0007080)
- N.G. Deshpande, X.G. He, *Phys. Rev. Lett.* **75**, 1703 (1995). <https://doi.org/10.1103/PhysRevLett.75.1703>. [arXiv:hep-ph/9412393](https://arxiv.org/abs/hep-ph/9412393)

44. M. Gronau, Phys. Lett. B **492**, 297 (2000). [https://doi.org/10.1016/S0370-2693\(00\)01119-9](https://doi.org/10.1016/S0370-2693(00)01119-9). arXiv:hep-ph/0008292
45. X.G. He, Y.K. Hsiao, J.Q. Shi, Y.L. Wu, Y.F. Zhou, Phys. Rev. D **64**, 034002 (2001). <https://doi.org/10.1103/PhysRevD.64.034002>. arXiv:hep-ph/0011337
46. H.K. Fu, X.G. He, Y.K. Hsiao, J.Q. Shi, Chin. J. Phys. **41**, 601 (2003). arXiv:hep-ph/0206199
47. D.A. Egolf, R.P. Springer, J. Urban, Phys. Rev. D **68**, 013003 (2003). <https://doi.org/10.1103/PhysRevD.68.013003>. arXiv:hep-ph/0211360
48. C.W. Chiang, M. Gronau, Z. Luo, J.L. Rosner, D.A. Suprun, Phys. Rev. D **69**, 034001 (2004). <https://doi.org/10.1103/PhysRevD.69.034001>. arXiv:hep-ph/0307395
49. C.W. Chiang, M. Gronau, J.L. Rosner, D.A. Suprun, Phys. Rev. D **70**, 034020 (2004). <https://doi.org/10.1103/PhysRevD.70.034020>. arXiv:hep-ph/0404073
50. C.W. Chiang, Y.F. Zhou, JHEP **0612**, 027 (2006). <https://doi.org/10.1088/1126-6708/2006/12/027>. arXiv:hep-ph/0609128
51. Y. Li, C.D. Lu, W. Wang, Phys. Rev. D **77**, 054001 (2008). <https://doi.org/10.1103/PhysRevD.77.054001>. arXiv:0711.0497 [hep-ph]
52. W. Wang, Y.M. Wang, D.S. Yang, C.D. Lu, Phys. Rev. D **78**, 034011 (2008). <https://doi.org/10.1103/PhysRevD.78.034011>. arXiv:0801.3123 [hep-ph]
53. C.W. Chiang, Y.F. Zhou, JHEP **0903**, 055 (2009). <https://doi.org/10.1088/1126-6708/2009/03/055>. arXiv:0809.0841 [hep-ph]
54. H.Y. Cheng, C.W. Chiang, A.L. Kuo, Phys. Rev. D **91**(1), 014011 (2015). <https://doi.org/10.1103/PhysRevD.91.014011>. arXiv:1409.5026 [hep-ph]
55. X.G. He, G.N. Li, D. Xu, Phys. Rev. D **91**(1), 014029 (2015). <https://doi.org/10.1103/PhysRevD.91.014029>. arXiv:1410.0476 [hep-ph]
56. X.G. He, G.N. Li, Phys. Lett. B **750**, 82 (2015). <https://doi.org/10.1016/j.physletb.2015.08.048>. arXiv:1501.00646 [hep-ph]
57. M. He, X.G. He, G.N. Li, Phys. Rev. D **92**(3), 036010 (2015). <https://doi.org/10.1103/PhysRevD.92.036010>. arXiv:1507.07990 [hep-ph]
58. Y.K. Hsiao, C.F. Chang, X.G. He, Phys. Rev. D **93**(11), 114002 (2016). <https://doi.org/10.1103/PhysRevD.93.114002>. arXiv:1512.09223 [hep-ph]
59. X.G. He, W. Wang, R.L. Zhu, J. Phys. G **44**(1), 014003 (2017). <https://doi.org/10.1088/0954-3899/44/1/014003>, <https://doi.org/10.1088/0022-3727/44/27/274003>. arXiv:1606.00097 [hep-ph]
60. C.K. Chua, Phys. Rev. D **95**(9), 096004 (2017). <https://doi.org/10.1103/PhysRevD.95.096004>. arXiv:1612.04249 [hep-ph]
61. X.G. He, S.F. Li, B. Ren, X.B. Yuan, Phys. Rev. D **96**(5), 053004 (2017). <https://doi.org/10.1103/PhysRevD.96.053004>. arXiv:1704.05788 [hep-ph]
62. C.D. Lu, W. Wang, F.S. Yu, Phys. Rev. D **93**(5), 056008 (2016). <https://doi.org/10.1103/PhysRevD.93.056008>. arXiv:1601.04241 [hep-ph]
63. H.Y. Cheng, C.W. Chiang, A.L. Kuo, Phys. Rev. D **93**(11), 114010 (2016). <https://doi.org/10.1103/PhysRevD.93.114010>. arXiv:1604.03761 [hep-ph]
64. H.Y. Cheng, C.W. Chiang, Phys. Rev. D **86**, 014014 (2012). <https://doi.org/10.1103/PhysRevD.86.014014>. arXiv:1205.0580 [hep-ph]
65. H.Y. Cheng, S. Oh, JHEP **1109**, 024 (2011). [https://doi.org/10.1007/JHEP09\(2011\)024](https://doi.org/10.1007/JHEP09(2011)024). arXiv:1104.4144 [hep-ph]
66. H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, T.M. Yan, H.L. Yu, Phys. Rev. D **49**, 5857 (1994) Erratum: [Phys. Rev. D **55**, 5851 (1997)] <https://doi.org/10.1103/PhysRevD.49.5857>. <https://doi.org/10.1103/PhysRevD.55.5851.2>. arXiv:hep-ph/9312304
67. H.Y. Cheng, C.Y. Cheung, W. Dimm, G.L. Lin, Y.C. Lin, T.M. Yan, H.L. Yu, Phys. Rev. D **48**, 3204 (1993). <https://doi.org/10.1103/PhysRevD.48.3204>. arXiv:hep-ph/9305340
68. L.L. Chau, H.Y. Cheng, Phys. Lett. B **333**, 514 (1994). [https://doi.org/10.1016/0370-2693\(94\)90176-7](https://doi.org/10.1016/0370-2693(94)90176-7). arXiv:hep-ph/9404207
69. L.L. Chau, H.Y. Cheng, Phys. Lett. B **280**, 281 (1992). [https://doi.org/10.1016/0370-2693\(92\)90067-E](https://doi.org/10.1016/0370-2693(92)90067-E)
70. H.N. Li, C.D. Lu, F.S. Yu, Phys. Rev. D **86**, 036012 (2012). <https://doi.org/10.1103/PhysRevD.86.036012>. arXiv:1203.3120 [hep-ph]
71. H.N. Li, C.D. Lu, Q. Qin, F.S. Yu, Phys. Rev. D **89**(5), 054006 (2014). <https://doi.org/10.1103/PhysRevD.89.054006>. arXiv:1305.7021 [hep-ph]
72. C. Patrignani et al. [Particle Data Group], Chin. Phys. C **40**(10), 100001 (2016). <https://doi.org/10.1088/1674-1137/40/10/100001>
73. M. Gronau, D. Wyler, Phys. Lett. B **265**, 172 (1991). [https://doi.org/10.1016/0370-2693\(91\)90034-N](https://doi.org/10.1016/0370-2693(91)90034-N)
74. M. Gronau, D. London, Phys. Lett. B **253**, 483 (1991). [https://doi.org/10.1016/0370-2693\(91\)91756-L](https://doi.org/10.1016/0370-2693(91)91756-L)
75. I. Dunietz, Phys. Lett. B **270**, 75 (1991). [https://doi.org/10.1016/0370-2693\(91\)91542-4](https://doi.org/10.1016/0370-2693(91)91542-4)
76. D. Atwood, I. Dunietz, A. Soni, Phys. Rev. Lett. **78**, 3257 (1997). <https://doi.org/10.1103/PhysRevLett.78.3257>. arXiv:hep-ph/9612433
77. D. Atwood, I. Dunietz, A. Soni, Phys. Rev. D **63**, 036005 (2001). <https://doi.org/10.1103/PhysRevD.63.036005>. arXiv:hep-ph/0008090
78. A. Giri, Y. Grossman, A. Soffer, J. Zupan, Phys. Rev. D **68**, 054018 (2003). <https://doi.org/10.1103/PhysRevD.68.054018>. arXiv:hep-ph/0303187
79. W. Wang, Phys. Rev. D **85**, 051301 (2012). <https://doi.org/10.1103/PhysRevD.85.051301>. arXiv:1110.5194 [hep-ph]
80. C.S. Kim, R.H. Li, W. Wang, Phys. Rev. D **88**(3), 034003 (2013). <https://doi.org/10.1103/PhysRevD.88.034003>. arXiv:1305.5320 [hep-ph]