Regular Article - Theoretical Physics



Non-commutative deformation of Chern–Simons theory

Vladislav G. Kupriyanov^{1,2,a}

¹ CMCC-Universidade Federal do ABC, Santo André, SP, Brazil

² Tomsk State University, Tomsk, Russia

Received: 23 May 2019 / Accepted: 22 December 2019 / Published online: 18 January 2020 © The Author(s) 2020

Abstract The problem of the consistent definition of gauge theories living on the non-commutative (NC) spaces with a non-constant NC parameter $\Theta(x)$ is discussed. Working in the L_{∞} formalism we specify the undeformed theory, 3d abelian Chern–Simons, by setting the initial ℓ_1 brackets. The deformation is introduced by assigning the star commutator to the ℓ_2 bracket. For this initial set up we construct the corresponding L_{∞} structure which defines both the NC deformation of the abelian gauge transformations and the field equations covariant under these transformations. To compensate the violation of the Leibniz rule one needs the higher brackets which are proportional to the derivatives of Θ . Proceeding in the slowly varying field approximation when the star commutator is approximated by the Poisson bracket we derive the recurrence relations for the definition of these brackets for arbitrary Θ . For the particular case of su(2)-like NC space we obtain an explicit all orders formulas for both NC gauge transformations and NC deformation of Chern-Simons equations. The latter are non-Lagrangian and are satisfied if the NC field strength vanishes everywhere.

Contents

1	Introduction	1		
2	Basic facts from L_{∞} -algebras			
3	Non-commutative deformation of the abelian gauge			
	transformations	4		
	3.1 Leading order contribution	4		
	3.2 Next to the leading order	4		
	3.3 Higher brackets and recurrence relations	5		
4	4 Non-commutative field dynamics and L_{∞} structure			
	4.1 Leading order contribution	7		
	4.2 Next to the leading order	8		
	4.3 Higher order relations	9		
5	Lie-algebra like deformation	11		

	5.1	NC $su(2)$ -like deformation	11	
	5.2	Non-commutative Chern–Simons theory	12	
	5.3	Non-commutative field strength	14	
	5.4	Action principle	15	
	5.5	Summary of the results	15	
6	Con	clusions	16	
7	App	endix: Important algebraic relations	16	
Re	References			

1 Introduction

In the standard approach to the definition of gauge theory one needs the notion of the covariant derivative, $\mathcal{D}_a = \partial_a - iA_a$, as a generalization of the partial derivative ∂_a which transforms covariantly, $\mathcal{D}_a \rightarrow e^{if(x)}\mathcal{D}_a$, under the gauge transformations $\delta_f A_a = \partial_a f$. This notion is based on the Leibniz rule. The non-commutativity is a fundamental feature of the space-time which manifests itself at very short distances [1,2]. It can be introduced in the theory through the star product, defined in the theory of deformation quantization [3] as,

$$f \star g = f \cdot g + \frac{i}{2} \Theta^{ab}(x) \,\partial_a f \,\partial_b g + \cdots, \qquad (1.1)$$

where $\Theta^{ab}(x)$ is the non-commutativity parameter depending on the specific physical model. In some cases, like the open string dynamics in the constant *B*-field [2], the noncommutativity parameter can be constant, however in general it is a function of coordinates. The coordinate dependence of Θ , in general, leads to the violation of the Leibniz rule,

$$\partial_c (f \star g) = (\partial_c f) \star g + f \star (\partial_c g) + \frac{\iota}{2} \partial_c \Theta^{ab}(x) \partial_a f \partial_b g + \cdots,$$
(1.2)

and makes it impossible to follow the standard path for the formulation of NC gauge theory. Let us note that in some particular cases, like the NC gauge theory on D-branes in non-

^a e-mail: vladislav.kupriyanov@gmail.com

geometric backgrounds [4] the type of non-commutativity is compatible with the Leibniz rule, so the standard reasoning can be used for the definition of the NC field strength. Because of the non-geometry one has to shift the field strength tensor by a closed two-form on the D-brane worldvolume to construct the NC Yang–Mills action.

The problem with the violation of the Leibniz rule can be taken under control if, e.g., instead of the partial derivative ∂_a one takes the inner one defined through the star commutator, $D_a = i[\cdot, x_a]_{\star}$, as was done in the approach of covariant coordinates [5]. This however may lead to a problem with the correct commutative limit. Another possibility discussed in the literature consists in using the deformed Leibniz rule constructed with the help of the twist element of a Hopf algebra [6,7]. Here we mention that the twist element is known for the very few examples of NC spaces [8].

In recent work [9] in collaboration with Ralph Blumenhagen, Ilka Brunner and Dieter Lüst we have formulated the L_{∞} -bootstrap approach to the construction of noncommutative gauge theories. On the one hand, in the physical literature L_{∞} structures were introduced for description of gauge theories [10], see also [11, 12] for more details and recent references. On the classical level such an L_{∞} structure contains all necessary information about the theory including the gauge symmetry, the field equations and the Noether identities. On the other hand, L_{∞} algebras (also known as strong homotopy Lie algebras) [13,14] provides a framework for dealing with the deformation since the Jacobi identities are required to hold only up to a total derivative or a higher coherent homotopy. We note in particular that the proof of the key result in deformation quantization, the Formality Theorem, is based on the concept of L_{∞} algebras [15].

The main idea of the L_{∞} bootstrap approach consists in two steps. The first one is to represent the original undeformed gauge theory, like the Chern–Simons or the Yang– Mills, as well as the deformation introduced through the star commutator as a part of a new L_{∞} algebra specifying the initial brackets ℓ_1 , ℓ_2 , etc. Then solving the L_{∞} relations, $\mathcal{J}_n = 0$, one determines the missing brackets ℓ_n and completes the L_{∞} algebra which governs the NC deformation of the gauge transformations and the equations of motion. In [9] we found the expressions for the gauge transformations and the field equations up to the order $\mathcal{O}(\Theta^2)$ in the noncommutativity parameter. However the calculations were extremely involved and it was not clear whether the procedure can be extended to the higher or potentially all orders in the deformation parameter.

The purpose of the current work is to develop the ideas and tools proposed in [9] for the construction of the solution for the L_{∞} bootstrap program and to apply them to the explicit example, the non-commutative deformation of the abelian Chern–Simons theory. The key observation made in this paper is that in each given order *n* there is a set of *consistency conditions* for solvability of the L_{∞} bootstrap equations, $\mathcal{J}_n = 0$. These conditions are satisfied as a consequence of the previously solved equations, $\mathcal{J}_m = 0$, with m < n. We use this observation to express the brackets ℓ_n in terms of those which have already been found. Note that the solutions ℓ_n of the equations, $\mathcal{J}_n = 0$, are not unique and often can be chosen to be zero. Aiming to provide explicit calculations we work in the slowly varying field approximation when the higher derivative terms in the star commutator are discarded and it is approximated by the Poisson bracket. We set,

$$\ell_2(f,g) = -\{f,g\} = -\Theta^{ab}(x)\,\partial_a f\,\partial_b g,\tag{1.3}$$

where $\Theta^{ab}(x)$ is any suitably symmetric function satisfying the Jacobi identity,

$$\Theta^{al} \partial_l \Theta^{bc} + \Theta^{cl} \partial_l \Theta^{ab} + \Theta^{bl} \partial_l \Theta^{ca} = 0.$$
(1.4)

The paper is organized as follows. We start with a brief review of basic facts from L_{∞} algebras in the Sect. 5.47. In the Sect. 5.48 we construct the NC deformation of the abelian gauge transformation,

$$\delta_f A_a = \partial_a f + \{A_a, f\} + \Gamma_a^k(A) \,\partial_k f, \tag{1.5}$$

which satisfies the gauge closure condition, $[\delta_f, \delta_g]A_a = \delta_{\{f,g\}}A_a$. Then, in the Sect. 5.49 we derive an expression for the field equations,

$$\mathcal{F}^a := P^{abc}(A) \ \partial_b A_c + R^{abc}(A) \ \{A_b, A_c\} = 0, \qquad (1.6)$$

which are covariant under the transformation (1.5), i.e., $\delta_f \mathcal{F}^a = \{\mathcal{F}^a, f\}$, and reproduce in the commutative limit, $\Theta \rightarrow 0$, the standard abelian Chern–Simons equations, $\varepsilon^{abc} \partial_b A_c = 0$. For arbitrary NC parameter $\Theta^{ij}(x)$ we provide the recurrence relations for the construction of the functions $\Gamma_a^k(A)$, $P^{abc}(A)$ and $R^{abc}(A)$ at any order in Θ . For deformation parameter, which is linear in coordinates, in the Sect. 5 we obtain an explicit all orders expressions for both NC gauge transformations and NC field equations. We define the NC field strength and show that just like in the commutative case the NC Chern–Simons equations are equivalent to the requirement that the NC field strength should vanish everywhere.

2 Basic facts from $L_\infty\text{-algebras}$

For the convenience of the reader in this Section we will briefly review the basic facts form the theory of L_{∞} -algebras and its relation to the gauge theories. We start with a formal definition. In fact, L_{∞} -algebras are generalized Lie algebras where one has not only a two-bracket but more general multilinear *n*-brackets with *n* inputs

$$\ell_n: \qquad X^{\otimes n} \to X x_1, \dots, x_n \mapsto \ell_n(x_1, \dots, x_n),$$
(2.1)

defined on a graded vector space $X = \bigoplus_m X_m$, where $m \in \mathbb{Z}$, denotes the grading of the corresponding subspace. Each element $x \in X$, has its own degree, meaning that if deg(x) = p, this element belongs to the subspace X_p . These brackets are graded anti-symmetric,

$$\ell_n(\dots, x_1, x_2, \dots) = (-1)^{1 + \deg(x_1)\deg(x_2)} \ell_n(\dots, x_2, x_1, \dots).$$
(2.2)

The degree of ℓ_n is,

$$p = \deg(\ell_n(x_1, \dots, x_n)) = n - 2 + \sum_{i=1}^n \deg(x_i).$$
 (2.3)

The set of higher brackets ℓ_n define an L_∞ algebra if they satisfy the infinitely many relations

$$\mathcal{J}_{n}(x_{1},...,x_{n}) \\ \coloneqq \sum_{i+j=n+1}^{i+j=n+1} (-1)^{i(j-1)} \sum_{\sigma} (-1)^{\sigma} \chi(\sigma;x) \\ \ell_{j} \Big(\ell_{i}(x_{\sigma(1)},...,x_{\sigma(i)}), x_{\sigma(i+1)},...,x_{\sigma(n)} \Big) = 0.$$
(2.4)

The permutations are restricted to the ones with

$$\sigma(1) < \dots < \sigma(i), \qquad \sigma(i+1) < \dots < \sigma(n), \tag{2.5}$$

and the sign $\chi(\sigma; x) = \pm 1$ can be read off from (2.2).

The framework of L_{∞} algebras is quite flexible and it has been suggested that every classical perturbative gauge theory including its dynamics, is organized by an underlying L_{∞} structure [11]. To see this, let us assume that the field theory has a standard type gauge structure, meaning that the variations of the fields can be organized unambiguously into a sum of terms each of a definite power in the fields. We take the graded space, $X = X_0 \oplus X_{-1} \oplus X_{-2}$, and all others trivial. The assignment is as follows. X_0 corresponds to the space of gauge parameters or functions f, X_{-1} is the space of gauge fields A_a and X_{-2} contains the left hand side of the equations of motion of the gauge theory, $\mathcal{F}(A) = 0$. General elements in X_{-2} are denoted by the letter E, i.e.,

$$\begin{array}{cccc} X_0 & X_{-1} & X_{-2} \\ f & A_a & E_a \end{array}$$
(2.6)

The gauge variations are defined in terms of the brackets $\ell_{n+1}(f, A^n) \in X_{-1}$ as,

$$\delta_f A = \sum_{n \ge 0} \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} \ell_{n+1}(f, \underbrace{A, \dots, A}_{n \text{ times}})$$

= $\ell_1(f) + \ell_2(f, A) - \frac{1}{2} \ell_3(f, A, A) + \cdots$ (2.7)

The equations of motion can be written as

$$\mathcal{F} := \sum_{n \ge 1} \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} \ell_n(A^n)$$
$$= \ell_1(A) - \frac{1}{2} \ell_2(A^2) - \frac{1}{3!} \ell_3(A^3) + \cdots .$$
(2.8)

Using the L_{∞} relations (2.4) one can show that the commutator of gauge variations yields [11,18]

$$\left[\delta_f, \delta_g\right] A = \delta_{-C(f,g,A)} A + \delta_{-C(f,g,\mathcal{F},A)}^T A$$
(2.9)

with

$$C(f, g, A) = \sum_{n \ge 0} \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} \ell_{n+2}(f, g, \underbrace{A, \dots, A}_{n \text{ times}}).$$
(2.10)

and where the second term on the right hand side of (2.9) vanishes on-shell. It can be expanded as

$$C(f, g, \mathcal{F}, A) = \ell_3(f, g, \mathcal{F}) + \cdots$$

= $\sum_{n \ge 0} \frac{1}{n!} (-1)^{\frac{n(n-1)}{2}} \ell_{n+3}(f, g, \mathcal{F}, A^n).$
(2.11)

The gauge variation of the equation of motion $\mathcal F$ reads

$$\delta_f \mathcal{F} = \ell_2(f, \mathcal{F}) + \ell_3(f, \mathcal{F}, A) - \frac{1}{2}\ell_4(f, \mathcal{F}, A^2) + \dots$$
(2.12)

reflecting that, as opposed to the gauge field A, it transforms covariantly.

The same gauge algebra may correspond to the different gauge theories. So, following [11] it is convenient to introduce two different L_{∞} algebras. The first of them denoted by L_{∞}^{gauge} describes only the gauge transformations of fields A_a and is concentrated in two component space $X^{gauge} = X_0 \oplus X_{-1}$. The second, L_{∞}^{full} is concentrated in three component space, $X^{full} = X_0 \oplus X_{-1} \oplus X_{-2}$, and includes the information about the dynamics. This can be extended further as in [12], adding the fourth component X_{-3} containing the Noether identities, etc. For example, to define the L_{∞}^{gauge} algebra corresponding to the abelian gauge transformation, $\delta_f A_a = \partial_a f$, one sets $\ell_1(f) = \partial_a f$ and all other brackets vanishing. The L_{∞}^{full} algebra corresponding to the abelian Chern–Simons theory is defined by setting $\ell_1(A)^a = \varepsilon^{abc} \partial_b A_c$, while for Yang–Mills we define $\ell_1(A)^a = \Box A^a - \partial^a (\partial \cdot A)$. In both cases it is obvious that, $\ell_1(\ell_1(f)) = 0$.

We stress that in principle, the L_{∞} algebra may have an infinite number of the brackets ℓ_n , which however, are not arbitrary, since they should satisfy L_{∞} relations (2.4). As it was already mentioned in the introduction the idea of the L_{∞} bootstrap approach consists in representing the original undeformed gauge theory together with a deformation as a part of a new L_{∞} structure by setting initial brackets and solving L_{∞} relations to determine the L_{∞} algebra, which corresponds to the consistent deformation of the original theory.

3 Non-commutative deformation of the abelian gauge transformations

In this section we discuss a non-commutative deformation of the abelian L_{∞}^{gauge} algebra defined by the bracket, $\ell_1(f) = \partial_a f$. A deformation is introduced through the star commutator of functions which, from the consideration of anti-symmetry, should be assigned to the bracket,

$$\ell_2(f,g) = i[f,g]_{\star}.$$
(3.1)

For the Hermitean associative star product¹ an expression for the star commutator up to the third order in Θ reads [20],

$$i[f,g]_{\star}$$

$$= -\Theta^{kl} \partial_{k} f \partial_{l} g$$

$$+ \frac{1}{12} \left[\Theta^{nl} \partial_{l} \Theta^{mk} \partial_{n} \partial_{m} \Theta^{ij} \left(\partial_{i} f \partial_{j} \partial_{k} g - \partial_{i} g \partial_{j} \partial_{k} f \right) \right.$$

$$+ \frac{1}{2} \Theta^{nk} \partial_{n} \Theta^{jm} \partial_{m} \Theta^{il} \left(\partial_{i} \partial_{j} f \partial_{k} \partial_{l} g - \partial_{i} \partial_{j} g \partial_{k} \partial_{l} f \right)$$

$$+ \Theta^{ln} \partial_{l} \Theta^{jm} \Theta^{ik} \left(\partial_{i} \partial_{j} f \partial_{k} \partial_{n} \partial_{m} g - \partial_{i} \partial_{j} g \partial_{k} \partial_{n} \partial_{m} f \right)$$

$$+ \frac{1}{2} \Theta^{jl} \Theta^{im} \Theta^{kn} \partial_{i} \partial_{j} \partial_{k} f \partial_{l} \partial_{n} \partial_{m} g$$

$$+ \frac{1}{2} \Theta^{nk} \Theta^{ml} \partial_{n} \partial_{m} \Theta^{ij} \left(\partial_{i} f \partial_{j} \partial_{k} \partial_{l} g - \partial_{i} g \partial_{j} \partial_{k} \partial_{l} f \right) \right]$$

$$+ \mathcal{O} \left(\Theta^{5} \right) . \qquad (3.2)$$

In principle, the higher order terms in the star commutator can be constructed following the prescription of the Formality Theorem [15] and involves the Kontsevich graphs and the Kontsevich weights. The construction of the Kontsevich graphs is straightforward, however to the best of our knowledge there is no regular way of the computing the Kontsevich weights corresponding to the Kontsevich wheel diagrams. So, for arbitrary non-commutativity parameter $\Theta(x)$ an explicit all orders expression for the star commutator (3.2) is not known.

In this work we will consider the limit of slowly varying, but not necessarily small fields, i.e., we discard the higher derivatives terms, like $\partial f \partial \partial g$, etc., in the star commutator and approximate it by the Poisson bracket,

$$\ell_2(f,g) \approx -\Theta^{ab}(x) \,\partial_a f \,\partial_b g = -\{f,g\}.$$
(3.3)

This is a "self-consistent" approximation of non-commutativity. If we work with the NC deformations induced by the associative star product, the star commutator (3.1) satisfies the Jacobi identity, so does the corresponding Poisson bracket (3.3). Below we will see that in this case the brackets of the type $\ell_{n+2}(f, g, A^n)$, n > 0, can be taken to be zero.

3.1 Leading order contribution

Having non-vanishing brackets $\ell_1(f)$ and $\ell_2(f, g)$, one has to check the L_{∞} relation, $\mathcal{J}_2(f, g) = 0$, which reads,

$$\ell_1(\ell_2(f,g)) = -\{\overleftarrow{\partial_a f}, g\} - \{f, \overleftarrow{\partial_a g}\} - (\partial_a \Theta^{ij}) \partial_i f \partial_j g$$
$$= \ell_2(\ell_1(f), g) + \ell_2(f, \ell_1(g)), \qquad (3.4)$$

and involves the yet undetermined bracket $\ell_2(f, A)$. It means that now the identity, $\mathcal{J}_2(f, g) = 0$, becomes an equation on $\ell_2(f, A)$. Solving this equation one may proceed to the next L_{∞} relation, $\mathcal{J}_3(f, g, h) = 0$, and define the next bracket $\ell_3(f, g, A)$, etc.

Let us see how it works on practice. From (3.4) one finds

$$\ell_2(f,A) = -\{f,A_a\} - \frac{1}{2}(\partial_a \Theta^{ij}) \partial_i f A_j.$$
(3.5)

Note that the solution is not unique, one may also set, e.g.,

$$\ell'_{2}(f,A) = \ell_{2}(f,A) + s_{a}^{ij}(x) \,\partial_{i} f A_{j} \,, \tag{3.6}$$

with $s_a^{ij}(x) = s_a^{ji}(x)$. By the definition of L_{∞} , $\ell'_2(A, f) := -\ell'_2(f, A)$. The symmetry of $s_a^{ij}(x)$ implies that this choice of the bracket $\ell'_2(f, A)$ also satisfies the equation (3.4). However, the symmetric part $s_a^{ij}(x) \partial_i f A_j$ can always be "gauged away" by L_{∞} -quasi-isomorphism, physically equivalent to a Seiberg-Witten map [2], see [19] for more details.

3.2 Next to the leading order

Then we have to define the bracket $\ell_3(f, g, A)$ from the identity $\mathcal{J}_3(f, g, h) = 0$. Taking into account that according to

¹ These requirements to the star product are justified by the physical applications.

$$\begin{aligned} \mathcal{J}_{3}(f,g,h) \\ &:= \ell_{2}(\ell_{2}(f,g),h) + \ell_{2}(\ell_{2}(g,h),f) + \ell_{2}(\ell_{2}(h,f),g) \\ &+ \ell_{3}(\ell_{1}(f),g,h) + \ell_{3}(f,\ell_{1}(g),h) + \ell_{3}(f,g,\ell_{1}(h)) = 0 . \end{aligned}$$
(3.7)

The first line in the above expression is a Jacobiator for the bracket $\ell_2(f, g)$ defined in (1.3),

$$\ell_2(\ell_2(f,g),h) + \ell_2(\ell_2(g,h),f) + \ell_2(\ell_2(h,f),g)$$

= {{f,g},h} + {{h, f}, g} + {{g,h}, f} \equiv 0. (3.8)

So we may just set, $\ell_3(A, f, g) = 0$.

The next step is the crucial for the whole construction. We have to analyze the relation $\mathcal{J}_3(f, g, A) = 0$, given by

$$0 = \ell_2(\ell_2(A, f), g) + \ell_2(\ell_2(f, g), A) + \ell_2(\ell_2(g, A), f) + \ell_1(\ell_3(A, f, g)) - \ell_3(A, \ell_1(f), g) - \ell_3(A, f, \ell_1(g)).$$
(3.9)

For simplicity, we replace it with $\mathcal{J}_3(g, h, \ell_1(f)) = 0$, written in the form

$$\ell_{3}(\ell_{1}(f), \ell_{1}(g), h) - \ell_{3}(\ell_{1}(f), \ell_{1}(h), g) = G(f, g, h),$$

$$G(f, g, h) := \ell_{1}(\ell_{3}(\ell_{1}(f), g, h)) + \ell_{2}(\ell_{2}(\ell_{1}(f), g), h) + \ell_{2}(\ell_{2}(g, h), \ell_{1}(f)) + \ell_{2}(\ell_{2}(h, \ell_{1}(f)), g).$$
(3.10)

We will follow the logic of [20] for the solution of the above algebraic equation. By construction, the equation (3.10) is antisymmetric with respect to the permutation of g and h. The graded symmetry of the ℓ_3 bracket, $\ell_3(\ell_1(f), \ell_1(g), h) = \ell_3(\ell_1(g), \ell_1(f), h)$, implies the identity on the l.h.s. of (3.10):

$$\ell_{3}(\ell_{1}(f), \ell_{1}(g), h) - \ell_{3}(\ell_{1}(f), \ell_{1}(h), g) + \ell_{3}(\ell_{1}(h), \ell_{1}(f), g) - \ell_{3}(\ell_{1}(h), \ell_{1}(g), f) + \ell_{3}(\ell_{1}(g), \ell_{1}(h), f) - \ell_{3}(\ell_{1}(g), \ell_{1}(f), h) \equiv 0.$$

Which in turn requires the graded cyclicity of r.h.s. of the eq. (3.10),

$$G(f, g, h) + G(h, f, g) + G(g, h, f) = 0.$$
(3.11)

The latter is nothing but the consistency condition for the eq. (3.10).

It is remarkable that the consistency condition (3.11) follows from the previously satisfied L_{∞} relations, namely $\mathcal{J}_2(f, g) = 0$, and $\mathcal{J}_3(f, g, h) = 0$. Indeed, taking the definition of G(f, g, h), one writes

$$G(f, g, h) + G(h, f, g) + G(g, h, f)$$

= $\ell_2(\ell_2(\ell_1(h), f), g) + \ell_2(\ell_2(f, g), \ell_1(h))$

$$\begin{aligned} &+\ell_2(\ell_2(g,\ell_1(h)),f) + \ell_2(\ell_2(\ell_1(g),h),f) \\ &+\ell_2(\ell_2(h,f),\ell_1(g)) + \ell_2(\ell_2(f,\ell_1(g)),h) \\ &+\ell_2(\ell_2(\ell_1(f),g),h) + \ell_2(\ell_2(g,h),\ell_1(f)) \\ &+\ell_2(\ell_2(h,\ell_1(f)),g) + \ell_1(\ell_3(\ell_1(f),g,h)) \\ &+\ell_1(\ell_3(f,\ell_1(g),h)) + \ell_1(\ell_3(f,g,\ell_1(h))). \end{aligned}$$

Using $\mathcal{J}_2(f, g) = 0$, we may push ℓ_1 out of the brackets and rewrite it as

$$\ell_1 \Big[\ell_2(\ell_2(f,g),h) + \ell_2(\ell_2(g,h),f) + \ell_2(\ell_2(h,f),g) \\ + \ell_3(\ell_1(f),g,h) + \ell_3(f,\ell_1(g),h) + \ell_3(f,g,\ell_1(h)) \Big] \\ = \ell_1 \Big[\mathcal{J}_3(f,g,h) \Big] \equiv 0.$$

Which means that the consistency condition (3.11) holds true as a consequence of the previously satisfied L_{∞} relations.

Taking into account (3.11) one may easily check that the following expression (symmetrization in f and g of the r.h.s. of the Eq. (3.10)):

$$\ell_3(\ell_1(f), \ell_1(g), h) = -\frac{1}{6} \Big(G(f, g, h) + G(g, f, h) \Big),$$
(3.12)

has the required graded symmetry and solves, $\mathcal{J}_3(g, h, \ell_1(f)) = 0$. For $\ell_1(f) = A$, and $\ell_1(g) = B$, one gets,

$$\ell_3(A, B, f)_a = -\frac{1}{6} \Big(G_a^{kij} + G_a^{kji} \Big) A_i B_j \partial_k f, \qquad (3.13)$$

with

$$G_{a}^{kij} = -\Theta^{im}\partial_{m}\partial_{a}\Theta^{jk} - \frac{1}{2}\partial_{a}\Theta^{jm}\partial_{m}\Theta^{ki} - \frac{1}{2}\partial_{a}\Theta^{km}\partial_{m}\Theta^{ij}.$$
(3.14)

At this point we would like to stress three main observations. First, the consistency condition (graded cyclicity) (3.11) holds true as a consequence of the L_{∞} relations. Second, one needs higher brackets to compensate the violation of the standard Leibniz rule, which is standard in deformation quantization. And the last one is that the order of the bracket $\ell_{n+1}(F, A^n)$ in gauge fields A^n corresponds to the order of this brackets in the non-commutativity parameter Θ^n .

3.3 Higher brackets and recurrence relations

Once the brackets $\ell_3(f, g, A)$ and $\ell_3(f, A, B)$ are determined we may proceed to the next L_∞ relations and find the brackets with four, five, etc., entries. First let us discuss the relations with three gauge parameters, $\mathcal{J}_{n+3}(f, g, h, A^n) = 0$. The relation with four entries, $\mathcal{J}_4(f, g, h, A) = 0$, can be represented schematically as follows,

$$\mathcal{J}_4 := \ell_1 \ell_4 - \ell_4 \ell_1 + \ell_3 \ell_2 - \ell_2 \ell_3 = 0.$$
(3.15)

Recall that by the consideration of the degree, $\ell_2(f, g) \in X_0$, and $\ell_2(f, A) \in X_{-1}$, and also, $\ell_3(f, g, h) = 0$, and $\ell_4(f, g, h, A) = 0$, since the corresponding brackets belong to the space X_1 which is empty in our construction. Taking into account now that the bracket, $\ell_3(f, g, A) = 0$, due to (3.7), the identity (3.15) becomes,

$$\ell_4(\ell_1(f), g, h, A) + \ell_4(f, \ell_1(g), h, A) + \ell_4(f, g, \ell_1(h), A) = 0.$$
(3.16)

The latter can be satisfied setting, $\ell_4(f, g, A, B) = 0$. The same arguments show that if the bracket $\ell_2(f, g)$ satisfies the Jacobi identity (3.8), then the L_∞ relations with three gauge parameters, $\mathcal{J}_{n+3}(f, g, h, A^n) = 0$, are satisfied for, $\ell_{n+2}(f, g, A^n) = 0$, with $n \ge 1$. On the other hand, this means that the closure condition for the gauge algebra becomes

$$[\delta_f, \delta_g]A = \delta_{\{f,g\}}A. \tag{3.17}$$

To proceed with the relations with two gauge parameters, $\mathcal{J}_{n+2}(g, h, A^n) = 0$, we replace them by the equations, $\mathcal{J}_{n+2}(g, h, \ell_1(f)^n) = 0$, which in turn can be represented in the form

$$\ell_{n+2}(\ell_1(f)^n, \ell_1(g), h) -\ell_{n+2}(\ell_1(f)^n, \ell_1(h), g) = G(f_1, \dots, f_n, g, h), (3.18)$$

where the right hand side, $G(f_1, ..., f_n, g, h)$, is defined in terms of the previously defined brackets $\ell_{m+2}(\ell_1(f)^m, \ell_1(g), h)$, with m < n. It is symmetric in the first *n* arguments and antisymmetric in the last two by the construction. The graded symmetry of $\ell_{n+2}(\ell_1(f)^n, \ell_1(g), h)$ implies the non-trivial consistency condition (since $G(f_1, ..., f_n, g, h)$ is symmetric in first *n* arguments, one needs to check the cyclicity relation with respect to the permutation of the last three slots),

$$G(f_1, \dots, f_n, g, h) + G(f_1, \dots, f_{n-1}, g, h, f_n)$$

+G(f_1, \dots, f_{n-1}, h, f_n, g) = 0, (3.19)

which follows from the previous L_{∞} relations and can be proved by induction. The solution of the Eq. (3.18) can be constructed taking the symmetrization of the r.h.s. in the first n + 1 arguments.

The order by order in Θ calculations indicates the ansatz for the brackets,

$$\ell_{n+2} \left(f, A^{n+1} \right)_a$$

= $(n+1)! (-1)^{\frac{n(n-1)}{2}} \Gamma_a^{ki_1 \dots i_{n+1}} \partial_k f A_{i_1} \dots A_{i_{n+1}},$
(3.20)

yielding following expression for the gauge variation,

$$\delta_f A_a = \partial_a f + \{A_a, f\} + \Gamma_a^k(A) \,\partial_k f, \qquad (3.21)$$

where

$$\Gamma_{a}^{k}(A) = \sum_{n=1}^{\infty} \Gamma_{a}^{k(n)}, \qquad \Gamma_{a}^{k(n)} = \Gamma_{a}^{ki_{1}\dots i_{n}}(x) A_{i_{1}}\dots A_{i_{n}}.$$
(3.22)

According to the slowly varying field approximation we take into account only the leading order contribution in derivatives ∂f and ∂A . But we do not restrict the orders in Θ . The order of the term $\Gamma_a^{k(n)}$ in the gauge fields A coincides with the order of this term in deformation parameter Θ . To determine $\Gamma_a^k(A)$ we use the closure condition (3.17). One finds after simplification that,

$$\begin{split} \delta_{f} \left(\delta_{g} A_{a} \right) &- \delta_{g} \left(\delta_{f} A_{a} \right) - \delta_{\{f,g\}} A_{a} \\ &= \left(\frac{\delta \Gamma_{a}^{l}}{\delta A_{k}} - \frac{\delta \Gamma_{a}^{k}}{\delta A_{l}} - \left(\delta_{a}^{b} + \Gamma_{a}^{b} \right) \partial_{b} \Theta^{kl} \\ &- \partial_{b} \Gamma_{a}^{l} \Theta^{bk} + \partial_{b} \Gamma_{a}^{k} \Theta^{bl} - \frac{\delta \Gamma_{a}^{k}}{\delta A^{b}} \Gamma_{b}^{l} + \frac{\delta \Gamma_{a}^{l}}{\delta A^{b}} \Gamma_{b}^{k} \right) \partial_{k} f \partial_{l} g, \end{split}$$

$$(3.23)$$

where we use the notation, $\partial_b \Gamma_a^{l(n)} = (\partial_b \Gamma_a^{li_1...i_n}) A_{i_1} \dots A_{i_n}$. So, the closure condition (3.17) yields the following equation on $\Gamma_a^k(x, A)$,

$$\frac{\delta\Gamma_a^l}{\delta A_k} - \frac{\delta\Gamma_a^k}{\delta A_l} = \left(\delta_a^b + \Gamma_a^b\right) \partial_b \Theta^{kl} + \partial_b \Gamma_a^l \Theta^{bk} - \partial_b \Gamma_a^k \Theta^{bl} + \frac{\delta\Gamma_a^k}{\delta A^b} \Gamma_b^l - \frac{\delta\Gamma_a^l}{\delta A^b} \Gamma_b^k.$$
(3.24)

In what follows we will construct the perturbative in Θ solution of the above equation. We substitute the decomposition (3.22) in the Eq. (3.24) and compare the same orders in Θ from the left and from the right. In the first order in Θ we obtain,

$$\frac{\delta \Gamma_a^{l(1)}}{\delta A^k} - \frac{\delta \Gamma_a^{k(1)}}{\delta A^l} = G_a^{kl(1)} := \partial_a \Theta^{kl}, \qquad (3.25)$$

yielding a solution, $\Gamma_a^{k(1)} = -\partial_a \Theta^{kl} A_l/2$, which corresponds to (3.5). The second order in Θ gives,

$$\frac{\partial \Gamma_a^{l(2)}}{\delta A^k} - \frac{\partial \Gamma_a^{k(2)}}{\delta A^l} = G_a^{kl(2)} := -\frac{1}{2} G_a^{kli} A_i, \qquad (3.26)$$

where G_a^{kli} was defined in (3.14). The consistency condition for the Eq. (3.26) reads,

$$\frac{\delta G_a^{kl(2)}}{\delta A^i} + \frac{\delta G_a^{li(2)}}{\delta A^k} + \frac{\delta G_a^{ik(2)}}{\delta A^l} = 0.$$
(3.27)

It is equivalent to the requirement (3.11) and is satisfied as a consequence of the L_{∞} construction as it was discussed in the previous subsection. A solution of the Eq. (3.26),

$$\Gamma_a^{k(2)} = -\frac{1}{3} G_a^{kl(2)} A_l = \frac{1}{6} G_a^{kli} A_l A_i, \qquad (3.28)$$

represents exactly the contribution of the bracket $\ell_3(f, A, A)$ defined in (3.13) to the gauge variation (2.7).

In the higher orders in the deformation parameter Θ the Eq. (3.24) results in,

$$\frac{\partial \Gamma_a^{l(n+1)}}{\delta A^k} - \frac{\partial \Gamma_a^{k(n+1)}}{\delta A^l} = G_a^{kl(n+1)},$$

$$G_a^{kl(n+1)} := \frac{1}{2} \Gamma_a^{b(n)} \partial_b \Theta^{kl} + \partial_b \Gamma_a^{l(n)} \Theta^{bk} - \frac{1}{2} \partial_a \Theta^{kb} \Gamma_b^{l(n)}$$

$$+ \sum_{m=1}^{n-1} \frac{\delta \Gamma_a^{k(m+1)}}{\delta A^b} \Gamma_b^{l(n-m)} - (k \leftrightarrow l).$$
(3.29)

The consistency condition in this case,

$$\frac{\delta G_a^{kl(n+1)}}{\delta A^i} + \frac{\delta G_a^{li(n+1)}}{\delta A^k} + \frac{\delta G_a^{ik(n+1)}}{\delta A^l} = 0, \qquad (3.30)$$

is equivalent to the relations (3.19) which follow from the L_{∞} construction. A solution of the equation (3.29),

$$\Gamma_a^{k(n+1)} = -\frac{1}{n+2} \, G_a^{kl(n+1)} A_l, \tag{3.31}$$

defines the functions $\Gamma_a^{l(n+1)}$ in terms of the previously determined $\Gamma_a^{l(m)}$ with, $m \le n$.

The situation here is quite similar to the construction of the star product in the deformation quantization. For the arbitrary non-commutativity parameter $\Theta^{ab}(x)$ the best we can do is to provide the recurrence relations (3.31) for the construction of the gauge variation (3.21). However for the specific choices of Θ , like e.g., the linear one, it is possible to address the question of the convergence of the series (3.22) and provide all orders explicit formula for $\Gamma^k_a(A)$ in (3.21).

4 Non-commutative field dynamics and L_{∞} structure

In this Section we discuss the consistent deformation of the field dynamics in the bootstrap approach, considering the example of the non-commutative deformation of the abelian Chern–Simons theory. In this case we write the initial brackets as

$$\ell_1(f) = \partial_a f , \qquad \ell_1(A) = \varepsilon^{abc} \,\partial_b A_c , \qquad \ell_2(f,g) = -\{f,g\}.$$
(4.1)

The brackets $\ell_{n+1}(f, A^n)$ and $\ell_{n+2}(f, g, A^n)$ were determined in the Sect. 5.48. The rest of the brackets $\ell_n(A^n)$, $\ell_{n+2}(f, E, A^n)$, and $\ell_{n+3}(f, g, E, A^n)$, can be found from the corresponding l_{∞} relations.

4.1 Leading order contribution

The first new non-trivial L_{∞} relation is

$$\mathcal{J}_2(f, A) := \ell_1(\ell_2(f, A)) - \ell_2(\ell_1(f), A) - \ell_2(f, \ell_1(A)) = 0,$$
(4.2)

which we rewrite as

$$\ell_2(\ell_1(f), A) + \ell_2(f, \ell_1(A)) = \ell_1(\ell_2(f, A)).$$
(4.3)

In the r.h.s. the bracket, $\ell_2(f, A) \in X_{-1}$, is given by (3.5), while $\ell_1(A)$ is determined in (4.1), so

$$\ell_{1}(\ell_{2}(f, A)) = -\varepsilon^{abc} \{\partial_{b} f, A_{c}\} - \{f, \varepsilon^{abc} \partial_{b} A_{c}\} - \varepsilon^{abc} \partial_{b} \Theta^{kl} \partial_{k} f \partial_{l} A_{c} - \frac{1}{2} \varepsilon^{abc} \partial_{c} \Theta^{kl} \partial_{k} \partial_{b} f A_{l} - \frac{1}{2} \varepsilon^{abc} \partial_{c} \Theta^{kl} \partial_{k} f \partial_{b} A_{l}.$$
(4.4)

The brackets $\ell_2(\ell_1(f), A)$ and $\ell_2(f, \ell_1(A))$ in the l.h.s. of (4.3) need to be determined. The bracket $\ell_2(f, E)$ should be antisymmetric with respect to the permutation of its arguments, so we identify

$$\ell_2(f, \ell_1(A)) = -\{f, \varepsilon^{abc} \partial_b A_c\}, \quad \text{thus} \quad \ell_2(f, E) = -\{f, E_a\}.$$
(4.5)

The rest of the Eq. (4.3) can be written in the form

$$\ell_2(\ell_1(f), A) = Q_1^{abck} \partial_k f \partial_b A_c + S_1^{abck} A_k \partial_b \partial_c f + T_1^{abckl} \partial_k A_b \partial_l \partial_c f,$$
(4.6)

where the coefficient functions P_1^{aijk} , Q_1^{aijk} and R_1^{aijkl} are given by

$$Q_1^{abck} = \varepsilon^{acm} \partial_m \Theta^{kb} - \frac{1}{2} \varepsilon^{abm} \partial_m \Theta^{kc},$$

$$S_1^{abck} = -\frac{1}{4} \left(\varepsilon^{abm} \partial_m \Theta^{ck} + \varepsilon^{acm} \partial_m \Theta^{bk} \right),$$

$$T_1^{abckl} = -\varepsilon^{abc} \Theta^{kl}.$$
(4.7)

The solution of the Eq. (4.6) will be constructed following the logic of the previous section. There is a non-trivial consistency condition coming from the graded symmetry of the bracket ℓ_2 , which is satisfied as a consequence of the previously solved L_{∞} relations. The relation $\mathcal{J}_2(f, \ell_1(g)) = 0$, can be written as

$$\ell_2(\ell_1(f), \ell_1(g)) = \ell_1(\ell_2(f, \ell_1(g))).$$
(4.8)

The graded symmetry of ℓ_2 bracket,

$$\ell_2(\ell_1(f), \ell_1(g)) = \ell_2(\ell_1(g), \ell_1(f)), \tag{4.9}$$

implies the consistency condition on the right hand side of (4.8),

$$\ell_1(\ell_2(f,\ell_1(g))) - \ell_1(\ell_2(g,\ell_1(f))) = 0.$$
(4.10)

The later is automatically satisfied due to L_{∞} relation, $\mathcal{J}_2(f,g) = 0$, since

$$\ell_1(\ell_2(f,\ell_1(g))) - \ell_1(\ell_2(g,\ell_1(f))) = \ell_1[\ell_1(\ell_2(f,g)) - \mathcal{J}_2(f,g)] \equiv 0.$$
(4.11)

In the specific case of the deformation of Chern–Simons theory, i.e., Eq. (4.4) the relation (4.10) implies

$$\begin{aligned} Q_1^{abck} \partial_k f \partial_b \partial_c g + S_1^{abck} \partial_k g \partial_b \partial_c f + T_1^{abckl} \partial_k \partial_b g \partial_l \partial_c f \\ &= Q_1^{abck} \partial_k g \partial_b \partial_c f + S_1^{abck} \partial_k f \partial_b \partial_c g + T_1^{abckl} \partial_k \partial_b f \partial_l \partial_c g, \end{aligned}$$

$$(4.12)$$

which in turn yields the following relations between the coefficients Q_1^{aibck} , S_1^{abck} and T_1^{abckl} :

$$S_1^{abck} = Q_1^{a(bc)k}$$
, and $T_1^{abckl} = T_1^{acblk}$. (4.13)

We stress that these relations can be checked explicitly using just (4.7), however they follow from the L_{∞} algebra constructed so far. Using (4.13) the origynal equation (4.6) becomes

$$\ell_2(\ell_1(f), A) = Q_1^{abck} \left(\partial_k f \, \partial_b A_c + A_k \, \partial_b \partial_c f \right) + T_1^{abckl} \, \partial_k A_b \, \partial_l \partial_c f ,$$

$$(4.14)$$

implying the solution

$$\ell_{2}(B, A) = Q_{1}^{abck} \left(B_{k} \partial_{b} A_{c} + A_{k} \partial_{b} B_{c} \right) + T_{1}^{abckl} \partial_{k} A_{b} \partial_{l} B_{c}$$

$$= - \varepsilon^{abc} \{A_{b}, B_{c}\} + \varepsilon^{acm} \partial_{m} \Theta^{kb} \left(A_{k} \partial_{b} B_{c} + B_{k} \partial_{b} A_{c} \right)$$

$$- \frac{1}{2} \varepsilon^{abm} \partial_{m} \Theta^{kc} \left(A_{k} \partial_{b} B_{c} + B_{k} \partial_{b} A_{c} \right),$$

(4.15)

which is in the perfect agreement with our previous result [9].

D Springer

4.2 Next to the leading order

At this order there appears higher brackets ℓ_3 . The expressions for $\ell_3(A, f, g)$ and $\ell_3(A, B, f)$ were found in the previous section. Taking into account that now X_{-2} is non trivial, one may also have non-vanishing brackets $\ell_3(E, f, g) \in X_{-1}$, $\ell_3(E, A, f) \in X_{-2}$ and $\ell_3(A, B, C) \in X_{-2}$.

Let us start with $\ell_3(E, f, g)$. Such a term contributes to the closure condition, $\mathcal{J}_3(f, g, A) = 0$, which is however satisfied without it. Therefore, we can set $\ell_3(E, f, g) = 0$. Next we consider the L_{∞} relation, $\mathcal{J}_3(E, f, g) = 0$, i.e.,

$$0 = \ell_2(\ell_2(E, f), g) + \ell_2(\ell_2(g, E), f) + \ell_2(\ell_2(f, g), E) + \ell_3(E, \ell_1(f), g) + \ell_3(E, f, \ell_1(g)).$$
(4.16)

Since, $\ell_2(E, f) = \{E_a, f\}$, by (4.5), the first line in the above equation vanishes and one derives,

$$\ell_3(E, A, f) = 0. \tag{4.17}$$

Finally, to define $\ell_3(A, B, C)$, one has to solve, $\mathcal{J}(A, B, f) = 0$, written as,

$$\ell_{3}(A, B, \ell_{1}(f)) = r_{3}(A, B, f),$$

$$r_{3}(A, B, f) = -\ell_{1}(\ell_{3}(A, B, f)) - \ell_{3}(\ell_{1}(A), B, f)$$

$$+\ell_{3}(A, \ell_{1}(B), f) - \ell_{2}(\ell_{2}(A, B), f)$$

$$-\ell_{2}(\ell_{2}(f, A), B) + \ell_{2}(\ell_{2}(B, f), A).$$
(4.18)

By the construction the r.h.s., $r_3(A, B, f)$, is symmetric with respect to the permutation of *A* and *B*. As before, the graded symmetry of the ℓ_3 bracket,

$$\ell_3(A, \ell_1(g), \ell_1(f)) = \ell_3(A, \ell_1(f), \ell_1(g)), \qquad (4.19)$$

implies a non-trivial *consistency condition* on the r.h.s. of (4.18),

$$r_3(A, \ell_1(g), f) = r_3(A, \ell_1(f), g).$$
(4.20)

Using the previously satisfied L_{∞} relations, $\mathcal{J}_2(f, g) = 0$, and, $\mathcal{J}_2(A, f) = 0$, one may check that,

$$r_{3}(A, \ell_{1}(g), f) - r_{3}(A, \ell_{1}(f), g)$$

= $\ell_{1} (\mathcal{J}_{3}(g, f, A)) - \mathcal{J}_{3}(\ell_{1}(A), g, f) \equiv 0.$ (4.21)

Again the consistency condition (4.20) is satisfied as a consequence of L_{∞} construction.

Now let us discuss the solution of the Eq. (4.18) for the non-commutative deformation of CS theory. The calculation of the r.h.s. is quite involved, but straightforward. We represent it as

$$\ell_{3}(A, B, \ell_{1}(f)) = Q_{2}^{aijkl} (A_{i} \partial_{j} B_{k} \partial_{l} f + B_{i} \partial_{j} A_{k} \partial_{l} f) + S_{2}^{aijkl} (A_{i} B_{l} \partial_{j} \partial_{k} f + B_{i} A_{l} \partial_{j} \partial_{k} f) + T_{2}^{aijklm} (\partial_{i} f \partial_{j} A_{k} \partial_{l} B_{m} + \partial_{i} f \partial_{j} B_{k} \partial_{l} A_{m}) + U_{2}^{aijklm} (A_{i} \partial_{j} B_{k} \partial_{l} \partial_{m} f + B_{i} \partial_{j} A_{k} \partial_{l} \partial_{m} f),$$

$$(4.22)$$

where

$$\begin{aligned} \mathcal{Q}_{2}^{aijkl} &= \varepsilon^{abj} \left(\frac{1}{2} \Theta^{lm} \partial_{b} \partial_{m} \Theta^{ki} \right. \\ &+ \left. + \frac{1}{6} \Theta^{km} \partial_{b} \partial_{m} \Theta^{il} + \frac{1}{6} \Theta^{im} \partial_{b} \partial_{m} \Theta^{kl} \right. \\ &+ \left. \frac{1}{6} \partial_{b} \Theta^{km} \partial_{m} \Theta^{il} - \frac{1}{3} \partial_{b} \Theta^{im} \partial_{m} \Theta^{kl} \right) \\ &+ \varepsilon^{abk} \left(\Theta^{lm} \partial_{b} \partial_{m} \Theta^{ij} - \frac{1}{2} \Theta^{jm} \partial_{m} \partial_{m} \Theta^{il} \right. \\ &+ \left. \partial_{b} \Theta^{im} \partial_{m} \Theta^{jl} - \frac{1}{2} \partial_{b} \Theta^{jm} \partial_{m} \Theta^{il} \right) \\ &+ \varepsilon^{abc} \left(\frac{1}{2} \partial_{b} \Theta^{ij} \partial_{c} \Theta^{kl} - \frac{1}{2} \partial_{b} \Theta^{ik} \partial_{c} \Theta^{jl} \right) , \\ S_{2}^{aijkl} &= \varepsilon^{abl} \left(\frac{1}{6} \Theta^{im} \partial_{b} \partial_{m} \Theta^{jk} + \frac{1}{3} \partial_{b} \Theta^{im} \partial_{m} \Theta^{jk} \right) , \\ T_{2}^{aijklm} &= \frac{1}{2} \varepsilon^{abk} \Theta^{jl} \partial_{b} \Theta^{im} , \\ U_{2}^{aijklm} &= \frac{1}{2} \varepsilon^{abk} \Theta^{jm} \partial_{b} \Theta^{il} - \frac{1}{2} \varepsilon^{abm} \Theta^{jl} \partial_{b} \Theta^{ik} . \end{aligned}$$

The Eq. (4.20) implies the following relations on the coefficient functions

$$Q_{2}^{aijkl} = Q_{2}^{aljki},$$

$$Q_{2}^{aijkl} = S_{2}^{aijkl} + S_{2}^{aljki},$$

$$U_{2}^{aijk(lm)} = T_{2}^{aijk(lm)} + T_{2}^{ai(lm)jk}.$$
(4.24)

We stress that the above relations are not manifest from the explicit form of the coefficient functions P_2^{aijkl} , Q_2^{aijkl} , R_2^{aijklm} and $S_2^{aijk(lm)}$ given by (4.23) correspondingly. They follow from the L_{∞} relations, $\mathcal{J}_3(g, f, A) = 0$, $\mathcal{J}_3(E, g, f) = 0$, etc., which were also used to obtain the Eq. (4.20). The situation here is absolutely the same as in the previous Section for the construction of L_{∞}^{gauge} -algebra. The solution of the L_{∞} relations in each given order *n* imply the non-trivial consistency conditions, which in turn are satisfied due to the previously solved lower order L_{∞} relations.

The following expression

$$\ell_3 (A, B, C)$$

= $\frac{1}{2} Q_2^{aijkl} (A_i \partial_j B_k C_l + C_i \partial_j A_k B_l + B_i \partial_j C_k A_l$
+ $C_i \partial_j B_k A_l + B_i \partial_j A_k C_l + A_i \partial_j C_k B_l)$

$$+ T_{2}^{aijklm} \left(A_{i} \partial_{j} B_{k} \partial_{l} C_{m} + C_{i} \partial_{j} A_{k} \partial_{l} B_{m} + B_{i} \partial_{j} C_{k} \partial_{l} A_{m} \right. \\ + C_{i} \partial_{j} B_{k} \partial_{l} A_{m} + B_{i} \partial_{j} C_{k} \partial_{l} B_{m} + A_{i} \partial_{j} C_{k} \partial_{l} B_{m} \right),$$

$$(4.25)$$

by construction is symmetric in all arguments and due to the relations (4.24) satisfies the Eq. (4.22).

4.3 Higher order relations

Following the same logic as in the beginning of the Sect. 3.3 we conclude that the L_{∞} relations of the type, $\mathcal{J}_n(A^{n-3}E, f, g) = 0$, are satisfied for, $\ell_n(A^{n-2}, E, f) = 0$, for n > 0. Thus, we conclude that the condition (2.12) on the gauge variation of the field equation, $\mathcal{F} = 0$, defined by (2.8) becomes

$$\delta_f \mathcal{F} = \ell_2(f, \mathcal{F}) = \{\mathcal{F}, f\}. \tag{4.26}$$

That is, the gauge variation of the field equation is proportional to the field equation itself, i.e., it is gauge invariant on-shell. The proper field equations, $\mathcal{F} = 0$, are constructed from the brackets $\ell_n(A^n)$ which should be determined from the L_{∞} relations, $\mathcal{J}_n(f, A^{n-1}) = 0$. The latter can be schematically represented in the form

$$\ell_n\left(A^{n-1}, \ell_1(f)\right) = r_n\left(A^{n-1}, f\right),$$
(4.27)

where the r.h.s. $r_n(A^{n-1}, f)$ written in terms of the lower order brackets $\ell_m, m < n$, by the construction is symmetric in first n - 1 arguments. Like in the case of first two orders given by the Eqs. (4.6) and (4.22) correspondingly, the Eq. (4.27) has a non-trivial consistency condition. The graded symmetry of the ℓ_n -bracket on the l.h.s. of (4.27) implies the relation,

$$r_n\left(A^{n-2}, \ell_1(g), f\right) = r_n\left(A^{n-2}, \ell_1(f), g\right).$$
 (4.28)

This relation can be proved by induction. To construct a solution of the Eq. (4.27) one may follow the prescription of the previous subsection, in particular (4.22) and (4.25).

The form of the lower order brackets indicates the anzats,

$$\ell_n \left(A^n \right)^a = n! (-1)^{\frac{n(n-1)}{2}} \left(P^{abc(n-1)}(A) \,\partial_b A_c + R^{abc(n-2)}(A) \,\{A_b, A_c\} \right),$$
(4.29)

where,

$$P^{abc(0)} = \varepsilon^{abc}, \qquad P^{abc(n)} = P^{abci_1\dots i_n}(x) A_{i_1}\dots A_{i_n},$$

$$R^{abc(0)} = \frac{1}{2} \varepsilon^{abc}, \qquad R^{abc(n)} = R^{abci_1\dots i_n}(x) A_{i_1}\dots A_{i_n}.$$
(4.30)

🖄 Springer

The latter in turn means that for the l.h.s. of the field equation one writes,

$$\mathcal{F}^{a} = P^{abc} \left(A \right) \, \partial_{b} A_{c} + R^{abc} \left(A \right) \left\{ A_{b}, A_{c} \right\}, \tag{4.31}$$

with

$$P^{abc}(A) = \sum_{n=0}^{\infty} P^{abc(n)}, \text{ and } R^{abc}(A) = \sum_{n=0}^{\infty} R^{abc(n)},$$

(4.32)

The notations used here are similar to those in (3.22). The order of the term $P^{abc(n)}$ in the gauge fields A^n coincides with the order of this term in the deformation parameter Θ^n . By construction, R^{abc} (A) should be antisymmetric in b and c, since it is contracted with the Poisson bracket, $\{A_b, A_c\}$.

To determine the functions $P^{abc}(A)$ and $R^{abc}(A)$ we use the gauge covariance condition (4.26). Introducing the notation

$$\delta_f = \bar{\delta}_f + \{\cdot, f\},\tag{4.33}$$

one obtains in the l.h.s. of (4.26):

$$\delta_{f} \mathcal{F}_{a} = \left(\bar{\delta}_{f} P^{abc}\right) \partial_{b} A_{c} + P^{abc} \partial_{b} \left(\bar{\delta}_{f} A_{c}\right) + P^{abc} \partial_{b} \Theta^{kl} \partial_{k} A_{c} \partial_{l} f$$

$$+ P^{abc} \left\{A_{c}, \partial_{b} f\right\} + \left\{P^{abc} \partial_{b} A_{c}, f\right\}$$

$$+ \left(\bar{\delta}_{f} R^{abc}\right) \left\{A_{b}, A_{c}\right\} + \left\{R^{abc}, f\right\} \left\{A_{b}, A_{c}\right\}$$

$$+ 2 R^{abc} \left\{\bar{\delta}_{f} A_{b}, A_{c}\right\} + R^{abc} \left\{\left\{A_{b}, f\right\}, A_{c}\right\}$$

$$+ R^{abc} \left\{A_{b}, \left\{A_{c}, f\right\}\right\}.$$

$$(4.34)$$

While the r.h.s. of (4.26) is just given by

$$\{P^{abc} \partial_b A_c + R^{abc} \{A_b, A_c\}, f\}.$$
(4.35)

Taking into account that due to Jacobi identity,

$$R^{abc}\left(\{A_b, \{A_c, f\}\} + \{A_c, \{f, A_b\}\} + \{f, \{A_b, A_c\}\}\right) \equiv 0,$$
(4.36)

the Eq. (4.26) for the ansatz (4.31) becomes

$$(\bar{\delta}_f P^{abc}) \partial_b A_c + P^{abc} \partial_b (\bar{\delta}_f A_c) + P^{abc} \partial_b \Theta^{kl} \partial_k A_c \partial_l f + P^{abc} \{A_c, \partial_b f\} + (\bar{\delta}_f R^{abc}) \{A_b, A_c\} + 2 R^{abc} \{\bar{\delta}_f A_b, A_c\} = 0.$$

$$(4.37)$$

We set separately

$$\left(\bar{\delta}_{f} P^{abc}\right) \partial_{b} A_{c} + P^{abc} \partial_{b} \left(\bar{\delta}_{f} A_{c}\right) + P^{abc} \partial_{b} \Theta^{kl} \partial_{k} A_{c} \partial_{l} f = 0,$$
(4.38)

🖄 Springer

and

Let us analyze first the Eq. (4.38). We remind that by (3.21) and (4.33), $\bar{\delta}_f A_a = \partial_a f + \Gamma_a^k(A) \partial_k f$. So the Eq. (4.38) can be written as

$$\begin{bmatrix} \frac{\delta P^{abc}}{\delta A^{l}} \left(\delta_{l}^{k} + \Gamma_{l}^{k} \right) + P^{abl} \frac{\delta \Gamma_{l}^{k}}{\delta A^{c}} + P^{alc} \partial_{l} \Theta^{bk} \end{bmatrix} \partial_{b} A_{c} \partial_{k} f + P^{abc} \left(\delta_{c}^{k} + \Gamma_{c}^{k} \right) \partial_{b} \partial_{k} f = 0.$$
(4.40)

Thus, we obtain two separate conditions on P^{abc} :

$$\frac{\delta P^{abc}}{\delta A^l} \left(\delta_l^k + \Gamma_l^k \right) + P^{abl} \frac{\delta \Gamma_l^k}{\delta A^c} + P^{alc} \,\partial_l \Theta^{bk} = 0, \quad (4.41)$$

and

$$P^{abc}\left(\delta_{c}^{k}+\Gamma_{c}^{k}\right)+P^{akc}\left(\delta_{c}^{b}+\Gamma_{c}^{b}\right)=0.$$
(4.42)

From this point we act in the same way as in the Sect. 3.3, we will construct the perturbative in Θ solution of the above equations using the expression for Γ_l^k obtained in the previous section. The Eq. (4.41) in the first order in Θ reads,

$$-\frac{\delta P^{abc(1)}}{\delta A^k} = Q^{abck(1)} := \varepsilon^{acm} \partial_m \Theta^{kb} - \frac{1}{2} \varepsilon^{abm} \partial_m \Theta^{kc}.$$
(4.43)

The r.h.s. of the above equation is exactly the coefficient Q_1^{abck} determined in (4.7). Its solution,

$$P^{abc(1)} = -Q^{abck(1)}A_k, (4.44)$$

reproduces the contribution to the equations of motion (2.8) from the corresponding part of the bracket $\ell_2(A, A)$ defined in (4.15). This choice for $P^{abc(1)}$ also satisfies the Eq. (4.41) up to the first order in Θ . It can be checked explicitly, but also follows from the first of the Eq. (4.13), that is from the L_{∞} construction.

In the higher orders in Θ the Eq. (4.41) results in,

$$-\frac{\delta P^{abc(n+1)}}{\delta A^k} = Q^{abck(n+1)},$$
$$Q^{abck(n+1)} := \sum_{m=1}^n \frac{\delta P^{abc(n+1-m)}}{\delta A^l} \Gamma_l^{k(m)}$$

$$+\sum_{m=0}^{n} P^{abl(n-m)} \frac{\delta \Gamma^{k(m+1)}}{\delta A^c} + P^{alc(n)} \partial_l \Theta^b k.$$
(4.45)

The consistency condition for this equation reads,

$$\frac{\delta Q^{abck(n+1)}}{\delta A^l} = \frac{\delta Q^{abcl(n+1)}}{\delta A^k}.$$
(4.46)

It follows from the relation (4.28). A solution is

$$P^{abc(n+1)} = -\frac{1}{n+1} Q^{abck(n+1)} A_k.$$
(4.47)

As before, for arbitrary non-commutativity parameter $\Theta^{kl}(x)$ we can only provide the recurrence relations (4.47) for definition of the functions $P^{abc(n+1)}$. In the next Section we will construct an explicit all orders expression for $P^{abc}(A)$ for linear Θ .

Once $P^{abc}(A)$ is found, from (4.42) one determines,

$$R^{abc} = \frac{1}{2} \left[(\delta + \Gamma)^{-1} \right]_{k}^{b} P^{akc}.$$
 (4.48)

The antisymmetry, $R^{abc} = -R^{acb}$, follows from (4.42).

5 Lie-algebra like deformation

The main goal of this Section is to work out the most simple and at the same time non-trivial situation taking the noncommutativity parameter Θ to be linear function of the coordinates and satisfying the Jacobi identity. Physically it corresponds, for example, to the *Q*-flux backgrounds in open string theory [22].

5.1 NC su(2)-like deformation

We choose the non-commutativity parameter, $\Theta^{ij}(x) = 2\theta \varepsilon^{ijk} x_k$, which correspond to the rotationally invariant 3d NC space [23–27]. For the brevity of the calculations in most cases we will suppress the symbol θ in this and the following subsections. However, we will restore θ in the Sect. 5.3 where we provide the summary of the main findings of this section. The corresponding Poisson bracket is

$$\{f,g\} = 2\varepsilon^{ijk} x_k \ \partial_i f \ \partial_j g. \tag{5.1}$$

For the first two brackets with one gauge parameter one finds,

$$\ell_2(f, A)_a = \{A_a, f\} + \varepsilon_a{}^{bc}A_b\partial_c f$$

$$\ell_3(f, A, A)_a = -\frac{2}{3} \left(\partial_a f A^2 - \partial_b f A^b A_a\right),$$
(5.2)

with $A^2 = A_b A^b$. Then, using the recurrence relations (3.31) we observe that the brackets $\ell_{n+3}(f, A^n)$ with the odd *n* vanish, while for even *n* they have the structure

$$\ell_{n+3}(f, A^n) = \left(\partial_a f A^2 - \partial_b f A^b A_a\right) \chi_n(A^2), \tag{5.3}$$

for some monomial function $\chi_n(A^2)$. The combination of (5.2) and (5.3) in (3.21) results in the following ansatz for the gauge variation:

$$\delta_f A_a = \partial_a f + \{A_a, f\} + \varepsilon^{abc} A_b \partial_c f + \left(\partial_a f A^2 - \partial_b f A^b A_a\right) \chi(A^2),$$
(5.4)

where the function, $\chi(A^2) = \sum_n \chi_n(A^2)$, should be determined from the closure condition (3.17).

For the convenience of the reader here we repeat the calculation (3.23) for the specific form of the gauge variation (5.4),

$$\begin{split} \delta_{f} \left(\delta_{g} A_{a} \right) &- \delta_{g} \left(\delta_{f} A_{a} \right) - \delta_{\{f,g\}} A_{a} \\ &= \{ \delta_{f} A_{a}, g \} + \varepsilon^{abc} \delta_{f} A_{b} \partial_{c} g \\ &+ \left(2 \partial_{a} g A_{b} \delta_{f} A^{b} - \partial_{b} g \delta_{f} A^{b} a_{a} - \partial_{b} g A^{b} \delta_{f} A_{a} \right) \chi(A^{2}) \\ &+ \left(\partial_{a} g A^{2} - \partial_{b} g A^{b} A_{a} \right) \chi'(A^{2}) 2 A_{c} \delta_{f} A^{c} \\ &- \{ \delta_{g} A_{a}, f \} - \varepsilon^{abc} \delta_{g} A_{b} \partial_{c} f \\ &- \left(2 \partial_{a} f A_{b} \delta_{g} A^{b} - \partial_{b} f \delta_{g} A^{b} a_{a} - \partial_{b} f A^{b} \delta_{g} A_{a} \right) \chi(A^{2}) \\ &- \left(\partial_{a} f A^{2} - \partial_{b} f A^{b} A_{a} \right) \chi'(A^{2}) 2 A_{c} \delta_{g} A^{c} \\ &- \partial_{a} \{ f, g \} - \{ A_{a}, \{ f, g \} \} - \varepsilon^{abc} A_{b} \partial_{c} \{ f, g \} \\ &- \left(\partial_{a} \{ f, g \} A^{2} - \partial_{b} \{ f, g \} A^{b} A_{a} \right) \chi(A^{2}). \end{split}$$

After tedious but straightforward calculations we rewrite the r.h.s. of (5.5) as

$$\begin{bmatrix} \partial_a g \partial_b f A^b - \partial_a f \partial_b g A^b \end{bmatrix}$$

$$\left(1 + 3\chi(A^2) + A^2\chi^2(A^2) + 2A^2\chi'(A^2)\right).$$
(5.6)

That is, requiring that

$$2t\chi'(t) + 1 + 3\chi(t) + t\chi^{2}(t) = 0, \quad \chi(0) = -\frac{1}{3}, \quad (5.7)$$

we will obtain zero in the r.h.s. of (5.5). The solution of (5.7) is

$$\chi(t) = \frac{1}{t} \left(\sqrt{t} \cot \sqrt{t} - 1 \right).$$
(5.8)

Thus, we have obtained in (5.4) and (5.8) with, $t = \theta^2 A^2$, an explicit form of the non-commutative su(2)-like deformation of the abelian gauge transformations in the slowly varying field approximation. Following the lines described in [28] this result can be generalized for the non-commutative deformations along any linear Poisson structure $\Theta^{ij}(x)$.

5.2 Non-commutative Chern–Simons theory

The initial data in this case were specified in (4.1). The brackets $\ell_2(A, A)$ and $\ell_3(A^3)$ were calculated in (4.15) and (4.25) correspondingly. The resulting expression for the NC *su*(2)-like deformation of the abelian Chern–Simons equations of motion up to the order $\mathcal{O}(\Theta^3)$ is given by:

$$\mathcal{F}^{a} := \varepsilon^{abc} \partial_{b} A_{c} + \frac{1}{2} \varepsilon^{abc} \{A_{b}, A_{c}\} + \theta (2A^{b} \partial^{a} A_{b} - A^{a} \partial^{b} A_{b} - A^{b} \partial_{b} A^{a}) + \theta^{2} \left[-\frac{8}{3} \varepsilon^{abc} A^{2} \partial_{b} A_{c} + \frac{2}{3} \varepsilon^{abm} A_{m} A^{c} \partial_{b} A_{c} - 2 \varepsilon^{acm} A_{m} A^{b} \partial_{b} A_{c} + 2 \varepsilon^{bcm} A_{m} A^{a} \partial_{b} A_{c} \right] - \theta \{A^{2}, A_{a}\} + \mathcal{O}(\theta^{3}) = 0.$$
(5.9)

Definition of the P-term

Taking into account an explicit form of the gauge variation $\delta_f A_a$ given by (5.4), the Eqs. (4.41) and (4.42) in case of linear Θ become,

$$\frac{\delta P^{abc}}{\delta A_e} \left(1 + A^2 \chi\right) + \frac{\delta P^{abc}}{\delta A^m} \left(\varepsilon^{mne} A_n - A^m A^e \chi\right) + P^{abm} \varepsilon^{cem} + 2 P^{amc} \varepsilon^{bem} + 2 P^{abe} A^c \left(\chi + A^2 \chi'\right) - P^{abc} A^e \chi - P^{abm} A_m \delta^{ce} \chi - 2 P^{abm} A_m A^c A^e \chi' = 0,$$
(5.10)

and

$$(P^{abc} + P^{acb}) (1 + A^2 \chi) - P^{abm} \varepsilon^{cnm} A_n - P^{acm} \varepsilon^{bnm} A_n - P^{abn} A_n A^c \chi - P^{acn} A_n A^b \chi = 0.$$

$$(5.11)$$

Again the lower order brackets $\ell_n(A^n)$ indicate the anzatz

$$P^{abc}(A) = \varepsilon^{abc} F(A^{2}) + \varepsilon^{abm} A_{m} A^{c} G(A^{2}) + \varepsilon^{acm} A_{m} A^{b} H(A^{2}) + \varepsilon^{bcm} A_{m} A^{a} J(A^{2}) + A^{a} A^{b} A^{c} K(A^{2}) + A^{a} \delta^{bc} L(A^{2}) + A^{b} \delta^{ac} M(A^{2}) + A^{c} \delta^{ab} N(A^{2}).$$
(5.12)

The Eq. (5.11) implies the following relations on the coefficient functions:

$$G + H (1 + A^{2} \chi) - \chi F - M = 0,$$

$$K - \chi (L + M) - J - H = 0,$$

$$L (1 + A^{2} \chi) + F + A^{2} J = 0,$$

$$M (1 + A^{2} \chi) + N - F + A^{2} H = 0.$$

(5.13)

Our strategy is to substitute (5.12) in (5.10) and collect the coefficients at the different powers of fields A, modulo the A^2 . Starting with a quartic in A contribution, $A^a A^b A^c A^e$, then cubic in A structures, like $\varepsilon^{abm} A_m A^c A^e$, etc. down to the zero order in A terms like $\delta^{ab} \delta^{ce}$. Equating to zero these coefficients we will obtain the system of differential equations on the coefficient functions F, \ldots, N . We stress that there are algebraic relations involving the Levi-Civita tensors ε^{abc} and vector fields A^e described in the appendix. Using them we will reduce the number of different structures and thus the number of the equations on F, G, etc. These relations is not overfull. The Eq. (5.10) does have the solution.

We start writing a term quartic in A term in the l.h.s. of (5.10):

$$A^{a} A^{b} A^{c} A^{e} \left[2 K' - 2 \chi K - 2 \chi' \left(L + M + A^{2} K \right) + 2 A^{2} \chi' K \right].$$
(5.14)

The cubic in the field A contribution is given by

$$\varepsilon^{abm} A_m A^c A^e \left[2 G' - \chi G + 2 A^2 \chi' G \right] + \varepsilon^{acm} A_m A^b A^e \left[2 H' - 3\chi H \right] + \varepsilon^{bcm} A_m A^a A^e \left[2 J' - 3\chi J \right] + \varepsilon^{aem} A_m A^b A^c \left[2 \left(\chi + A^2 \chi' \right) H - K \right] + \varepsilon^{bem} A_m A^a A^c \left[2 \left(\chi + A^2 \chi' \right) J + K \right].$$
(5.15)

At this point for the first time we make use of the algebraic relation from the appendix to reduce the number of structures. Namely employing the identity

$$\varepsilon^{aem} A_m A^b - \varepsilon^{bem} A_m A^a = -\varepsilon^{abe} A^2 + \varepsilon^{abm} A_m A^e,$$
(5.16)

and setting, J = -H, one rewrites (5.15) as

$$\varepsilon^{abm} A_m A^c A^e \left[2 G' - \chi G + 2 A^2 \chi' G + 2 (\chi + A^2 \chi') H - K \right] + \varepsilon^{acm} A_m A^b A^e \left[2 H' - 3\chi H \right]$$

$$+ \varepsilon^{bcm} A_m A^a A^e \left[2 J' - 3 \chi J \right] + \varepsilon^{abe} A^c \left[A^2 K - 2 A^2 \left(\chi + A^2 \chi' \right) H \right].$$
(5.17)

We stress that now there has appeared the linear in A contribution coming from the cubic ones.

We continue with the quadratic in the fields A terms in the l.h.s. of (5.10),

$$\delta^{ae} A^{b} A^{c} \left[(1 + A^{2}\chi) K + G + 2(\chi + A^{2}\chi') M \right] + \delta^{be} A^{a} A^{c} \left[(1 + A^{2}\chi) K + G + 2(\chi + A^{2}\chi') L \right] + \delta^{ce} A^{a} A^{b} [K - J - H - \chi (L + M)] + \delta^{bc} A^{a} A^{e} [2L' - 2\chi L + J] + \delta^{ac} A^{b} A^{e} [2M' - 2\chi M - H] + \delta^{ab} A^{c} A^{e} [2N' - 2G] + \varepsilon^{acm} A_{m} \varepsilon^{ben} A_{n} H - \varepsilon^{aem} A_{m} \varepsilon^{bcn} A_{n} J.$$
(5.18)

Using the identity (7.5) from the appendix which we write here for the convenience of the reader,

$$\varepsilon^{acm} A_m \varepsilon^{ben} A_n$$

$$= \left(\delta^{ab} \,\delta^{ce} - \delta^{ae} \,\delta^{bc}\right) A^2$$

$$+ \,\delta^{bc} A^a \,A^e - \delta^{ce} A^a \,A^b - \delta^{ab} \,A^c \,A^e + \delta^{ae} \,A^b \,A^c,$$
(5.19)

we rewrite (5.18) as

$$\delta^{ab} \,\delta^{ce} \,A^{2} \,[H - J] - \delta^{ae} \,\delta^{bc} \,A^{2} \,H + \delta^{ac} \,\delta^{be} \,A^{2} \,J \\ + \,\delta^{ae} \,A^{b} \,A^{c} \left[(1 + A^{2}\chi) \,K + G + 2(\chi + A^{2}\chi') \,M + H \right] \\ + \,\delta^{be} \,A^{a} \,A^{c} \left[(1 + A^{2}\chi) \,K + G + 2(\chi + A^{2}\chi') \,L - J \right] \\ + \,\delta^{ce} \,A^{a} \,A^{b} \,[K - J - H - \chi(L + M) - H + J] \\ + \,\delta^{bc} \,A^{a} \,A^{e} \left[2 \,L' - 2\chi \,L + J + H \right] \\ + \,\delta^{ac} \,A^{b} \,A^{e} \left[2 \,M' - 2\chi \,M - H - J \right] \\ + \,\delta^{ab} \,A^{c} \,A^{e} \left[2 \,N' - 2G - H + J \right].$$
(5.20)

At this point it is convenient to invert the order. First we will analize the zero order in *A* contributions in the Eq. (5.10) and only then the linear in the fields *A* terms. Taking into account the first line of (5.20) the zero order in *A* terms in the l.h.s. of (5.10) are given by

$$\delta^{ac} \,\delta^{be} \left[F + \left(1 + A^2 \,\chi \right) M + J \right] \\ + \,\delta^{ae} \,\delta^{bc} \left[F + \left(1 + A^2 \,\chi \right) L - H \right] \\ + \,\delta^{ab} \,\delta^{ce} \left[N - 2F + H - J \right].$$
(5.21)

The significant simplification occurs if we set

$$H = -J = 0. (5.22)$$

In order to the Eq. (5.10) be satisfied the coefficients at the different structures in the l.h.s. should be equal to zero. Thus from (5.21) we get

$$L = M = -\frac{F}{1 + A^2 \chi}$$
, and $N = 2 F$. (5.23)

Equating to zero the coefficient at $\delta^{ce} A^a A^b$ in (5.20) one finds,

$$K = \chi \left(L + M \right). \tag{5.24}$$

Now let us return to the linear in A contributions to the left hand side of the Eq. (5.10). Taking into account (5.17) it can be written as

$$\varepsilon^{abc} A^{e} \left[2 F' - \chi F \right] + \varepsilon^{abe} A^{c} \left[\left(1 + A^{2} \chi \right) G + 2N + 2 \left(\chi + A^{2} \chi' \right) F \right] + A^{2} K - 2 A^{2} \left(\chi + A^{2} \chi' \right) H \right] + \varepsilon^{ace} A^{b} \left[\left(1 + A^{2} \chi \right) H + M \right] + \varepsilon^{bce} A^{a} \left[\left(1 + A^{2} \chi \right) J - L \right] + \varepsilon^{abm} A_{m} \delta^{ce} \left[G - \chi F \right] + \varepsilon^{acm} A_{m} \left(1 + A^{2} \chi \right) H - \varepsilon^{aem} A_{m} \delta^{bc} L + \varepsilon^{bcm} A_{m} \delta^{ae} \left(1 + A^{2} \chi \right) J + \varepsilon^{bem} A_{m} \delta^{ac} M.$$
(5.25)

Here we recall that because of the algebraic identities from the appendix not all structures in the above expression are independent. Now using these identities and previously defined coefficients we will reduce the number of terms in (5.25). First, using (7.4) and (5.23) we get rid of the terms,

$$-\varepsilon^{aem} A_m \,\delta^{bc} \,L + \varepsilon^{bem} A_m \,\delta^{ac} \,M,$$

substituting them with,

$$\varepsilon^{abe} A^c L - \varepsilon^{abm} A_m \delta^{ec} L$$

Then we utilize the identity (7.1) to convey the terms,

$$\varepsilon^{ace} A^b M - \varepsilon^{bce} A^a L,$$

through the

$$-\varepsilon^{abc} A^e L + \varepsilon^{abe} A^c L.$$

We use that, H = -J = 0, from (5.22), and also notice that due to (5.23) and (5.24) the coefficients *K*, *L* and *F* satisfy the relation,

$$2L + A^2 K = -2F,$$

We conclude that the linear in A contribution to the l.h.s. of the Eq. (5.10) given initially by (5.25) becomes,

$$\varepsilon^{abc} A^{e} \left[2 F' - \chi F - L \right] + \varepsilon^{abe} A^{c} \left[\left(1 + A^{2} \chi \right) G + 2 F + 2 \left(\chi + A^{2} \chi' \right) F \right] + \varepsilon^{abm} A_{m} \delta^{ce} \left[G - \chi F - L \right].$$
(5.26)

Again we set to zero the coefficients in (5.26) and obtain the relations

$$2F' = \chi F + L.$$
(5.27)

$$\left(1 + A^{2} \chi\right) G + 2 F + 2 \left(\chi + A^{2} \chi'\right) F = 0, \qquad (5.28)$$

and

$$G = \chi F + L, \tag{5.29}$$

The solution of the Eq. (5.27) with the initial condition, F(0) = 1, is

$$F(t) = \frac{\sin\sqrt{t}\cos\sqrt{t}}{\sqrt{t}}.$$
(5.30)

The relation (5.29) defines the function *G* in terms of previously found ones χ and *F*. The Eq. (5.28) is satisfied as a consequence of the relation (5.29) and the differential Eq. (5.7). The same happens, for exemple, with the equation,

$$L' - \chi \, L = 0, \tag{5.31}$$

resulting from the quadratic contribution (5.20). To show (5.31) one needs (5.7), (5.23) and (5.27). The careful check shows that the rest of the coefficients also vanishes. We stress that in order to the Eq. (5.10) hold the function $\chi(t)$ cannot be arbitrary, but necessarily the one which guarantees the condition (3.17), i.e., $[\delta_f, \delta_g]A = \delta_{\{f,g\}}A$.

Definition of R-term

Now using (5.4) and (5.12) in (4.48) we obtain after simplification,

$$R^{abc}(A) = \left(\varepsilon^{abc} + \delta^{ab} A^{c} - \delta^{ac} A^{b}\right) S\left(A^{2}\right) + \left(\varepsilon^{abm} A_{m} A^{c} - \varepsilon^{acm} A_{m} A^{b}\right) T\left(A^{2}\right),$$
(5.32)

where

$$S = \frac{F}{2(1+A^2\chi)}$$
, and $T = \frac{\chi F}{2(1+A^2\chi)}$.
(5.33)

Comparison to the lower order brackets

As a consistency check let us calculate the first order contributions to the equations of motion. Since

$$L = M = -\frac{\sin^2 \sqrt{t}}{t}, \qquad S = V = \frac{\sin^2 \sqrt{t}}{2t},$$
 (5.34)

one finds, L(0) = M(0) = -1. Then N(0) = 2F(0) = 2, and S(0) = 1/2, so the first order contribution is given by

$$2A_b\partial_a A_b - A_a\partial_b A_b - A_b\partial_b A_a + \frac{1}{2}\varepsilon^{abc}\{A_b, A_c\}, \quad (5.35)$$

which is in the perfect agreement with (5.9). Now,

$$F'(0) = -\frac{2}{3}, \quad G(0) = -\frac{4}{3}, \quad V(0) = \frac{1}{2},$$
 (5.36)

which results in

$$-\frac{2}{3}\varepsilon^{abc}A^2\partial_bA_c - \frac{4}{3}\varepsilon^{abm}A_mA^c\partial_bA_c + \{A_aA^2\}.$$
 (5.37)

The term with the Poisson bracket is exactly the same as in (5.9), but the coefficients at the first two terms are different. However, adding to the (5.37) the algebraic identity (7.2) from the Appendix multiplied by the factor -2,

$$-2\varepsilon^{abc}A^2 + 2\varepsilon^{bcm}A_mA^a - 2\varepsilon^{acm}A_mA^b + 2\varepsilon^{abm}A_mA^c \equiv 0,$$

we arrive exactly to the Eq. (5.9).

5.3 Non-commutative field strength

So far working in 3*d* we have constructed a vector $\mathcal{F}^{a}(A)$, which transforms covariantly under the NC gauge transformations (5.4), i.e., $\delta_{f}\mathcal{F}^{a} = \{f, \mathcal{F}^{a}\}$. The commutative limit of this vector gives the l.h.s. of the abelian Chern–Simons equations, $\lim_{\theta \to 0} \mathcal{F}^{a}(A) = \varepsilon^{abc} \partial_{b}A_{c}$. Now, using the Levi-Ciivita epsilon we may define the tensor

$$\mathcal{F}^{ab} := \varepsilon^{abc} \mathcal{F}_c = P^{abcd} (A) \ \partial_c A_d + R^{abcd} (A) \ \{A_c, A_d\} ,$$
(5.38)

which also transforms covariantly, $\delta_f \mathcal{F}^{ab} = \{f, \mathcal{F}^{ab}\},\$ and reproduces in the commutative limit the abelian field strength,

$$\lim_{\theta \to 0} \mathcal{F}_{ab} = \partial_a A_b - \partial_b A_a \,. \tag{5.39}$$

We call this tensor the non-commutative field strength. Its explicit form is given by,

$$\mathcal{F}^{ab} = \partial^{a} \left(A^{b} F(\theta^{2} A^{2}) \right) - \partial^{b} \left(A^{a} F(\theta^{2} A^{2}) \right) + \theta \varepsilon^{abc} \partial^{d} \left(A_{c} A_{d} L(\theta^{2} A^{2}) \right) + \theta F(\theta^{2} A^{2}) \varepsilon^{abc} \partial_{c} A^{2} - \frac{1}{2} \{ A^{a} L\left(\theta^{2} A^{2}\right), A^{b} \} - \frac{1}{2} \{ A^{a}, A^{b} L(\theta^{2} A^{2}) \} - \frac{\theta}{2} L(\theta^{2} A^{2}) \varepsilon^{abc} \{ A_{c}, A^{2} \},$$
(5.40)

where

$$F(\theta^2 A^2) = \frac{\sin\left(2\sqrt{\theta^2 A^2}\right)}{2\sqrt{\theta^2 A^2}}, \quad \text{and}$$
$$L(\theta^2 A^2) = -\frac{\sin^2\sqrt{\theta^2 A^2}}{\theta^2 A^2}.$$

Thus, just like in the commutative case, the non-commutative equations of motion, $\mathcal{F}^a(A) = 0$, are satisfied if the non-commutative field strength vanishes everywhere, $\mathcal{F}^{ab}(A) = 0$.

5.4 Action principle

To check whether exist an action *S* yielding the field equations, $\mathcal{F}_a = 0$, we use the criterium of second variational derivatives. If the equations are Lagrangian, i.e., $\mathcal{F}_a = \delta S / \delta A^a$, then

$$\frac{\delta \mathcal{F}_a}{\delta A^b} = \frac{\delta \mathcal{F}_b}{\delta A^a}.$$
(5.41)

One may easily check that, for $\mathcal{F}^a = P^{abc}(A) \partial_b A_c + R^{abc}(A) \{A_b, A_c\}$, this condition does not hold. In particular, because P^{abc} given by (5.12) is not an antisymmetric in *a* and *c*.

On the other hand, there is a known result about the rigidity of the Chern–Simons action [29], meaning essentially that up to the field redefinition any consistent deformation of the Chern–Simons action is proportional to the trivial one. Thus the absence of the action principle for the Eq. (4.31) means that possibly we obtained here some non-trivial deformation of the Chern–Simons theory.

5.5 Summary of the results

Let us summarise the main results of the Sect. 5. Consider the three dimensional space endowed with the Poisson bracket,

$$\{x^i, x^j\} = 2\theta \,\varepsilon^{ijk} x_k. \tag{5.42}$$

In the Sect. 5.1 we have constructed the non-commutative deformation of the abelian gauge transformation, given by,

$$\delta_f A_a = \partial_a f + \{A_a, f\} + \theta \,\varepsilon^{abc} A_b \partial_c f + \theta^2 \left(\partial_a f A^2 - \partial_b f A^b A_a \right) \chi \left(\theta^2 A^2 \right), \qquad (5.43)$$

where

$$\chi(t) = \frac{1}{t} \left(\sqrt{t} \cot \sqrt{t} - 1 \right), \qquad \chi(0) = -\frac{1}{3}, \tag{5.44}$$

which close the algebra

$$[\delta_f, \delta_g]A_a = \delta_{\{f,g\}}A_a. \tag{5.45}$$

The field equations covariant under the gauge transformation (5.43) and reproducing the abelian Chern–Simons equations in the commutative limit were obtained in the Sect. 5.2. An explicit form is,

 $\mathcal{F}^{a} := P^{abc}(A) \ \partial_{b}A_{c} + R^{abc}(A) \ \{A_{b}, A_{c}\} = 0, \qquad (5.46)$

where

$$P^{abc}(A) = \varepsilon^{abc} F\left(\theta^{2} A^{2}\right) + \theta^{2} \varepsilon^{abm} A_{m} A^{c} G\left(\theta^{2} A^{2}\right) + \theta^{3} A^{a} A^{b} A^{c} K\left(\theta^{2} A^{2}\right) + \theta A^{a} \delta^{bc} L\left(\theta^{2} A^{2}\right) + \theta A^{b} \delta^{ac} M\left(\theta^{2} A^{2}\right) + \theta A^{c} \delta^{ab} N\left(\theta^{2} A^{2}\right),$$
(5.47)

and

$$R^{abc}(A) = \varepsilon^{abc} S\left(\theta^2 A^2\right) + \theta^2 \left(\varepsilon^{abm} A_m A^c - \varepsilon^{acm} A_m A^b\right) T\left(\theta^2 A^2\right) + \theta \left(\delta^{ab} A^c - \delta^{ac} A^b\right) S\left(\theta^2 A^2\right), \qquad (5.48)$$

and the coefficient functions are determined in (5.22-5.24), (5.29), (5.30) and (5.33) as

$$F(t) = \frac{N(t)}{2} = \frac{\sin\sqrt{t}\cos\sqrt{t}}{\sqrt{t}},$$

$$G(t) = \frac{2\sqrt{t}\cos 2\sqrt{t} - \sin 2\sqrt{t}}{2t\sqrt{t}},$$

$$K(t) = -4T(t) = -\frac{2\sin\sqrt{t}}{t^2} \left(\sqrt{t}\cos\sqrt{t} - \sin\sqrt{t}\right),$$

$$L(t) = M(t) = -2S(t) = -\frac{\sin^2\sqrt{t}}{t}.$$
 (5.49)

The Eq. (5.46) are non-Lagrangian and as in the commutative case are equivalent to the requirement that the noncommutative field strength (5.40) should vanish everywhere. Further physical properties and applications will be discussed elsewhere.

6 Conclusions

To construct the L_{∞} structure with given initial terms one has to solve the L_{∞} relations, $\mathcal{J}_n = 0$. The key observation we made in this work is that in each given order n there is the consistency condition of the equation, $\mathcal{J}_n = 0$, which is satisfied as a consequence of the previously solved L_{∞} relations, $\mathcal{J}_m = 0, m < n$. Using this observation we were able to derive the recurrence relations for the construction of the L_{∞} algebra describing the NC deformation of the abelian Chern-Simons theory in the slowly varying field approximation. Using these recurrence relations we made a conjecture regarding the explicit all orders formula for the NC su(2)-like deformation of the gauge transformations $\delta_f A$ and the corresponding field equations, $\mathcal{F}(A) = 0$. The functional coefficients in the proposed ansatz were fixed from the closure conditions of the gauge algebra (3.17) and the requirement of the gauge covariance of the equations of motion (4.26)correspondingly.

We conclude that the problem formulated in the introduction regarding the existence of the solution to the L_∞ bootstrap program has a positive answer. Moreover we were able to find an explicit example of such a solution. Thus we can see that L_∞ algebra is not only a correct mathematical framework to deal with the deformations but also is a powerful tool for the construction of these deformations.

Acknowledgements I am grateful to Ralph Blumenhagen for fruitful discussions and to the Max Planck Institute for Physics for hospitality. I am also appreciative to the anonymous referees for the careful reading of the manuscript and valuable suggestions and remarks. This work was supported by the Grant 305372/2016-5 from the Conselho Nacional de Pesquisa (CNPq, Brazil) and the Grant 2019/11475-6 from the Fundação de Amparo á Pesquisa do Estado de São Paulo (FAPESP, Brazil).

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: There are no external data associated with the manuscript.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecomm ons.org/licenses/by/4.0/. Funded by SCOAP³.

7 Appendix: Important algebraic relations

Since we are in 3*d*, for any vector A^e one may check that,

$$\varepsilon^{abc} A^e - \varepsilon^{bce} A^a + \varepsilon^{cea} A^b - \varepsilon^{eab} A^c \equiv 0.$$
 (7.1)

The latter reflects the fact that in 3*d* any totally antisymmetric tensor of rank four vanishes. Contracting the above identity with A_e we arrive at,

$$\varepsilon^{abc} A^2 - \varepsilon^{bcm} A_m A^a + \varepsilon^{acm} A_m A^b - \varepsilon^{abm} A_m A^c \equiv 0.$$
(7.2)

Taking the derivative of (7.2) with respect to A_e one finds,

$$2 \varepsilon^{abc} A^{e} - \varepsilon^{abe} A^{c} - \varepsilon^{abm} A_{m} \delta^{ce} + \varepsilon^{ace} A^{b} + \varepsilon^{acm} A_{m} \delta^{be} - \varepsilon^{bce} A^{a} - \varepsilon^{bcm} A_{m} \delta^{ae} = 0.$$
(7.3)

Now using (7.1) in (7.3) we end up with

$$\varepsilon^{abc} A^e - \varepsilon^{abm} A_m \,\delta^{ce} + \varepsilon^{acm} A_m \,\delta^{be} - \varepsilon^{bcm} A_m \,\delta^{ae} \equiv 0.$$
(7.4)

One more identity we need is

$$\varepsilon^{acm} A_m \varepsilon^{ben} A_n = \left(\delta^{ab} \, \delta^{ce} - \delta^{ae} \, \delta^{bc} \right) A^2 + \delta^{bc} \, A^a \, A^e - \delta^{ce} \, A^a \, A^b - \delta^{ab} \, A^c \, A^e + \delta^{ae} \, A^b \, A^c.$$
(7.5)

It can be obtained from (7.2) contracting it with ε^{cen} and then renaming the indices.

References

- S. Doplicher, K. Fredenhagen, J.E. Roberts, The quantum structure of space-time at the Planck scale and quantum fields. Commun. Math. Phys. **172**, 187 (1995)
- 2. N. Seiberg, E. Witten, String theory and noncommutative geometry. JHEP **9909**, 032 (1999)
- F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, Quantum mechanics as a deformation of classical mechanics. Lett. Math. Phys. 1, 521 (1977)
- C. Hull, R.J. Szabo, Noncommutative gauge theories on D-branes in non-geometric backgrounds. arXiv:1903.04947 [hep-th]
- J. Madore, S. Schraml, P. Schupp, J. Wess, Gauge theory on noncommutative spaces. Eur. Phys. J. C 16, 161 (2000)
- M. Dimitrijevic, F. Meyer, L. Moller, J. Wess, Gauge theories on the kappa Minkowski space-time. Eur. Phys. J. C 36, 117 (2004)
- 7. D.V. Vassilevich, Twist to close. Mod. Phys. Lett. A 21, 1279 (2006)
- R.J. Szabo, Symmetry, gravity and noncommutativity. Class. Quant. Gravit. 23, R199 (2006)
- 9. R. Blumenhagen, I. Brunner, V. Kupriyanov, D. Lüst, Bootstrapping non-commutative gauge theories from L_{∞} algebras. JHEP **1805**, 097 (2018)
- B. Zwiebach, Closed string field theory: quantum action and the B–V master equation. Nucl. Phys. B **390**, 33 (1993)
- 11. O. Hohm, B. Zwiebach, L_{∞} algebras and field theory. Fortsch. Phys. **65**(3–4), 1700014 (2017)

- 12. B. Jurco, L. Raspollini, C. Saemann, M. Wolf, L_{∞} -algebras of classical field theories and the Batalin–Vilkovisky formalism. arXiv:1809.09899 [hep-th]
- T. Lada, J. Stasheff, Introduction to SH Lie algebras for physicists. Int. J. Theor. Phys. 32, 1087 (1993)
- 14. J. Stasheff, L_{∞} and A_{∞} structures: then and now. arXiv:1809.02526 [math.QA]
- M. Kontsevich, Deformation quantization of Poisson manifolds. 1. Lett. Math. Phys. 66, 157 (2003)
- 16. V.G. Kupriyanov, L_{∞} -bootstrap approach to non-commutative gauge theories. Fortsch. Phys. https://doi.org/10.1002/prop. 201910010. arXiv:1903.02867 [hep-th]
- F.A. Berends, G.J.H. Burgers, H. van Dam, On the theoretical problems in constructing interactions involving higher spin massless particles. Nucl. Phys. B 260, 295 (1985)
- 18. R. Fulp, T. Lada, J. Stasheff, sh-Lie algebras induced by Gauge transformations. Commun. Math. Phys. **231**, 25 (2002)
- 19. R. Blumenhagen, M. Brinkmann, V. Kupriyanov, M. Traube, On the uniqueness of L_{∞} bootstrap: quasi-isomorphisms are Seiberg–Witten maps. J. Math. Phys. **59**(12), 123505 (2018)
- V.G. Kupriyanov, D.V. Vassilevich, Star products made (somewhat) easier. Eur. Phys. J. C 58, 627–637 (2008)

- V.G. Kupriyanov, Recurrence relations for symplectic realization of (quasi)-Poisson structures. J. Phys. A 52(22), 225204 (2019)
- D. Luest, E. Malek, E. Plauschinn, M. Syvari, Open-string nonassociativity in an R-flux background. arXiv:1903.05581 [hep-th]
- 23. A.B. Hammou, M. Lagraa, M.M. Sheikh-Jabbari, Coherent state induced star product on R_{λ}^3 and the fuzzy sphere. Phys. Rev. D 66, 025025 (2002)
- J.M. Gracia-Bondia, F. Lizzi, G. Marmo, P. Vitale, Infinitely many star products to play with. JHEP 0204, 026 (2002). arXiv:hep-th/0112092
- P. Vitale, J.C. Wallet, Noncommutative field theories on R³_λ: toward UV/IR mixing freedom. JHEP **1304**, 115 (2013)
- V. Galikova, S. Kovacik, P. Presnajder, Laplace–Runge–Lenz vector in quantum mechanics in noncommutative space. J. Math. Phys. 54, 122106 (2013)
- V.G. Kupriyanov, A hydrogen atom on curved noncommutative space. J. Phys. A 46, 245303 (2013)
- 28. V.G. Kupriyanov, P. Vitale, Noncommutative \mathbb{R}^d via closed star product. JHEP **1508**, 024 (2015)
- 29. G. Barnich, M. Henneaux, Consistent couplings between fields with a gauge freedom and deformations of the master equation. Phys. Lett. B **311**, 123 (1993)