



# Hyperbolic and trigonometric hypergeometric solutions to the star-star equation

Erdal Catak<sup>1,a</sup>, Ilmar Gahramanov<sup>2,3,4,b</sup>, Mustafa Mullahasanoglu<sup>2,c</sup>

<sup>1</sup> Department of Physics, Istanbul University, 34134 Istanbul, Turkey

<sup>2</sup> Department of Physics, Bogazici University, Bebek, 34342 Istanbul, Turkey

<sup>3</sup> Department of Mathematics, Khazar University, Mehseti St. 41, AZ1096 Baku, Azerbaijan

<sup>4</sup> Institute of Radiation Problems, Azerbaijan National Academy of Sciences, B.Vahabzade St. 9, AZ1143 Baku, Azerbaijan

Received: 29 May 2022 / Accepted: 1 August 2022 / Published online: 5 September 2022  
© The Author(s) 2022

**Abstract** We construct the hyperbolic and trigonometric solutions to the star-star relation via the gauge/YBE correspondence by using the three-dimensional lens partition function and superconformal index for a certain  $\mathcal{N} = 2$  supersymmetric gauge dual theories. This correspondence relates supersymmetric gauge theories to exactly solvable models of statistical mechanics. The equality of partition functions for the three-dimensional supersymmetric dual theories can be written as an integral identity for hyperbolic and basic hypergeometric functions.

## Contents

1 Introduction	1
2 Star-star relation	2
3 Solutions to the star-star equation	2
4 Conclusions	6
Appendix A: Derivation of the star-star relation (3.9)	6
References	7

## 1 Introduction

The gauge/YBE correspondence [1,2] connecting supersymmetric gauge theories and integrable lattice models of statistical mechanics provides a powerful tool for studying spin models. It turns out that most of integrable edge-interacting (Ising-like) models in statistical mechanics [3–5] (and some IRF models [6–8]) can be obtained by this correspondence.

We will start with a very short account of this topic, the interested reader can find an exhaustive review on the subject in [9,10], where necessary information about the correspondence presented. Similar identities appear in integrable models of statistical mechanics. In this work, we present some new hypergeometric integral identities of hyperbolic and trigonometric types.

One of the striking features of recent developments in non-perturbative supersymmetric gauge theories is their deep relationship with interesting mathematical structures, see, e.g. [11–13]. At present, they provide the main source of many new identities for hypergeometric functions [14–18].

In a recent work [19], the authors constructed a new solution to the star-triangle equation. This was achieved by using a certain three-dimensional supersymmetric dual theories on the lens space  $S_b^3/\mathbb{Z}_r$ . The sufficient condition for the integrability of the lattice spin models is the star-star relation [20]. In this paper we present the corresponding star-star relation for the model studied in [19] and also for models discussed in [21–23] (the corresponding gauge theories live on the squashed sphere  $S_b^3$  and  $S^2 \times S^1$ ). In the context of the gauge/YBE correspondence, this relation can be obtained from the equality of three-dimensional  $\mathcal{N} = 2$  supersymmetric partition functions for a certain dual SQED theories. Our first two solutions to the star-star relation are given in terms of hyperbolic hypergeometric integrals (they are written in terms of hyperbolic gamma functions) and the last solution is a trigonometric type written in terms of basic hypergeometric integrals.

The paper is organized as follows. In Sect. 2 we briefly recall the star-star relation for the IRF models. Then we present solutions to the star-star equation.

<sup>a</sup> e-mail: [ecatak@istanbul.edu.tr](mailto:ecatak@istanbul.edu.tr)

<sup>b</sup> e-mail: [ilmar.gahramanov@boun.edu.tr](mailto:ilmar.gahramanov@boun.edu.tr) (corresponding author)

<sup>c</sup> e-mail: [mustafa.mullahasanoglu@boun.edu.tr](mailto:mustafa.mullahasanoglu@boun.edu.tr)

## 2 Star-star relation

We deal here with the integrable interaction-round-a-face (IRF) lattice spin models [24, 25]. In the IRF models, spin variables are located on the sites of the square lattice and interact via face. The sufficient condition for the integrability is the Yang–Baxter equation. In [26] it was shown that the Yang–Baxter equation for the face models can be reduced to study of the so-called star-star relation. The star-star relation contains essentially all the information needed to solve the lattice spin model. Here we mainly follow the work of Baxter<sup>1</sup> [26], therefore we refer the reader to the original paper for details.

We consider an IRF model with the face Boltzmann weight

$$R \begin{pmatrix} \sigma_4 & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix} = \sum_{m_i} \int dx_i W(\sigma_1, \sigma_i) W(\sigma_2, \sigma_i) \times W(\sigma_3, \sigma_i) W(\sigma_4, \sigma_i), \quad (2.1)$$

where the  $\sigma_i = (x_i; m_i)$  stands for the continuous valued spin  $x_i$  and discrete valued spin  $m_i$  and  $W(\sigma_i, \sigma_j)$  denotes the interaction between spins  $\sigma_i$  and  $\sigma_j$ . The Boltzmann weight  $W(\sigma_i, \sigma_j)$  solves the star-triangle relation for a certain integrable Ising-type lattice model

$$\sum_{m_i} \int dx_i W_{\alpha_1}(\sigma_1, \sigma_i) W_{\alpha_2}(\sigma_2, \sigma_i) W_{\alpha_3}(\sigma_3, \sigma_i) = \mathcal{R}(\alpha_1, \alpha_2, \alpha_3) W_{\eta-\alpha_1}(\sigma_1, \sigma_2) \times W_{\eta-\alpha_2}(\sigma_1, \sigma_3) W_{\eta-\alpha_3}(\sigma_2, \sigma_3), \quad (2.2)$$

where  $\alpha_1 + \alpha_2 + \alpha_3 = \eta$ . Here  $\alpha_i$  stands for the rapidity parameter (spectral parameter) and  $\mathcal{R}$  is a spin-independent function.

Now let us define the following Boltzmann weight

$$R \begin{pmatrix} \sigma_1 & \\ \sigma_2 & \sigma_3 \\ & \sigma_4 \end{pmatrix} = W(\sigma_1, \sigma_3) W(\sigma_2, \sigma_4) R \begin{pmatrix} \sigma_4 & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix}. \quad (2.3)$$

The system is integrable if the Boltzmann weight satisfies the following star-star relation

$$R_{(left)} \begin{pmatrix} \sigma_1 & \\ \sigma_2 & \sigma_3 \\ & \sigma_4 \end{pmatrix} = R_{(right)} \begin{pmatrix} \sigma_1 & \\ \sigma_2 & \sigma_3 \\ & \sigma_4 \end{pmatrix}. \quad (2.4)$$

The identity (2.4) is illustrated in Fig. 1, where we skipped the rapidity lines for convenience, for the full picture and explicit expressions the reader is referred to [20].

One can also see (2.4) as the following way with the definition (2.3) to be convinced with the pictorial representation of the star-star relation in Fig. 1

$$R \begin{pmatrix} \sigma_4 & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix} = \frac{W(\sigma_1, \sigma_2) W(\sigma_1, \sigma_4)}{W(\sigma_3, \sigma_2) W(\sigma_3, \sigma_4)} R \begin{pmatrix} \sigma_4 & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix}. \quad (2.5)$$

where  $R \begin{pmatrix} \sigma_4 & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix}$  functions differ with the spectral parameters [20] omitted in this study.

By using the star-star relation one obtains the following IRF Yang–Baxter equation (it is depicted in Fig. 2)

$$\sum_{m_i \in \mathbb{Z}} \int dx_i R \begin{pmatrix} \sigma_5 & \\ \sigma_6 & \sigma_i \\ & \sigma_1 \end{pmatrix} R \begin{pmatrix} \sigma_i & \sigma_3 \\ \sigma_1 & \sigma_2 \end{pmatrix} R \begin{pmatrix} \sigma_5 & \sigma_4 \\ \sigma_i & \sigma_3 \end{pmatrix} = \sum_{m_j \in \mathbb{Z}} \int dx_j R \begin{pmatrix} \sigma_6 & \sigma_j \\ \sigma_1 & \sigma_2 \end{pmatrix} R \begin{pmatrix} \sigma_5 & \sigma_4 \\ \sigma_6 & \sigma_j \end{pmatrix} R \begin{pmatrix} \sigma_4 & \\ \sigma_j & \sigma_3 \\ & \sigma_2 \end{pmatrix}, \quad (2.6)$$

where the summation and integration stand for the discrete and continuous spin variables, respectively. Note that there are several solutions to the IRF Yang–Baxter equation obtained via gauge/YBE correspondence [2, 3, 5, 7, 8].

## 3 Solutions to the star-star equation



By using the gauge/YBE correspondence one can systematically derive solution of the Yang–Baxter equation from calculations of supersymmetric gauge theory. In the context of this correspondence the Yang–Baxter equation expresses the identity of partition functions for supersymmetric dual pairs. Therefore the main step is to choose appropriate supersymmetric duality. Here we consider the following three-dimensional  $\mathcal{N} = 2$  dual theories [19, 23, 41]

- *theory A* has  $U(1)$  gauge group, six chiral multiplets with  $SU(3) \times SU(3) \times U(1)$  global symmetry group
- *theory B* consists of nine free “mesons” with the same global symmetry group as theory A.

The supersymmetric localization technique<sup>2</sup> [29] enables us to calculate the partition function on different manifolds. The results of Coulomb branch localization on  $S_b^3$ ,  $S_b^3/\mathbb{Z}_r$  and  $S^2 \times S^1$  are known (see, e.g. [30–32]) and we will use these results in order to construct the star-triangle relation and corresponding star-star relation in the next sections.

<sup>1</sup> We use the same notation as in [19].

<sup>2</sup> The short review of the three-dimensional supersymmetric localization can be found in [27, 28].

**Fig. 1** The star-star relation**Fig. 2** Yang–Baxter relation via the star-star relation. Spin variables live on vertices and interact via edges

### 3.1 Notations

Let us introduce some definitions and notations of special functions which we use in the paper. The  $q$ -Pochhammer symbol is defined as follows

$$(z; q)_{\infty} = \prod_{i=0}^{\infty} (1 - zq^i). \quad (3.1)$$

We use the shorthand notation

$$(z, x; q)_{\infty} = (z; q)_{\infty} (x; q)_{\infty}. \quad (3.2)$$

We also use hyperbolic gamma function which can be defined as

$$\gamma^{(2)}(z; \omega_1, \omega_2) = e^{\frac{\pi i}{2} B_{2,2}(z; \omega_1, \omega_2)} \frac{(e^{-2\pi i \frac{z}{\omega_2}} \tilde{q}; \tilde{q})_{\infty}}{(e^{-2\pi i \frac{z}{\omega_1}}; q)_{\infty}}, \quad (3.3)$$

with the parameters  $\tilde{q} = e^{2\pi i \omega_1 / \omega_2}$  and  $q = e^{-2\pi i \omega_2 / \omega_1}$  and the  $B_{2,2}(z; \omega_1, \omega_2)$  stands for the Bernoulli polynomial

$$B_{2,2}(z; \omega_1, \omega_2) = \frac{z^2 - z(\omega_1 + \omega_2)}{\omega_1 \omega_2} + \frac{\omega_1^2 + 3\omega_1 \omega_2 + \omega_2^2}{6\omega_1 \omega_2}. \quad (3.4)$$

The hyperbolic gamma function also has an integral representation<sup>3</sup>

$$\gamma^{(2)}(z; \omega_1, \omega_2) = \exp \left( - \int_0^{\infty} \frac{dx}{x} \left[ \frac{\sinh x(2z - \omega_1 - \omega_2)}{2 \sinh(x\omega_1) \sinh(x\omega_2)} - \frac{2z - \omega_1 - \omega_2}{2x\omega_1 \omega_2} \right] \right), \quad (3.5)$$

<sup>3</sup> One can find different integral representations in [33, 34].

where  $\text{Re}(\omega_1), \text{Re}(\omega_2) > 0$  and  $\text{Re}(\omega_1 + \omega_2) > \text{Re}(z) > 0$ .

Additionally, we will use the following reflection property for hyperbolic gamma function

$$\gamma^{(2)}(\omega_1 + \omega_2 - z; \omega_1, \omega_2) \gamma^{(2)}(z; \omega_1, \omega_2) = 1. \quad (3.6)$$

### 3.2 Solution via $S_b^3$ supersymmetric partition function

The equivalence of the partition functions for dual theories on  $S_b^3$  gives the following hyperbolic hypergeometric integral identity [1, 35, 36]

$$\begin{aligned} & \int_{-i\infty}^{i\infty} \prod_{i=1}^3 \gamma^{(2)}(a_i - x; \omega_1, \omega_2) \gamma^{(2)}(b_i + x; \omega_1, \omega_2) \frac{dx}{i\sqrt{\omega_1 \omega_2}} \\ &= \prod_{i,j=1}^3 \gamma^{(2)}(a_i + b_j; \omega_1, \omega_2), \end{aligned} \quad (3.7)$$

with the balancing condition  $\sum_{i=1}^3 (a_i + b_i) = \omega_1 + \omega_2$ . In [21, 37] it was shown that this integral gives the star-triangle relation for the Faddeev–Volkov model which has the following Boltzmann weight

$$\begin{aligned} W_{\alpha}(x_i, x_j) &= \gamma^{(2)}(-\alpha + x_i - x_j; \omega_1, \omega_2) \\ &\quad \gamma^{(2)}(-\alpha - x_i + x_j; \omega_1, \omega_2), \end{aligned} \quad (3.8)$$

where we introduced new variables  $a_i = -\alpha_i + x_i$  and  $b_i = -\alpha_i - x_i$ . Here  $\alpha$  is a spectral parameter,  $x_i$  is a spin variable and  $\omega_1, \omega_2$  are temperature-like parameters.

The corresponding star-star relation for this model (for details, see Appendix A) has the following form

$$\begin{aligned} & \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(a_i - x; \omega_1, \omega_2) \gamma^{(2)}(b_i + x; \omega_1, \omega_2) \frac{dx}{i\sqrt{\omega_1\omega_2}} \\ &= \frac{\prod_{i,j=1}^2 \gamma^{(2)}(a_i + b_j; \omega_1, \omega_2)}{\prod_{i,j=3}^4 \gamma^{(2)}(\tilde{a}_i + \tilde{b}_j; \omega_1, \omega_2)} \\ & \times \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(\tilde{a}_i - z; \omega_1, \omega_2) \gamma^{(2)}(\tilde{b}_i + z; \omega_1, \omega_2) \frac{dz}{i\sqrt{\omega_1\omega_2}}, \end{aligned} \quad (3.9)$$

with a new balancing condition  $\sum_{i=1}^4 (a_i + b_i) = 2(\omega_1 + \omega_2)$  and we used the following notations

$$\begin{aligned} \tilde{a}_i &= a_i + s, & \tilde{b}_i &= b_i + s, & \text{if } i &= 1, 2, \\ \tilde{a}_i &= a_i - s, & \tilde{b}_i &= b_i - s, & \text{if } i &= 3, 4, \end{aligned} \quad (3.10)$$

with

$$\begin{aligned} s &= \frac{1}{2}(\omega_1 + \omega_2 - a_1 - a_2 - b_1 - b_2) \\ &= \frac{1}{2}(-\omega_1 - \omega_2 + a_3 + a_4 + b_3 + b_4). \end{aligned} \quad (3.11)$$

This integral identity was obtained in [38]. The physical interpretation of this identity discussed in [39].

### 3.3 Solution via $S_b^3/\mathbb{Z}_r$ supersymmetric partition function

We again start with the equivalence of the partition functions for dual theories. This time we consider the dual theories on  $S_b^3/\mathbb{Z}_r$  and obtain the following hyperbolic hypergeometric integral identity<sup>4</sup> [19]

$$\begin{aligned} & \sum_{y=0}^{[r/2]} \epsilon(y) e^{-\pi i y} \int_{-\infty}^{\infty} \prod_{i=1}^3 \\ & \gamma^{(2)}(-i(a_i - x) - i\omega_1(u_i - y); -i\omega_1 r, -i\omega) \\ & \gamma^{(2)}(-i(a_i - x) - i\omega_2(r - (u_i - y)); -i\omega_2 r, -i\omega) \\ & \gamma^{(2)}(-i(b_i + x) - i\omega_1(v_i + y); -i\omega_1 r, -i\omega) \\ & \gamma^{(2)}(-i(b_i + x) - i\omega_2(r - (v_i + y)); \\ & \quad -i\omega_2 r, -i\omega) \frac{dx}{r\sqrt{-\omega_1\omega_2}} = e^{\frac{-\pi i}{2} \sum_{i=1}^3 (u_i - v_i)} \\ & \prod_{i,j=1}^3 \gamma^{(2)}(-i(a_i + b_j) - i\omega_1(u_i + v_j); \\ & \quad -i\omega_1 r, -i\omega) \gamma^{(2)}(-i(a_i + b_j) - i\omega_2 \end{aligned}$$

<sup>4</sup> Note that this integral identity was obtained via the reduction procedure and it needs to be proven rigorously.

$$\times (r - (u_i + v_j)); -i\omega_2 r, -i\omega), \quad (3.12)$$

with the balancing conditions  $\sum_{i=1}^3 a_i + b_i = \omega_1 + \omega_2$  and  $\sum_{i=1}^3 u_i + v_i = 0$ . The  $\epsilon(y)$  function is defined as  $\epsilon(0) = \epsilon(\lfloor \frac{r}{2} \rfloor) = 1$  and  $\epsilon(y) = 2$  otherwise. The main difference from the expression (3.7) is that here the summation is over the holonomies  $y = \frac{r}{2\pi} \int A_\mu dx^\mu$ , where the integration is over a non-trivial cycle on  $S_b^3/\mathbb{Z}_r$  and  $A_\mu$  is the gauge field, see, e.g. [31].

By introducing new variables  $a_i = -\alpha_i + x_i$  and  $b_i = -\alpha_i - x_i$  with the condition  $u_i = -v_i$  for  $i = 1, 3$ , one can rewrite the integral identity (3.12) as the star-triangle equation with the following Boltzmann weight

$$\begin{aligned} & W_\alpha(x_i, x_j, u_i, u_j) \\ &= e^{-\pi i(u_i + u_j)} \gamma^{(2)}(-i(-\alpha + x_i - x_j) \\ & \quad - i\omega_1(u_i - u_j); -i\omega_1 r, -i\omega) \\ & \times \gamma^{(2)}(-i(-\alpha + x_i - x_j) \\ & \quad - i\omega_2(r - (u_i - u_j)); -i\omega_2 r, -i\omega) \\ & \times \gamma^{(2)}(-i(-\alpha - x_i + x_j) \\ & \quad - i\omega_1(u_j - u_i); -i\omega_1 r, -i\omega) \\ & \times \gamma^{(2)}(-i(-\alpha - x_i + x_j) \\ & \quad - i\omega_2(r - (u_j - u_i)); -i\omega_2 r, -i\omega). \end{aligned} \quad (3.13)$$

The model with the Boltzmann weight (3.13) is an exactly solvable lattice spin model with discrete and continuous spin variables living on sites, where  $x_i$  represents continuous spin and  $u_i$  represents discrete spin variable. The  $r = 1$  case corresponds to the Faddeev–Volkov model from the previous section<sup>5</sup>.

Using the similar technique presented in Appendix A one can construct the star-star relation for this model

$$\begin{aligned} & \sum_{y=0}^{[r/2]} \epsilon(y) \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(-i(a_i - x) - i\omega_1(u_i - y); -i\omega_1 r, -i\omega) \\ & \times \gamma^{(2)}(-i(a_i - x) - i\omega_2(r - (u_i - y)); -i\omega_2 r, -i\omega) \\ & \times \gamma^{(2)}(-i(b_i + x) - i\omega_1(v_i + y); -i\omega_1 r, -i\omega) \\ & \times \gamma^{(2)}(-i(b_i + x) - i\omega_2(r - (v_i + y)); -i\omega_2 r, -i\omega) \frac{dx}{r\sqrt{-\omega_1\omega_2}} \\ &= \frac{e^{\frac{\pi i}{2} \sum_{i=1}^2 (u_i - v_i)}}{e^{\frac{\pi i}{2} \sum_{i=3}^4 (\tilde{u}_i - \tilde{v}_i)}} \\ & \times \frac{\prod_{i,j=1}^2 \gamma^{(2)}(-i(a_i + b_j) - i\omega_1(u_i + v_j); -i\omega_1 r, -i\omega)}{\prod_{i,j=3}^4 \gamma^{(2)}(-i(\tilde{a}_i + \tilde{b}_j) - i\omega_1(\tilde{u}_i + \tilde{v}_j); -i\omega_1 r, -i\omega)} \\ & \times \frac{\prod_{i,j=1}^2 \gamma^{(2)}(-i(a_i + b_j) - i\omega_2(r - (u_i + v_j)); -i\omega_2 r, -i\omega)}{\prod_{i,j=3}^4 \gamma^{(2)}(-i(\tilde{a}_i + \tilde{b}_j) - i\omega_2(r - (\tilde{u}_i + \tilde{v}_j)); -i\omega_2 r, -i\omega)} \end{aligned}$$

<sup>5</sup> The  $r = 2$  case is also special, in [40] it was used for proving orthogonality and completeness relation of the Clebsch–Gordan coefficients for the self-dual continuous series of  $U_q(osp(1|2))$ .

$$\begin{aligned}
& \times \sum_{m=0}^{\lfloor r/2 \rfloor} \epsilon(m) \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(-i(\tilde{a}_i - z) - i\omega_1(\tilde{u}_i - m); -i\omega_1 r, -i\omega) \\
& \times \gamma^{(2)}(-i(\tilde{a}_i - z) - i\omega_2(r - (\tilde{u}_i - m)); -i\omega_2 r, -i\omega) \\
& \times \gamma^{(2)}(-i(\tilde{b}_i + z) - i\omega_1(\tilde{v}_i + m); -i\omega_1 r, -i\omega) \\
& \times \gamma^{(2)}(-i(\tilde{b}_i + z) - i\omega_2(r - (\tilde{v}_i + m)); -i\omega_2 r, -i\omega) \\
& \times \frac{dz}{r\sqrt{-\omega_1\omega_2}}, \quad (3.14)
\end{aligned}$$

where the balancing conditions are  $\sum_{i=1}^4 a_i + b_i = 2(\omega_1 + \omega_2)$  and  $\sum_{i=1}^4 u_i + v_i = 0$ , and we used the following notations

$$\begin{aligned}
\tilde{a}_i &= a_i + s, \tilde{b}_i = b_i + s, \tilde{u}_i = u_i + p, \\
\tilde{v}_i &= v_i + p, \text{ if } i = 1, 2, \\
\tilde{a}_i &= a_i - s, \tilde{b}_i = b_i - s, \tilde{u}_i = u_i - p, \\
\tilde{v}_i &= v_i - p, \text{ if } i = 3, 4,
\end{aligned} \quad (3.15)$$

where

$$\begin{aligned}
s &= \frac{1}{2}(\omega_1 + \omega_2 - a_1 - a_2 - b_1 - b_2) \\
&= \frac{1}{2}(-\omega_1 - \omega_2 + a_3 + a_4 + b_3 + b_4), \\
p &= -\frac{1}{2}(u_1 + u_2 + v_1 + v_2) = \frac{1}{2}(u_3 + u_4 + v_3 + v_4).
\end{aligned} \quad (3.16)$$

### 3.4 Solution via $S^2 \times S^1$ supersymmetric partition function

In this section we present a new trigonometric solution to the star-triangle equation and to the star-star equation. In

$$\begin{aligned}
& \sum_{y=-\infty}^{\infty} \oint \prod_{i=1}^3 \frac{(q^{1+(y+u_i)/2}(a_i x)^{-1}, q^{1+(v_i-y)/2} x b_i^{-1}; q)_{\infty}}{(q^{(y+u_i)/2} a_i x, q^{(v_i-y)/2} b_i x^{-1}; q)_{\infty}} \frac{1}{x^{3y}} \frac{dx}{2\pi i x} \\
&= \frac{1}{\prod_{i=1}^3 a_i^{u_i} b_i^{v_i}} \prod_{i,j=1}^3 \frac{(q^{1+(u_i+v_j)/2}(a_i b_j)^{-1}; q)_{\infty}}{(q^{(u_i+v_j)/2} a_i b_j; q)_{\infty}}, \quad (3.17)
\end{aligned}$$

where the balancing conditions are  $\prod_{i=1}^3 a_i b_i = q$  and  $\sum_{i=1}^3 u_i + v_i = 0$ .

This identity can be written as the star-triangle relation by introducing the new fugacities  $a_i = \alpha_i^{-1} x_i$  and  $b_i = \alpha_i^{-1} x_i^{-1}$  and using the condition  $u_i = -v_i$ . The resulting Boltzmann weight then has the following form

$$\begin{aligned}
W_{\alpha}(x_i, x_j, u_i, u_j) &= \frac{(q^{1+(u_i-u_j)/2}(\alpha^{-1} x_i x_j^{-1})^{-1}; q)_{\infty}}{(q^{(u_i-u_j)/2} \alpha^{-1} x_i x_j^{-1}; q)_{\infty}} \\
&\times \frac{(q^{1+(u_j-u_i)/2}(\alpha^{-1} x_i^{-1} x_j)^{-1}; q)_{\infty}}{(q^{(u_j-u_i)/2} \alpha^{-1} x_i^{-1} x_j; q)_{\infty}}, \quad (3.18)
\end{aligned}$$

where  $\alpha$  stands for the spectral parameter. The corresponding statistical mechanics model is a square lattice model with edge interaction and discrete and continuous spin variables. This model is a special case (with broken gauge symmetry) of the integrable lattice spin model considered in [3] and it is a trigonometric analogue of the Faddeev–Volkov model.

One we can construct the star-star relation for this model

$$\begin{aligned}
& \sum_{y=-\infty}^{\infty} \oint \prod_{i=1}^4 \frac{(q^{1+(y+u_i)/2}(a_i x)^{-1}, q^{1+(v_i-y)/2} x(b_i)^{-1}; q)_{\infty}}{(q^{(y+u_i)/2} a_i x, q^{(v_i-y)/2} b_i x^{-1}; q)_{\infty}} \frac{1}{x^{5y}} \frac{dx}{2\pi i x} = \frac{1}{\prod_{i=1}^2 a_i^{u_i} b_i^{v_i}} \frac{\prod_{i,j=1}^2 \frac{(q^{1+(u_i+v_j)/2}(a_i b_j)^{-1}; q)_{\infty}}{(q^{(u_i+v_j)/2} a_i b_j; q)_{\infty}}}{\prod_{i=3}^4 (\tilde{a}_i)^{u_i} (\tilde{b}_i)^{v_i}} \frac{\prod_{i,j=3}^4 \frac{(q^{1+(\tilde{u}_i+\tilde{v}_j)/2}(\tilde{a}_i \tilde{b}_j)^{-1}; q)_{\infty}}{(q^{(\tilde{u}_i+\tilde{v}_j)/2} \tilde{a}_i \tilde{b}_j; q)_{\infty}}}{\prod_{i=3}^4 (\tilde{a}_i)^{u_i} (\tilde{b}_i)^{v_i}} \\
& \times \sum_{m=-\infty}^{\infty} \oint \prod_{i=1}^4 \frac{(q^{1+(m+\tilde{u}_i)/2}(\tilde{a}_i z)^{-1}, q^{1+(\tilde{v}_i-m)/2} z \tilde{b}_i^{-1}; q)_{\infty}}{(q^{(m+\tilde{u}_i)/2} \tilde{a}_i z, q^{(\tilde{v}_i-m)/2} \tilde{b}_i z^{-1}; q)_{\infty}} \frac{1}{z^{5m}} \frac{dz}{2\pi i z}, \quad (3.19)
\end{aligned}$$

[22,23,41] authors considered the following basic hypergeometric sum/integral identity<sup>6</sup> which represents the equivalence of the superconformal indices for dual theories (partition function on  $S^2 \times S^1$ ) discussed in Sect. 3

with the new balancing conditions  $\prod_{i=1}^4 a_i b_i = q^2$  and  $\sum_{i=1}^4 u_i + v_i = 0$ . In the latter expression we used the following notations

$$\begin{aligned}
\tilde{a}_i &= a_i s, \tilde{b}_i = b_i s, \tilde{u}_i = u_i + p, \\
\tilde{v}_i &= v_i + p, \text{ if } i = 1, 2, \\
\tilde{a}_i &= a_i s^{-1}, \tilde{b}_i = b_i s^{-1}, \tilde{u}_i = u_i - p, \\
\tilde{v}_i &= v_i - p, \text{ if } i = 3, 4,
\end{aligned} \quad (3.20)$$

<sup>6</sup> This identity can be written as a pentagon identity which is related to the triangulation of 3-manifolds.

where

$$s = \sqrt{\frac{q}{a_1 a_2 b_1 b_2}} = \sqrt{\frac{a_3 a_4 b_3 b_4}{q}},$$

$$p = -\frac{1}{2}(u_1 + u_2 + v_1 + v_2) = \frac{1}{2}(u_3 + u_4 + v_3 + v_4). \quad (3.21)$$

## 4 Conclusions

In this work, we constructed hyperbolic and trigonometric solutions to the star-star equation. We obtained new solutions from the equality of three-dimensional  $\mathcal{N} = 2$  supersymmetric partition functions for certain dual SQED theories via the gauge/YBE correspondence.

There are several ways of constructing solutions to the star-star equation. One can use the Bailey pair construction starting from the star-triangle relation for the models discussed here. It is possible to obtain the solution by breaking the gauge symmetry from  $SU(2)$  group to the  $U(1)$  for the supersymmetric dual theories with  $SU(2)$  gauge group and  $SU(6)$  flavor group considered in [42].

There are many interesting limits of the solutions considered here, for instance it would be interesting to construct solution to the star-star equation in terms of Euler gamma functions [43].

The gauge/YBE correspondence has revealed various interesting connections among integrable models and supersymmetric gauge theories. There are underlying mathematical structures such as quantum algebras related to the solutions of the Yang–Baxter equations obtained via gauge/YBE correspondence. It would be interesting to pursue this direction for these solutions to the star-star relation.

**Acknowledgements** We are grateful to Deniz N. Bozkurt for valuable discussions. The work of Ilmar Gahramanov is partially supported by the Bogazici University Research Fund under Grant Number 20B03SUP3 and TUBITAK Grant 220N106. Mustafa Mullahasanoglu is supported by the 2209-TUBITAK National/International Research Projects Fellowship Programme for Undergraduate Students under Grant Number 1919B011902237.

**Data Availability Statement** This manuscript has no associated data or the data will not be deposited. [Authors' comment: Our manuscript has no associated data or the data will not be deposited.]

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permit-

ted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP<sup>3</sup>. SCOAP<sup>3</sup> supports the goals of the International Year of Basic Sciences for Sustainable Development.

## Appendix A: Derivation of the star-star relation (3.9)

Here we follow the approach presented in [38].

Let us consider the following double integral

$$\int_{-\infty}^{\infty} \prod_{i=1}^2 \gamma^{(2)}(a_i - x) \gamma^{(2)}(b_i + x) \\ \times \int_{-\infty}^{\infty} \prod_{i=3}^4 \gamma^{(2)}(a_i - s - z) \gamma^{(2)}(b_i - s + z) \\ \times \gamma^{(2)}(s + z - x) \gamma^{(2)}(s - z + x) \frac{dx}{i\sqrt{\omega_1 \omega_2}} \frac{dz}{i\sqrt{\omega_1 \omega_2}}. \quad (A.1)$$

Here we used the shorthand notation  $\gamma^{(2)}(x) = \gamma^{(2)}(x; \omega_1, \omega_2)$ . First we integrate the integral (A.1) over the  $x$  variable using the identity (3.7). We end up with the following result

$$\gamma^{(2)}(2s) \prod_{i,j=1}^2 \gamma^{(2)}(a_i + b_j) \\ \times \int_{-\infty}^{\infty} \prod_{i=3}^4 \gamma^{(2)}(a_i - s - z) \gamma^{(2)}(b_i - s + z) \\ \times \prod_{i=1}^2 \gamma^{(2)}(s + z + b_i) \prod_{i=1}^2 \gamma^{(2)}(a_i + s - z) \frac{dz}{i\sqrt{\omega_1 \omega_2}}. \quad (A.2)$$

Then integrating (A.1) over the  $z$  variable one finds that

$$\gamma^{(2)}(2s) \prod_{i,j=3}^4 \gamma^{(2)}(a_i + b_j - 2s) \\ \times \int_{-\infty}^{\infty} \prod_{i=1}^2 \gamma^{(2)}(a_i - x) \gamma^{(2)}(b_i + x) \\ \times \prod_{i=3}^4 \gamma^{(2)}(b_i + x) \prod_{i=3}^4 \gamma^{(2)}(a_i - x) \frac{dx}{i\sqrt{\omega_1 \omega_2}}. \quad (A.3)$$

The latter two expressions are results of the same integral expression, therefore we find that

$$\gamma^{(2)}(2s) \prod_{i,j=1}^2 \gamma^{(2)}(a_i + b_j)$$



$$\begin{aligned}
& \times \int_{-\infty}^{\infty} \prod_{i=3}^4 \gamma^{(2)}(a_i - s - z) \gamma^{(2)}(b_i - s + z) \\
& \times \prod_{i=1}^2 \gamma^{(2)}(s + z + b_i) \prod_{i=1}^2 \gamma^{(2)}(a_i + s - z) \frac{dz}{i\sqrt{\omega_1 \omega_2}} \\
& = \gamma^{(2)}(2s) \prod_{i,j=3}^4 \gamma^{(2)}(a_i + b_j - 2s) \\
& \times \int_{-\infty}^{\infty} \prod_{i=1}^2 \gamma^{(2)}(a_i - x) \gamma^{(2)}(b_i + x) \\
& \times \prod_{i=3}^4 \gamma^{(2)}(b_i + x) \prod_{i=3}^4 \gamma^{(2)}(a_i - x) \frac{dx}{i\sqrt{\omega_1 \omega_2}}. \quad (\text{A.4})
\end{aligned}$$

One can rewrite this expression in a compact form

$$\begin{aligned}
& \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(a_i - x) \gamma^{(2)}(b_i + x) \frac{dx}{i\sqrt{\omega_1 \omega_2}} \\
& = \frac{\prod_{i,j=1}^2 \gamma^{(2)}(a_i + b_j)}{\prod_{i,j=3}^4 \gamma^{(2)}(\tilde{a}_i + \tilde{b}_j)} \\
& \times \int_{-\infty}^{\infty} \prod_{i=1}^4 \gamma^{(2)}(\tilde{a}_i - z) \gamma^{(2)}(\tilde{b}_i + z) \frac{dz}{i\sqrt{\omega_1 \omega_2}}, \quad (\text{A.5})
\end{aligned}$$

with the related balancing condition  $\sum_{i=1}^4 a_i + b_i = 2(\omega_1 + \omega_2)$  by introducing new variables

$$\begin{aligned}
\tilde{a}_i &= a_i + s, & \tilde{b}_i &= b_i + s, & \text{if } i &= 1, 2, \\
\tilde{a}_i &= a_i - s, & \tilde{b}_i &= b_i - s, & \text{if } i &= 3, 4,
\end{aligned} \quad (\text{A.6})$$

where

$$\begin{aligned}
s &= \frac{1}{2}(\omega_1 + \omega_2 - a_1 - a_2 - b_1 - b_2) \\
&= \frac{1}{2}(-\omega_1 - \omega_2 + a_3 + a_4 + b_3 + b_4).
\end{aligned} \quad (\text{A.7})$$

In order to obtain the final result one needs to use the reflection property (3.6) for the hyperbolic gamma function.

## References

- V.P. Spiridonov, Elliptic beta integrals and solvable models of statistical mechanics. *Contemp. Math.* **563**, 181–211 (2012). [arXiv:1011.3798](#) [hep-th]
- M. Yamazaki, Quivers, YBE and 3-manifolds. *JHEP* **05**, 147 (2012). [arXiv:1203.5784](#) [hep-th]
- I. Gahramanov, V.P. Spiridonov, The star-triangle relation and 3d superconformal indices. *JHEP* **08**, 040 (2015). [arXiv:1505.00765](#) [hep-th]
- A.P. Kels, New solutions of the star-triangle relation with discrete and continuous spin variables. *J. Phys. A* **48**(43), 435201 (2015). [arXiv:1504.07074](#) [math-ph]
- A.P. Kels, M. Yamazaki, Lens elliptic gamma function solution of the Yang–Baxter equation at roots of unity. *J. Stat. Mech.* **1802**(2), 023108 (2018). [arXiv:1709.07148](#) [math-ph]
- M. Yamazaki, New integrable models from the gauge/YBE correspondence. *J. Stat. Phys.* **154**, 895 (2014). [arXiv:1307.1128](#) [hep-th]
- M. Yamazaki, W. Yan, Integrability from 2d  $\mathcal{N} = (2, 2)$  dualities. *J. Phys. A* **48**, 394001 (2015). [arXiv:1504.05540](#) [hep-th]
- I. Gahramanov, S. Jafarzade, Comments on the multi-spin solution to the Yang–Baxter equation and basic hypergeometric sum/integral identity. *Mod. Phys. Lett. A* **34**(18), 1950140 (2019). [arXiv:1710.09106](#) [math-ph]
- I. Gahramanov, S. Jafarzade, Integrable lattice spin models from supersymmetric dualities. *Phys. Part. Nucl. Lett.* **15**(6), 650–667 (2018). [arXiv:1712.09651](#) [math-ph]
- M. Yamazaki, Integrability as duality: the gauge/YBE correspondence. [arXiv:1808.04374](#) [hep-th]
- I. Gahramanov, Mathematical structures behind supersymmetric dualities. *Arch. Math.* **51**, 273–286 (2015). [arXiv:1505.05656](#) [math-ph]
- V.P. Spiridonov, Superconformal indices, Seiberg dualities and special functions. *Phys. Part. Nucl.* **51**(4), 508–513 (2020). [arXiv:1912.11514](#) [hep-th]
- Y. Tachikawa, On ‘categories’ of quantum field theories, in *International congress of mathematicians*, p. 12 (2017). [arXiv:1712.09456](#) [math-ph]
- V.P. Spiridonov, G.S. Vartanov, Elliptic hypergeometry of supersymmetric dualities. *Commun. Math. Phys.* **304**, 797–874 (2011). [arXiv:0910.5944](#) [hep-th]
- V.P. Spiridonov, G.S. Vartanov, Elliptic hypergeometry of supersymmetric dualities II. Orthogonal groups, knots, and vortices. *Commun. Math. Phys.* **325**(2), 421–486 (2014). [arXiv:1107.5788](#) [hep-th]
- C. Krattenthaler, V.P. Spiridonov, G.S. Vartanov, Superconformal indices of three-dimensional theories related by mirror symmetry. *JHEP* **06**, 008 (2011). [arXiv:1103.4075](#) [hep-th]
- I.B. Gahramanov, G.S. Vartanov, Superconformal indices and partition functions for supersymmetric field theories, in *XVIIIth Intern. Cong. Math. Phys.*, pp. 695–703 (2013). [arXiv:1310.8507](#) [hep-th]
- F.A.H. Dolan, V.P. Spiridonov, G.S. Vartanov, From 4d superconformal indices to 3d partition functions. *Phys. Lett. B* **704**, 234–241 (2011). [arXiv:1104.1787](#) [hep-th]
- D.N. Bozkurt, I. Gahramanov, M. Mullahasanoglu, Lens partition function, pentagon identity, and star-triangle relation. *Phys. Rev. D* **103**(12), 126013 (2021). [arXiv:2009.14198](#) [hep-th]
- R.J. Baxter, Star-triangle and star-star relations in statistical mechanics. *Int. J. Mod. Phys. B* **11**, 27–37 (1997)
- V.V. Bazhanov, V.V. Mangazeev, S.M. Sergeev, Exact solution of the Faddeev–Volkov model. *Phys. Lett. A* **372**, 1547–1550 (2008). [arXiv:0706.3077](#) [cond-mat.stat-mech]
- I. Gahramanov, H. Rosengren, A new pentagon identity for the tetrahedron index. *JHEP* **11**, 128 (2013). [arXiv:1309.2195](#) [hep-th]
- I. Gahramanov, H. Rosengren, Integral pentagon relations for 3d superconformal indices. [arXiv:1412.2926](#) [hep-th]. [*Proc. Symp. Pure Math.* **93**, 165 (2016)]
- R. Baxter, Fundamental problems in statistical mechanics v, in *Proceedings of the 1980 Enschede Summer School* (1980)
- R.J. Baxter, Exactly solved models in statistical mechanics (Elsevier, 2016)
- R.J. Baxter, Star-triangle and star-star relations in statistical mechanics. *Int. J. Mod. Phys. B* **11**(01n02), 27–37 (1997)

27. K. Hosomichi, A review on SUSY gauge theories on  $S^3$ . [arXiv:1412.7128](#) [hep-th]
28. B. Willett, Localization on three-dimensional manifolds. *J. Phys. A* **50**(44), 443006 (2017). [arXiv:1608.02958](#) [hep-th]
29. V. Pestun, Localization of gauge theory on a four-sphere and supersymmetric Wilson loops. *Commun. Math. Phys.* **313**, 71–129 (2012). [arXiv:0712.2824](#) [hep-th]
30. A. Kapustin, B. Willett, I. Yaakov, Nonperturbative tests of three-dimensional dualities. *JHEP* **10**, 013 (2010). [arXiv:1003.5694](#) [hep-th]
31. Y. Imamura, D. Yokoyama,  $S^3/Z_n$  partition function and dualities. *JHEP* **11**, 122 (2012). [arXiv:1208.1404](#) [hep-th]
32. Y. Imamura, S. Yokoyama, Index for three dimensional superconformal field theories with general R-charge assignments. *JHEP* **1104**, 007 (2011). [arXiv:1101.0557](#) [hep-th]
33. L. Faddeev, Discrete Heisenberg–Weyl group and modular group. *Lett. Math. Phys.* **34**, 249–254 (1995). [arXiv:hep-th/9504111](#)
34. S. Woronowicz, Quantum exponential function. *Rev. Math. Phys.* **12**(06), 873–920 (2000)
35. R. Kashaev, F. Luo, G. Vartanov, A TQFT of Turaev–Viro type on shaped triangulations. *Ann. Henri Poincaré* **17**(5), 1109–1143 (2016). [arXiv:1210.8393](#) [math.QA]
36. F.J. van de Bult, Hyperbolic hypergeometric functions. PhD thesis, University of Amsterdam (2007)
37. V.V. Bazhanov, V.V. Mangazeev, S.M. Sergeev, Faddeev–Volkov solution of the Yang–Baxter equation and discrete conformal symmetry. *Nucl. Phys. B* **784**, 234–258 (2007). [arXiv:hep-th/0703041](#)
38. F.J. van de Bult, E.M. Rains, J.V. Stokman, Properties of generalized univariate hypergeometric functions. *Commun. Math. Phys.* **275**(1), 37–95 (2007)
39. T. Dimofte, D. Gaiotto, An E7 surprise. *JHEP* **10**, 129 (2012). [arXiv:1209.1404](#) [hep-th]
40. L. Hadasz, M. Pawelkiewicz, V. Schomerus, Self-dual continuous series of representations for  $\mathcal{U}_q(sl(2))$  and  $\mathcal{U}_q(osp(1|2))$ . *JHEP* **10**, 091 (2014). [arXiv:1305.4596](#) [hep-th]
41. I. Gahramanov, H. Rosengren, Basic hypergeometry of supersymmetric dualities. *Nucl. Phys. B* **913**, 747–768 (2016). [arXiv:1606.08185](#) [hep-th]
42. I. Gahramanov, A.P. Kels, The star-triangle relation, lens partition function, and hypergeometric sum/integrals. *JHEP* **02**, 040 (2017). [arXiv:1610.09229](#) [math-ph]
43. E. Eren, I. Gahramanov, S. Jafarzade, G. Mogol, Gamma function solutions to the star-triangle equation. *Nucl. Phys. B* **963**, 115283 (2021). [arXiv:1912.12271](#) [math-ph]