Erratum

Erratum to: Numerical solutions of thin-film equations for polymer flows

Thomas Salez^{1,a}, Joshua D. McGraw², Sara L. Cormier², Oliver Bäumchen², Kari Dalnoki-Veress², and Elie Raphaël¹

¹ Laboratoire de Physico-Chimie Théorique, UMR CNRS Gulliver 7083, ESPCI, Paris, France

 $^2\,$ Department of Physics & Astronomy and the Brockhouse Institute for Materials Research, McMaster University, Hamilton, Canada

Original article: Eur. Phys. J. E (2012) 35: 114, DOI: 10.1140/epje/i2012-12114-x

Received 22 August 2013 Published online: 30 September 2013 – © EDP Sciences / Società Italiana di Fisica / Springer-Verlag 2013

Equation (36h) must be replaced by

$$\partial_T \mathcal{H}_3 = rac{8}{3} \; rac{\mathcal{A}_3 \mathcal{H}_{\bar{R}R\bar{R},3} - \mathcal{A}_4 \mathcal{H}_{\bar{R}R\bar{R},4}}{\Delta R}$$

which corresponds to the limit of eq. (28) when $R \to 0$.

Note that the 4/3 factor between the two expressions applies on a single point of the grid, the others being described by the full eq. (33). We have verified that this error is insignificant up to a typical 10^{-4} relative error on the output profiles, with the chosen spatial increment ($\Delta R = 0.01$) reported in table 2. The results and figures of the article are thus unchanged.

However, the missing 4/3 factor may lead to larger discrepancies for a coarser grid.

We take advantage of this *Erratum* to mention also that the conditions of eqs. (36a), (36d) and (36e) are not needed in the algorithm.

We thank Matilda Backholm and Michael Benzaquen for a recent independent study that contributed to raise this issue.

^a e-mail: thomas.salez@espci.fr