

RESEARCH ARTICLE

Open Access

Obesity and overweight in Bangladeshi children and adolescents: a scoping review

Sholeh Rahman[†], Md Tauhidul Islam[†] and Dewan S Alam^{*}

Abstract

Background: Obesity and overweight in children and adolescents is an emerging public health concern alongside under-nutrition in low and middle income countries. Our aim was to conduct a scoping review of literature to ascertain what is known about childhood and adolescent overweight and/or obesity in Bangladesh.

Method: Using the scoping review based on York methodology, a comprehensive search of published academic articles, conference proceedings and grey literature was carried out through PubMed, BanglajOL, Google and Google scholar limited to English-written papers. We summarized prevalence, risk factors and health outcomes of obesity/overweight in young children and adolescents aged between 0 to 19 years old in Bangladesh and highlighted use of different reference standards to measure childhood obesity.

Results: In total 21 studies met the inclusion criteria. Nearly all of the reviewed articles used data from cross sectional studies, while only two used case-control design. Overall thirteen studies (62%) were primary research and eight (38%) included secondary data. Studies indicated an increasing trend in childhood obesity over time. Prevalence ranged from less than 1% to 17.9% based on different reference standards, with higher percentage amongst urban children across different age groups and sexes.

Conclusion: This review demonstrated paucity of comprehensive literature on childhood obesity in Bangladesh, which might be explored through population-based prospective studies based on strong methodology and uniform reference standards. Sustainable and scalable preventative measures targeting high risk groups are required to avoid further rise.

Keywords: Overweight, Obesity, Children, Adolescents, Prevalence, Bangladesh

Background

Obesity in children and adolescents is rising alarmingly and approaching epidemic proportion in many economically developed countries, particularly in USA, Canada, Australia and several European countries [1]. Likewise in developing countries this issue is emerging as a public health crisis. According to a recent report, out of an estimated 43 million obese children worldwide in 2010, approximately 81% were from developing countries, half of which (18 million) were reported to be living in Asia despite of huge burden of under-nutrition. By 2020, it is estimated that the global prevalence of childhood obesity will reach approximately 60 million [2].

Factors contributing to the rising levels of childhood obesity in developing countries include socio-economic development, changes in lifestyle characterized by physical inactivity and unhealthy diet, living patterns, as well as rapid epidemiological and demographic transition [3,4].

WHO (2013) considers childhood obesity as “one of the most serious public health challenges of 21st century” [5]. It accounts for a wide range of psychosocial and medical consequences [6]. Lower self-esteem, social isolation, poor academic achievement and peer problems are the most apparent immediate consequences in obese children [7]. Obesity in children and adolescents predisposes them to and is causally linked with cardio-metabolic disorders such as hypertension, dyslipidaemia and insulin resistance which are well-established illnesses [8]. As overweight and obesity is likely to follow through into adulthood, in the longer term, there is a greater risk of developing

* Correspondence: dsalam@icddr.org

[†]Equal contributors

Centre for Control of Chronic Diseases (CCCD), icddr,b, 68 Shaheed Tajuddin Ahmed Sharani, Mohakhali Dhaka 1212, Bangladesh

cardiovascular diseases [9,10]. Obesity at a young age seems to have substantial impact on reducing life expectancy [11]. Evidence also suggests a link between obesity in young girls with potential menstrual disorders, hypertension in pregnancy and sub-fertility [12].

In Bangladesh, the context of obesity and overweight has been underexplored, more so amongst younger age groups. Understanding the current situation and trends will provide useful insights into its risk factors and will assist health professionals and policy-makers in decision-making and developing future research agenda. This paper explored the availability of literature on childhood and adolescent obesity/overweight and its determinants in Bangladesh through a scoping review which, unlike a systematic review, offers a much broader perspective in the respective field which makes it more appropriate method to assess the present situation of childhood and adolescent obesity in Bangladesh [13].

Methods

A scoping review was performed based on the York methodology outlined by Arksey and O' Malley from the University of York, United Kingdom [13]. The 'York framework' suggested five stages that we have followed for this review:

- Stage 1: Identifying the research question
- Stage 2: Identifying the relevant studies
- Stage 3: Study selection
- Stage 4: Charting the data
- Stage 5: Collating, summarizing and reporting the results

Initially, we defined research questions and developed search strategy and discussed them in a team meeting. Relevant literatures were then identified through a comprehensive search across different databases including Pubmed, Google, Google scholar and Bangladesh's country specific search engine (BanglaJOL). We used specific key words that included obesity, overweight, adiposity, BMI, childhood, adolescents and Bangladesh. Search terms were combined using Boolean operators (AND, OR, NOT) to narrow the results [14]. In addition, hand searching of appropriate reference lists were undertaken to identify any additional literatures or grey publications. Figure 1 shows different steps of the review conducted.

Selection of literature

We included the publication if:

- obesity and/or overweight was its primary or secondary outcome or an associated risk factor;
- study participants were Bangladeshi children and adolescents aged between 0 to 19 year old;




Figure 1 Steps in performing the scoping review on childhood obesity in Bangladesh. After determining the research question and search strategy, the authors conducted literature search using specific keywords followed by review of the selected literature to generate the results.

- it was relevant to any aspect of obesity, such as its prevalence, determinants, nutrition and growth in children and adolescents, anthropometry and health status; and
- it was written in English; no other limits on publication date of the articles were considered.

Papers were excluded if the study sample was older than 19 years of age or resided in countries other than Bangladesh. The initial search of databases revealed 241 literatures of which 159 were excluded after screening their titles and abstracts. One paper was not available even after contacting the library services and the publisher. The remaining 82 papers were screened for relevancy of their objectives, out of which a further 39 papers were excluded due to duplication or overlapping. Full copies of 43 literatures were read first independently and then reviewed again by the authors, which resulted in further exclusion of 35 papers as they did not meet our inclusion criteria. We retrieved 14 papers through manual search, from which one was excluded because of its apparently unrealistic results as well as lack of access to the full article [15]. The final number of papers

included in this review narrowed down to 21, constituting of 19 full papers, 1 journal abstract and 1 conference proceeding [16-36] (Figure 2). Details of each of the included papers were then summarized (Table 1).

Results

Application of inclusion and exclusion criteria resulted in overall selection of 21 papers for this review; searching BanglaJOL, PubMed, Google and Google Scholar revealed 1, 7 and 13 relevant papers respectively. Literatures selected were from a 15 year period with the oldest from 1998 to the most recent in 2013. The search did not retrieve any relevant literature beyond these dates.

Characteristics of the selected data

Seventy one percent of the studies (n = 15) included participants from both sexes and 29% involved females only (n = 6). Age range of the participants varied considerably across different studies. To make the interpretation easier,

we decided to divide the studies into three categories according to their participants' age range: group A (included those 5 years of age or younger), group B (included those over 5 years of age), group C (mixed group included subjects younger and older than 5 years of age). Accordingly twelve studies that examined obesity fell in group B, five in group A and only four in group C which included both age categories. In terms of the study site, majority of the studies (n = 14) were conducted in urban setting, six studies examined obesity in both urban and rural, and only one paper reported obesity in rural setting.

Study design, measurement and reference standards

Sixty two percent (n = 13) of the studies were primary research and thirty eight percent (n = 8) used secondary data for their result synthesis. Bangladesh Demographic and Health Survey (BDHS 2004 and 2007), Nutrition Surveillance Project (NSP 2000–2004), WHO database (1996) and Diarrheal Disease Surveillance System (DDSS




Figure 2 Flowchart of the number of literature searched and selected. Sequential steps undertaken for final inclusion of the literature in our review.

Table 1 Studies of childhood and adolescent obesity included in the review by study design, sample size, anthropometric indices and reference standards

Author(s) and year	n	Study design	Anthropometric index	Reference standard
Das et al., 2013 [16]	33,482	Cross-sectional	Weight for age (under 5) & BMI for age (5-19 years)	WHO, 2009 ¹ National Obesity Observatory, 2011 ²
Bhuiyan, 2013 [17]	198	Case-Control	BMI	CDC, 2000 ³
Hossain et al., 2012 [18]	10,115*	Cross-sectional	BMI	WHO, 2006 ⁴
Mohsin et al., 2012 [19]	161	Cross-sectional	BMI	CDC, 2000 ³
Saha et al., 2011-2012 [20]	96	Cross-sectional	Not defined	N/A
Corsi et al., 2011 [21]	1439	Cross-sectional	BMI	WHO, 2004 ⁵
Jesmin et al., 2011 [22]	380	Cross-sectional	Weight for height	WHO, 2006 ⁴
Mohsin et al., 2011 [23]	161	Cross-sectional	BMI	CDC, 2000 ³
Mohsin et al., 2010 [24]	468	Cross-sectional	BMI	CDC, 2000 ³
Sultana, 2010 [25]	1200	Cross-sectional	Weight for height	WHO, 1995 ⁶
Kurshed et al., 2010 [26]	352	Cross-sectional	BMI	WHO, 1995 ⁶
Rahman et al., 2009 [27]	652	Cross-sectional	BMI	CDC, 2000 ³
Balarajan et al., 2009 [28]	1328	Cross-sectional	BMI	WHO, 1995 ⁶
Khan et al., 2009 [29]	425	Cross-sectional	BMI	WHO, 2004 ⁵
Sultan et al., 2008 [30]	172	Cross-sectional	BMI	Dietz & Robinson, 1998 ⁷
Shafique et al., 2007 [31]	16404	Cross-sectional	BMI	WHO, 2004 ⁵
BBS/UNICEF, 2007 [32]	3797	Cross-sectional	BMI	WHO, 1995 ⁶
Rahman et al., 2002 [33]	5000	Case control preceded by cross-sectional survey	Weight for Height	Not mentioned
de Onis et al., 2000 [34]	4787	Cross-sectional	Weight for Height	WHO, 1995 ⁶
Rahman et al., 1998 [35]	316	Cross-sectional	Weight for Height	WHO, 1993 ⁸
Ferdousi et al., 2011 [36]	202	Cross-sectional	BMI	WHO, 2004 ⁵

¹WHO Growth Standards for Infants and Young Children.

²A Simple Guide to Classifying Body Mass Index in Children, Public Health Observatories, England.

³2000 CDC Growth Charts for the United States: Methods and development.

⁴WHO Child Growth Standards.

⁵Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies, report of a WHO expert committee.

⁶Physical status: the use and interpretation of anthropometry, report of a WHO expert committee.

⁷Use of body mass index as a measure of overweight in children and adolescents.

⁸Measuring changes in nutritional status.

*Overall sample size included women aged between 15 to 49 years, but sample size for the adolescent group was not mentioned.

1993–2011) were used as the sources of secondary data. The design of the selected studies was mostly cross-sectional, except for two studies which employed case-control design. Amongst the case control studies, one study was preceded by a cross-sectional survey. Different

standards were used for defining overweight and obesity across the studies (Table 1). BMI cut-off value was used as a measure of overweight/obesity in 62% (n = 13) of the studies reviewed, including three studies in under 5 year age group. The remaining 28% (n = 6) adopted

weight for height as an index of body size. Only one study did not mention the body size measurement index [20]. Another study used weight-for-age amongst children who were under 5 years of age and BMI-for-age amongst 5–19 years old [16]. No direct body composition based measure of adiposity was used in any of the reviewed articles. WHO standard for obesity and overweight has been changed and modified over time; however, it was the most commonly used reference for cut-off values in the studies. Few studies used CDC growth chart as their reference standard.

Prevalence of obesity, associated factors and health outcome

Prevalence of overweight and/or obesity was reported in seventeen of the studies and it was the primary outcome in eight of them (47%). The rest of the studies addressed obesity as their secondary outcome. Since the studies did not use the same reference standards, the prevalence of overweight and obesity could not be compared across the studies. However, the reported prevalence varied widely and has been summarized according to the age groups

and presented in Table 2. Recent survey by Bangladesh Bureau of Statistics (BBS) and UNICEF has reported 1.4% prevalence of overweight amongst children less than 5 years of age [32]. A study conducted amongst school children in Dhaka aged between 3 to 18 years of age, found 17.9% obese and 23.6% overweight children and adolescents [24]. In another study, the prevalence of obesity was found to be 13% and that of overweight was 11.5% amongst urban children aged between 2 to 10 years old; and obesity showed an increase with higher family income [35]. Most recent study (2013) among urban children, reported approximately a five fold increase in overweight and obesity over the past two decades [16]. In addition, two unpublished studies reported 17.8% and 7.6% obesity in children of different age categories [25,33]; obesity was significantly more in the highest income group (OR = 2.06) [25].

Only two studies assessed metabolic syndrome in children and adolescents. One of them reported metabolic syndrome in more than 36% of obese children and adolescents with greater number amongst girls [23]. Another one reported 17% IGT, 2% Diabetes Mellitus and 26% high

Table 2 Prevalence (%) of obesity and overweight amongst children and adolescents by sex, age range and setting of the study

	Author(s) and year	Age range	Settings	Sex	Prevalence of overweight/obesity
Group A	Jesmin et al., 2011	0-59 months	Urban	Both	Obesity: 3.9%
	Rahman et al., 2009	24-59 months	Rural	Both	At risk of overweight- male: 0.3%, female: 0.6% Overweight- male: 1.1%, female: 0.6%
	Sultan et al., 2008	1-12 months	Urban	Both	Obesity: 14%; Overweight: 25.6%
	BBS/UNICEF, 2007	0-59 months	Rural and urban	Both	Overweight/Obesity 1.4% (male: 1.2%, female: 1.6%) Severe overweight/obesity: 0.3%
	de Onis et al., 2000	< 5 years	Rural and urban	Both	Overweight: 1.1%
Group B	Hossain et al., 2012	15-19 years ¹	Rural and urban	Females only	Obesity: less than 1%
	Corsi et al., 2011	15-19 years ¹	Rural and urban	Females only	Overweight: 1.6%
	Kurshed et al., 2010	10-18 years	Urban	Females only	Overweight/obesity: 4.6%
	Sultana, 2010	6-13 years	Urban	Both	Overweight: 13.2%; Obesity: 17.8%
	Khan et al., 2009	13-19 years ²	Urban	Females only	Overweight: 2.1%; Obesity: 0.9%
	Balarajan et al., 2009	15-18 years ¹	Urban	Females only	Overweight/obesity: 1.8%
	Shafique et al., 2007	15-18 years ¹	Rural and urban	Females only	Zero prevalence of overweight or obesity
	Ferdousi et al., 2011	6-10 years	Urban	Both	Overweight: 18.32% and higher among males (10.9%)
Group C	Mohsin et al., 2010	3-18 years	Urban	Both	Overweight: 23.6%; Obesity: 17.9%
	Rahman et al., 2002	2-10 years	Urban	Both	Obesity: 7.6%
	Rahman et al., 1998	2-10 years	Urban	Both	Overweight: 11.5%; Obesity: 13%
	Das et al., 2013	< 5 years ³	Urban	Both	Overweight & obesity in < 5: 5.15%
		5-19 years			Overweight & obesity in 5-19 years: 6.7%

Group A: ≤5years of age.

Group B: >5 years of age.

Group C: Mixed group including subjects younger and older than 5 years of age.

¹Age range in overall sample was between 15 to 49 years.

²Age range of overall sample was between 13 to 49 years.

³Overall sample also included > 19 years adults.

total cholesterol among obese children and adolescents [19]. Majority of the studies ($n = 14$) assessed determinants and associated risk factors of obesity. Overall higher socio-economic status, parents higher education (maternal), higher rank occupation, parental obesity, urban residence and physical inactivity were found to be positively associated with overweight and obesity. One study suggested that vitamin A supplements ($OR = 1.665$) and mother's higher education ($OR = 1.98$) were positively associated with healthy, non-obese children [27].

Discussion

Obesity has been studied extensively in many developed countries, but in Bangladesh studies and data related to obesity in children and adolescents are relatively scarce. To our knowledge, this study is the first and only scoping review that has been conducted on the topic in Bangladesh which has employed a comprehensive search of published and un-published literature.

In our review, it was found that the prevalence of overweight and obesity varies significantly across the studies. However, lack of a uniform definition and reference standard makes it difficult to compare the data across studies and predict the trends accurately. We observed greater data scarcity from rural settings than urban, as majority of the studies represented urban children. Although, socio-economic stratification had not been defined in most of the studies, overall greater prevalence of obesity and overweight was reported amongst urban children and adolescents compared to those from rural backgrounds which clearly suggest socio-economic and lifestyle differences in two different settings. In one study overweight and obesity was stated as high as 23.6% and 17.9% respectively amongst affluent urban children [24]. On the other hand, less than 2% of rural children had been reported as being overweight [27]. These findings are in line with similar studies from neighboring India and Thailand which reported greater prevalence of obesity amongst urban children than rural ones [37,38].

Substantial amount of evidence is available on the occurrence of metabolic syndrome in obese children and adolescents. Several studies have reported an association between obesity with a wide range of complications in childhood as well as an increased likelihood of premature onset of chronic illnesses in later life [39-42]. Evidences show simultaneous rise in the prevalence of diabetes and overweight in children, which suggests their close inter-relationship [43-45]. However, little is known in this regard in developing countries, due to limited number of studies. In our review too, we came across very few studies which estimated the prevalence of metabolic syndrome in relation to childhood obesity [19,23], which made it difficult to predict an association between the two. More than half of the studies included in our

review assessed the risk factors of childhood obesity/overweight, irrespective of their design. A positive association between obesity with higher socio-economic status, lack of physical activity and urban residence has been reported. These findings are in accordance with studies from other developing countries which identified the similar risk factors [46,47]. In terms of maternal education, findings were contradictory; one study reported a positive association between child obesity and higher maternal education [16], while another one reported no association [17]. Overall, in our country there is a knowledge gap regarding other risk factors of obesity such as gender, birth weight and socio-cultural factors.

Our review has some limitations. Unlike a systematic review, we did not assess the quality of the studies included in our review, therefore, there is a possibility that the prevalence of overweight and/or obesity might have been exaggerated or underestimated as a result of methodological constraints. The reference standards that have been used in the studies reviewed have not been consistent. Some information might be missing due to inaccessibility of certain databases. In addition, there might have been other unpublished studies that are not available online which could not be revealed. Moreover, information on certain groups, such as tribal or ethnic minorities, is missing. Nevertheless, the findings of this review provide useful insights for future research needs in this area.

Conclusion

This review demonstrated that there is a noticeable published literature gap regarding child and adolescence obesity/overweight in Bangladesh. Although, in general, young age obesity is not considered as a major public health problem in Bangladesh yet, it is alarmingly high and on rise amongst certain groups, particularly amongst urban children from affluent households. Addressing the problem at its earliest could be achieved through identifying high risk groups and designing sustainable interventions which could be implemented on a larger scale to prevent further rise in overweight and obesity. In addition, determining the prevalence of metabolic syndrome amongst them might play a vital role in preventing socio-economic and public health burden in later life.

Implementing school-based programs that could be expanded to population level to promote physical activity and healthy food habits could have a potential impact. Furthermore, large scale prospective studies based on context-specific definition and reference standards of obesity in South-Asian population are required to enhance data comparability and prediction of future trends.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SR, MTI and DSA contributed equally to reviewing the literature, interpreting the results, writing the final report and they are responsible for the final content of this paper. All authors read and approved the final manuscript.

Acknowledgments

This study was supported by icddr. icddr,b acknowledges with gratitude the commitment of the Centre's donors for their generous support to its research efforts. The content of this paper is solely the responsibility of the authors and does not necessarily represent the official view of their employing institution.

Received: 10 September 2013 Accepted: 13 January 2014

Published: 22 January 2014

References

1. Wang Y, Lobstein T: **Worldwide trends in childhood overweight and obesity.** *Pediatr Obes* 2006, **1**(1):11–25. doi:10.1080/17477160600586747.
2. de Onis M, Blossner M, Borghi E: **Global prevalence and trends of overweight and obesity among preschool children.** *Am J Clin Nutr* 2010, **92**(5):1257–1264. doi:10.3945/ajcn.2010.29786.
3. Popkin BM: **An overview on the nutrition transition and its health implications: The Bellagio meeting.** *Public Health Nutr* 2002, **5**(1A):93–103. doi:10.1079/PHN2001280.
4. Davison KK, Birch LL: **Childhood overweight: a contextual model and recommendations for future research.** *Obes Rev* 2001, **2**(3):159–171. doi:10.1046/j.1467-789x.2001.00036.x.
5. World Health Organization: *Childhood overweight and obesity on the rise: Global Strategy on Diet, Physical Activity and Health.* 2004. <http://www.who.int/dietphysicalactivity/childhood/en/>.
6. Lee YS: **Consequences of childhood obesity.** *Ann Acad Med Singapore* 2009, **38**(1):75–77.
7. Dietz WH: **Health consequences of obesity in youth: childhood predictors of adult disease.** *Am Acad Pediatrics* 1998, **101**(Suppl 2):518–525.
8. Reilly JJ: **Descriptive epidemiology and health consequences of childhood obesity.** *Best Pract Res Clin Endocrinol Metab* 2005, **19**(3):327–341. doi:10.1016/j.beem.2005.04.002.
9. Baker JL, Olsen LW, Sorensen TIA: **Childhood body-mass index and the risk of coronary heart disease in adulthood.** *N Engl J Med* 2007, **357**(23):2329–2337. doi:10.1056/NEJMoa072515.
10. Gunnell DJ, Frankel SJ, Nanchahal K, Peters TJ, Smith GD: **Childhood obesity and adult cardiovascular mortality: a 57-y follow-up study based on the Boyd Orr cohort.** *Am J Clin Nutr* 1998, **67**(6):1111–1118.
11. St-Onge MP, Heymsfield SB: **Overweight and obesity status are linked to lower life expectancy.** *Nutr Rev* 2003, **61**(9):313–316. doi:10.1301/nr.2003.sept.313-316.
12. Lake JK, Power C, Cole TJ: **Women's reproductive health: the role of body mass index in early and adult life.** *Int J Obes Relat Metab Disord* 1997, **21**(6):432–438. doi:10.1038/sj.ijo.0800424.
13. Arksey H, O'Malley L: **Scoping studies: towards a methodological framework.** *Int J Soc Res Methodol* 2005, **8**(1):19–32. doi:10.1080/1364557032000119616.
14. MIT Libraries: *Database Search Tips: Boolean operators.* <http://libguides.mit.edu/content.php?pid=36863&sid=271372>.
15. Ahmed S, Karim R: **Socioeconomic differentials of childhood obesity among school children of affluent society in Dhaka city.** *J Paediatr Child Health* 2010, **46**(2):17–23. doi:10.1111/j.1440-1754.2010.01723.x (abstr).
16. Das SK, Chisti MJ, Huq S, Malek MA, Vanderlee L, Salam MA, Ahmed T, Faruque ASG, Al Mamun A: **Changing trend of overweight and obesity and their associated factors in an urban population of Bangladesh.** *Food Nutr* 2013, **4**:678–689. doi:10.4236/fns.2013.46087.
17. Bhuiyan MU, Zaman S, Ahmed T: **Risk factors associated with overweight and obesity among urban school children and adolescents in Bangladesh: a case-control study.** *BMC Pediatr* 2013, **13**(1):72. doi:10.1186/1471-2431-13-72.
18. Hossain MG, Bharati P, Aik SAW, Lestrel PE, Abeer A, Kamarul T, Aekplakorn W, Mo-Suwan L, Al-Isa AN, Bendixen H: **Body mass index of married Bangladeshi women: trends and association with socio-demographic factors.** *J Biosoc Sci* 2012, **44**(4):385. doi:10.1017/S002193201200003X.
19. Mohsin F, Mahbuba S, Begum T, Azad K, Nahar N: **Prevalence of impaired glucose tolerance among children and adolescents with obesity.** *Mymensingh Med J* 2012, **21**(4):684–690.
20. Saha S, Zahid MK, Rasheed S: **The Study of the level of knowledge, attitude, practices (KAP) as well as the effects of school environment on the nutritional status of children (7–12) coming from affluent families in the Dhaka city in Bangladesh.** *J Nutr* 2011–2012, **24**–25:31–48. doi:10.3329/bjnut.v24i0.14035.
21. Corsi DJ, Kyu HH, Subramanian SV: **Socioeconomic and geographic patterning of under-and overnutrition among women in Bangladesh.** *J Nutr* 2011, **141**(4):631–638. doi:10.3945/jn.110.131599.
22. Jesmin A, Yamamoto SS, Malik AA, Haque MA: **Prevalence and determinants of chronic malnutrition among preschool children: a cross-sectional study in Dhaka City, Bangladesh.** *J Health Popul Nutr* 2011, **29**(5):494–499.
23. Mohsin F, Baki A, Nahar J, Akhtar S, Begum T, Azad K, Nahar N: **Prevalence of metabolic syndrome among obese children and adolescents.** *Birdem Med J* 2011, **1**(1):21–25. doi:10.3329/birdem.v1i1.12382.
24. Mohsin F, Tayyeb S, Baki A, Sarker S, Zabeen B, Begum T, Azad K, Nahar N: **Prevalence of obesity among affluent school children in Dhaka.** *Mymensingh Med J* 2010, **19**(4):549–554.
25. Sultana S: *Prevalence and risk factor of childhood overweight and obesity in primary school children of Dhaka city,* Master thesis. University of Oslo: Faculty of Medicine; 2010. <http://urn.nb.no/URN:NBN:no-26615>.
26. Kurshed AAM, Rana MM, Khan S, Azad TMA, Begum J, Bhuyan MAH: **Dietary intake, physical activities and nutritional status of adolescent girls in an urban population of Bangladesh.** *Ibrahim Medical College Journal* 2010, **4**(2):78–82. doi:10.3329/imcj.v4i2.6502.
27. Rahman M, Mostofa G, Nasrin SO: **Nutritional status among children aged 24–59 months in rural Bangladesh: an assessment measured by BMI index.** In *Internet Journal of Biological Anthropology. Volume 3.* Texas: Internet Scientific Publications LLC; 2009.
28. Balarajan Y, Villamor E: **Nationally representative surveys show recent increases in the prevalence of overweight and obesity among women of reproductive age in Bangladesh, Nepal, and India.** *J Nutr* 2009, **139**(11):2139–2144. doi:10.3945/jn.109.112029.
29. Khan MMH, Kraemer A: **Factors associated with being underweight, overweight and obese among ever-married non-pregnant urban women in Bangladesh.** *Singapore Med J* 2009, **50**(8):804–813.
30. Sultan K, Habiba T: **Prevalence of overweight and obesity in infancy.** *Bangladesh Med Res Counc Bull* 2008, **34**(2):69–70. doi:10.3329/bmrcb.v34i2.1180.
31. Shafique S, Akhter N, Stallkamp G, de Pee S, Panagides D, Bloem MW: **Trends of under-and overweight among rural and urban poor women indicate the double burden of malnutrition in Bangladesh.** *Int J Epidemiol* 2007, **36**(2):449–457. doi:10.1093/ije/dyl306.
32. BBS/UNICEF: *Child and Mother Nutrition Survey 2005.* Dhaka: Bangladesh Bureau of Statistics and UNICEF; 2007.
33. Rahman SMM, Kabir I, Khaled MA, Bhuyan MAH, Rashid HA, Malek MA, Khan MR: **Prevalence and Determinants of Childhood Obesity in Dhaka City [abstract].** In *Proceedings of 10th ASCON.* Dhaka; 2002.
34. de Onis M, Blossner M: **Prevalence and trends of overweight among preschool children in developing countries.** *Am J Clin Nutr* 2000, **72**(4):1032–1039.
35. Rahman SMM, Akter BMD, Siddiqui MZA: **Prevalence of childhood Obesity in Dhaka city.** *Mymensingh Med J* 1998, **7**(1):3–6.
36. Ferdousi J, Alamgir AKM: **Prevalence and Determinants of Overweight in School Students: A Developing Country Perspective [abstract].** *Can J Diabetes* 2011, **35**(2):195.
37. Saraswathi S, Najafi M, Gangadhar MR, Malini SS: **Prevalence of childhood obesity in school children from rural and urban areas in Mysore, Karnataka.** *India. J Life Sci* 2011, **3**(1):51–55.
38. Sakamoto N, Wansom S, Tontisirin K, Marui E: **A social epidemiologic study of obesity among preschool children in Thailand.** *Int J Obes Relat Metab Disord* 2001, **25**(3):389–394. doi:10.1038/sj.ijo.0801477.
39. Weiss R, Caprio S: **The metabolic consequences of childhood obesity.** *Best Pract Res Clin Endocrinol Metab* 2005, **19**(3):405–419. doi:10.1016/j.beem.2005.04.009.
40. Ebbeling CB, Pawlak DB, Ludwig DS: **Childhood obesity: public-health crisis, common sense cure.** *Lancet* 2002, **360**(9331):473–482. doi:10.1016/S0140-6736(02)09678-2.

41. Csabi G, Torok K, Jeges S, Molnar D: **Presence of metabolic cardiovascular syndrome in obese children.** *Eur J Pediatr* 2000, **159**(1-2):91-94.
42. Pelin A-M, Matasaru S: **Metabolic syndrome in obese children and adolescents.** *Revista medico-chirurgicala a Societatii de Medici si Naturalisti din Iasi* 2012, **116**(4):957-961.
43. Hannon TS, Rao G, Arslanian SA: **Childhood obesity and type 2 diabetes mellitus.** *Am Acad Pediatrics* 2005, **116**(2):473-480. doi:10.1542/peds.2004-2536.
44. Fagot-Campagna A: **Emergence of type 2 diabetes mellitus in children: epidemiological evidence.** *J Pediatr Endocrinol Metab* 2000, **13**:1395-1402.
45. Fagot-Campagna A, Pettitt DJ, Engelgau MM, Burrows NR, Geiss LS, Valdez R, Beckles GLA, Saaddine J, Gregg EW, Williamson DF: **Type 2 diabetes among North American children and adolescents: an epidemiologic review and a public health perspective.** *J Pediatr* 2000, **136**(5):664-672. doi:10.1067/mpd.2000.105141.
46. Gupta N, Goel K, Shah P, Misra A: **Childhood obesity in developing countries: epidemiology, determinants, and prevention.** *Endocr Rev* 2012, **33**(1):48-70. doi:10.1210/er.2010-0028.
47. Kelishadi R: **Childhood overweight, obesity, and the metabolic syndrome in developing countries.** *Epidemiol Rev* 2007, **29**(1):62-76. doi:10.1093/epirev/mxm003.

doi:10.1186/1471-2458-14-70

Cite this article as: Rahman et al.: Obesity and overweight in Bangladeshi children and adolescents: a scoping review. *BMC Public Health* 2014 **14**:70.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

