

RESEARCH ARTICLE

Open Access

A Chinese herbal formula “Gan-Lu-Yin” suppresses vascular smooth muscle cell migration by inhibiting matrix metalloproteinase-2/9 through the PI3K/AKT and ERK signaling pathways

Yi-Chung Chien¹, Ming-Jyh Sheu^{2*}, Chieh-Hsi Wu², Wen-Hsin Lin², Ying-Yi Chen², Po-Liang Cheng¹ and Hsu-Chen Cheng^{1*}

Abstract

Background: This study was to explore the effects of Gan-Lu-Yin (GLY) on the migration of vascular smooth muscle cells (VSMCs) induced by fetal bovine serum and on neointima formation in a rat model of carotid artery balloon injury.

Methods: VSMCs were treated with different concentrations of GLY, and then analyzed with Flow cytometric analysis, zymography, transwell, and western blotting. SD rats received balloon-injury were analyzed with H&E staining.

Results: Our results showed that GLY significantly decreased the thickness of neointima. The inhibition by non-cytotoxic doses of GLY of VSMCs migration was through its negative regulatory effects on phosphorylated ERK1/2, PI3K/AKT, and FAK. The data showed that GLY can inhibit the migration of VSMCs cells, and might block injury-induced neointima hyperplasia via the inhibition of VSMCs migration, without inducing apoptosis.

Conclusions: These observations provide a mechanism of GLY in attenuating cell migration, thus as a potential intervention for restenosis.

Keywords: Chinese herbs, Gan-Lu-Yin, Migration, Restenosis, Vascular smooth muscle cell

Background

Angiographic restenosis is still a major limitation for the clinical application of percutaneous transluminal coronary angiography (PTCA). It has been well documented that restenosis is a complex and multifactorial process involving arterial remodeling and neointima hyperplasia, which might be caused by endothelial disruption, inflammation, cell proliferation and migration of vascular smooth muscle cells (VSMCs) [1]. Thereby, the pharmacological effects of drug-eluting stents for clinical trials

were focused on the prevention of migration and proliferation [2].

Gan-Lu-Yin (GLY), a Chinese herbal formula, consists of *Rehmannia glutinosa*, *Liriope spicata* (Thunb.) Lour, *Eriobotrya japonica* (Thunb.) Lindl, *Citrus sinensis* Osbeck, *Glycyrrhiza uralensis* Fisch, *Artemisia capillaris* Thunb., *Dendrobium nobile* Lindl., and *Scutellaria baicalensis* Georgi. Many of the GLY components, such as *R. glutinosa*, *L. spicata* (Thunb.) Lour, *E. japonica* (Thunb.) Lindl., *C. sinensis* Osbeck, *G. uralensis* Fisch, *A. capillaris* Thunb., *D. nobile* Lindl., and *S. baicalensis* Georgi, have been used as health foods for a long history in Taiwan. Besides, GLY has been used as a popular drink to manage constipation or oral ulcers. This formula is used to expel the heat, remove the dampness,

* Correspondence: soybean13mtdtw@gmail.com; hardway19800710@gmail.com

¹Department of Life Sciences, and Agricultural Biotechnology Center, National Chung Hsing University, Taichung 402, Taiwan

²School of Pharmacy, China Medical University, 91 Hsueh-Shih Road, Taichung 404, Taiwan

resolve inflammation, and clean the blood according to traditional Chinese medicinal prescriptions, Tai Ping Hui Min He Ji Ju Fang. Numerous studies mentioned that some of the single ingredients within the GLY formula have biological effects on antitumor potential by inhibiting cell proliferation or inflammation [3-5]. Our recent observations suggested that GLY extract has an inhibitory effect on angiogenesis, which in turn may prevent tumor growth, and its mechanism might be partially associated with blocking vascular endothelial growth factor (VEGF) protein expression of HUVEC [6]. However, the investigation regarding the attenuation of balloon injury-induced neointima formation has never been explored. In this study, our results have shown that GLY extract has an inhibitory effect on balloon injury-induced neointima formation. Moreover, the effect of GLY formula on fetal bovine serum (FBS)-induced VSMCs cells migration was studied. The present study was designed to explore the anti-proliferative and -migration effects of GLY on vascular smooth muscle cells and to evaluate its molecular mechanisms.

Methods

Reagents and chemicals

Dulbecco's Modified Eagle's Medium (DMEM), 3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide (MTT), RNase A, and other chemicals were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Trypsin - EDTA, fetal bovine serum (FBS), and penicillin/streptomycin were from Gibco Life Technologies, Inc. (Paisley, UK). Cell culture supplies were purchased from Costar (Corning, Inc., Cypress, CA, USA). The antibody against PI3K, Akt, MAPK/extracellular signal-regulated kinase (ERK) 1/2, and phosphorylated proteins were purchased from Cell Signaling Technology (Beverly, MA, USA). Anti-ERK1/2, anti-PI3K, antifocal adhesion kinase (FAK), anti-p-FAK, and horseradish peroxidase-conjugated goat antimouse IgG antibody were purchased from Santa Cruz Biotechnology Co. (Santa Cruz, CA, USA).

Materials

The ingredients of GLY, *R. glutinosa* (rehmannia), *L. spicata* (Thunb.) Lour (lilyturf), *E. japonica* (Thunb.) Lindl. (loquat leaves), *C. sinensis* Osbeck (sweet orange), *G. uralensis* Fisch (licorice root), *A. capillaris* Thunb. (capillaris), *D. nobile* Lindl. (dendrobium), and *S. baicalensis* Georgi (baical skullcap root), were provided from the Pharmacy Department of China Medical University Hospital, Taichung, Taiwan. All other reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Preparation of GLY extract

The ingredients of the GLY formula were equally weighed (about 1 kg) and soaked in 10 L of 50% ethanol solution (extractive solvent) for 3 days at room temperature. The solid residue of the above soaked herbs was filtered and discarded through a Buchner funnel lined with Whatman filter paper, and the filtrate was concentrated to paste by distillation under reduced pressure. The series concentrations (0.0625, 0.125, 0.25, 0.5, 1.0 and 2.0 mg/mL) of GLY extract were further diluted with deionized water for the subsequent studies.

Cell culture

VSMCs derived from rat thoracic aorta were obtained from Food Industry Research and Development Institute (Hsinchu, Taiwan). Cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) fetal bovine serum (FBS), 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units of penicillin, and 100 mg of streptomycin per mL. The cells were kept in a humidified 5% CO₂/95% air incubator at 37°C.

Cytotoxicity assay

MTT assay was performed to measure the cytotoxicity of GLY on VSMCs were seeded in ninety-six-well plates with 1×10^4 cells/well in DMEM supplemented with 10%FBS. After 24 h, cells were washed with PBS and then exposed to either 10% FBS alone or serial dilutions (0.0625, 0.125, 0.25, 0.5, 1.0 and 2.0 mg/mL) of GLY. After 24 h, the number of viable cells was determined by ELISA reader (Anthos 2001; Anthos Labtec, Salzburg, Austria).

Flow cytometric analysis

Cellular total DNA contents of the treated cells were assessed using flow cytometry following propidium iodide (PI) staining. Cells were analysed using a FACScan flow cytometer (Becton Dickinson, San Jose, CA, USA). PI fluorescence was linearly amplified and both the area and width of the fluorescence pulse were measured. Ten thousand events were acquired, and the percentages of hypodiploid (apoptotic, sub-G₁) events and percentages of cells in the G₀/G₁, S and G₂-M phases were determined using the DNA analysis software ModFitLT, version 2.0 (Verity Software, Topsham, ME, USA).

Wound healing assay

For cell motility determination, VSMCs (3×10^4) were seeded in a 6-well tissue culture plate and grown to 80-90% confluence. After aspiration of the medium, the center of the cell monolayers was scraped with a sterile micropipet tip to create a denuded zone (gap) of constant width. Subsequently, cellular debris was washed with PBS, and VSMCs cells were exposed to various

concentrations of GLY (0, 0.0625, 0.125, 0.25, and 0.5 mg/mL). Wound closure was monitored and photographed at 0 and 18 h with a Nikon inverted microscope. To quantify the migrated cells, pictures of the initial wounded monolayers were compared with the corresponding pictures of cells at the end of the incubation. Artificial lines fitting the cutting edges were drawn on pictures of the original wounds and overlaid on the pictures of cultures after incubation. Cells that had migrated across the white lines were counted in six random fields from each triplicate treatment.

Cell migration assay

VSMCs migration was assayed in transwell chambers (Millipore) according to the method reported by Huang et al. with some modifications [7]. Briefly, transwell chambers with 6.5 mm polycarbonate filters of 8 μ m pore size were used. VSMCs cells (5×10^5 mL⁻¹) and 0, 0.0625, 0.125, 0.25, and 0.5 mg/mL of GLY were suspended in DMEM (100 μ L, serum free), placed in the upper transwell chamber, and incubated for 18 h at 37°C. Then, the cells on the upper surface of the filter were completely wiped away with a cotton swab, and the lower surface of the filter was fixed in methanol, stained with Giemsa, and counted under a microscope at a magnification of 100 \times and 200 \times . For each replicate, the VSMCs cells in 10 randomly selected fields were determined and the counts were averaged.

Determination of MMP-2 and MMP-9 by zymography

MMP in the medium released from VSMCs was assayed using gelatin zymography (7.5% zymogram gelatin gels) according to the methods reported by Huang et al., with some modification [7]. Briefly, the culture medium was electrophoresed (120 V for 90 min) in a 7.5% SDS-PAGE gel containing 0.1% gelatin. The gel was then washed at room temperature in a solution containing 2.5% (v/v) Triton X-100 with two changes and subsequently transferred to a reaction buffer for enzymatic reaction containing 1% NaN₃, 10 mM CaCl₂ and 40 mM Tris-HCl, pH 8.0, at 37°C with shaking overnight (for 12-15 h). Finally, the electrophoretic gel was stained for 30 min with 0.25% (w/v) coomassie blue in 10% acetic acid (v/v) and 20% methanol (v/v) and destained in 10% acetic acid (v/v) and 20% methanol (v/v).

Western blotting analysis

VSMCs cultured in six-well plates were incubated with GLY at 0.125, 0.25 and 0.5 μ g/mL in DMEM containing 10% FBS for 24 h. The cells were then lysed in a buffer containing 2% SDS, 50 mM-dithiothreitol, 62.5 mM-Tris-HCl, pH 6.8, followed by incubation at 95°C for 5 min. Samples were separated using SDS-PAGE, transferred to polyvinylidene fluoride (PVDF) membranes,

blocked with 5% non-fat dry milk in PBS-Tween for 1 h, and then probed with the desired antibodies (anti-phosphorylated focal adhesion kinase (p-FAK; 1:1000), anti-phosphorylated ERK1/2 (1:1000), anti PI3K (1:2000), anti-AKT (1: 200), anti-pAKT (1:200), anti-MMP-2 (1:2000), anti-MMP-9 (1:1000), anti-TIMP-1 (1:1000) and anti-TIMP-2 (1:200); Novus Biologicals, Littleton, CO, USA) overnight at 4°C. The blots were then incubated with horseradish peroxidase-linked secondary antibody for 1 h followed by development with the electrochemical luminescence (ECL) reagent and exposure to Hyperfilm (Amersham, Arlington Heights, IL, USA).

Balloon angioplasty

Sprague-Dawley rats weighing 350-400 g were purchased from National Science Council (Taipei, Taiwan). Sixteen rats were used and divided as two groups including total injury control without GLY (n = 8), and 0.5 g/mL (n = 8) of GLY-treated groups. Animals were housed in a 12 h light-dark cycle with free access to food and water. All experimental procedures involving animals were approved by the ethics committee of the Institutional Animal Care and Use Committee of China Medical University. The rats were anaesthetised with 3.6% (w/v) chlorohydrate (1 mL/100 g, intraperitoneally). Angioplasty of the carotid artery was performed using a balloon embolectomy catheter as described previously [8].

Statistical analysis

Values are expressed as means \pm SD and analyzed using one-way ANOVA followed by LSD test for comparisons of group means. All statistical analyses were performed using SPSS for Windows, version 10 (SPSS, Inc.). The level of significance was set at *P* value <0.05.

Results

GLY inhibited FBS-induced VSMCs proliferation

Since outgrowth of VSMCs has been regarded as the major factor leading to restenosis, we performed the MTT assay to determine the inhibitory effects of GLY on cell viability of VSMCs. As shown in Figure 1A, GLY inhibited VSMCs viability in a concentration-dependent manner. The inhibitory effect of GLY on cell viability became significant at 0.5 mg/mL (*P* < 0.05), 1 mg/mL (*P* < 0.01) and 2 mg/mL (*P* < 0.001) after 24 h incubation.

Effects of GLY on cell cycle

Since the MTT showed that GLY at 0.5, 1 and 2 mg/mL significantly suppressed cell viability, we postulated that the inhibitory effects of GLY on cell viability might be mediated by apoptosis. We chose GLY at 0.0625, 0.125, 0.25, 0.5 and 1.0 mg/mL to determine its effects on cell cycle and apoptosis. The results demonstrate that




Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 Effects of GLY on cell viability and cell cycle analysis. (A). Effects of GLY on cell growth of VSMCs by MTT assay. The cells were incubated for 24 h with 10% fetal bovine serum alone (control) or with different concentrations of GLY (0.0625, 0.125, 0.25, 0.5, 1 and 2 mg/mL). (B). Flow cytometric analysis of GLY on the cell cycle of VSMCs cells. All the cells were treated with 15% fetal bovine serum with the addition of GLY at 0.0625, 0.125, 0.25, 0.5, 1 mg/mL for 18 h. (C). The percentage of subG₁ under flow cytometric analysis. Values are means of three separate experiments, with standard errors represented by vertical bars. Mean value was significantly different from that of the control group: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

treatment for 24 h with GLY at 0.0625, 0.125, 0.25 and 0.5 mg/mL had no effects on the cells apoptotic in the sub-G₁ phase, however, treatment for 24 h. with GLY at 1.0 mg/mL demonstrated significant apoptosis (Figures 1B and 1C).

Non-cytotoxic doses of GLY inhibited FBS-stimulated VSMCs motility in wound healing assay

We assessed the effect of GLY on the migration of VSMCs cells using the wound healing assay in which the confluent monolayer was scraped with a sterile micropipet tip to create a scratch wound. As shown in our data, GLY at 0.25 and 0.5 mg/mL inhibited the migration of VSMCs cells (Figures 2 A and 2B).

Non-cytotoxic doses of GLY inhibited FBS-stimulated VSMCs migration

The transwell assay was used to investigate the migration of VSMCs cells at 18 h after GLY treatment. We found that GLY added at 0.125, 0.25 and 0.5 mg/mL significantly decreased the migration (Figures 2 C and 2D) of VSMCs cells.

Non-cytotoxic doses of GLY inhibits the Release of MMP-2 and MMP-9 in VSMCs Cells

To examine the possible antimetastatic mechanisms of GLY, we determined the activity of MMP-2 and MMP-9 in culture media of VSMCs cells by zymographic analysis. In the absence of treatment, VSMCs cells constitutively secreted both MMP-9 and MMP-2. As shown in Figure 3A, GLY inhibited MMP-9 and MMP-2 activities in a concentration-dependent manner after incubation for 18 h (Figures 3 B and 3C).

Non-cytotoxic doses of GLY activated TIMPs, MMP-2, and MMP-9 expressions in VSMCs cells

To further explore the modulation of pro-MMP and MMP activation by GLY, we determined TIMP-1/2 and MMP2/9 proteins expression levels. As shown in Figure 4A, GLY strongly increased TIMP-1/2 activity and also decreased MMP-2/9 activity in a concentration-dependent manner.

Non-cytotoxic doses of GLY inhibited phosphorylated focal adhesion kinase and phosphorylated extracellular signal-regulated kinase

To evaluate the effect of GLY on FAK and ERK1/2 protein expression, VSMCs was treated with GLY at 0.125, 0.25 and 0.5 mg/mL for 30 min. As shown in Figure 4B, GLY had no effect on FAK expression. In addition, GLY at 0.25 and 0.5 mg/mL significantly suppressed the phosphorylation of FAK (p-FAK) in VSMCs cells. (Figure 4B). We analyzed the phosphorylation of ERK1/2 in VSMCs cells after treatment with GLY (0.125, 0.25 and 0.5 mg/mL) for 30 min. Data in Figure 4B showed that GLY significantly affect phosphorylated ERK1/2 at concentrations used (0.25 and 0.5 mg/mL).

Non-cytotoxic doses of GLY inhibited the PI3K/Akt signaling in VSMCs cells

To further investigate the involvement of PI3K/Akt, a series of experiments was performed to measure the expression of candidate signaling molecules upon GLY stimulation. The results showed that incubation of VSMCs cells with GLY (0.125, 0.25 and 0.5 mg/mL) led to a concentration-dependent decrease of PI3K and p-Akt levels (Figure 4B).

Extract of GLY inhibited balloon injury-induced neointima formation on the carotid artery


To test the efficacy of GLY in inhibiting neointima formation, Sprague–Dawley rats were applied with different concentrations of GLY (0.5 g/mL) for 14 days following balloon injury. After 2 weeks of balloon injury, the injured arteries were harvested and subjected to histological analysis for neointima formation assay. Intimal hyperplasia induced by balloon injury was evident as compared with the normal control (Figure 5A). The present results showed that GLY (0.5 g/mL) were effective in preventing neointima formation (Figure 5B). However, GLY at 0.125 g/mL and 0.25 g/mL did not show any influence on balloon injury-induced neointimal formation (data not shown). Using computerized image analysis, we calculated the area ratio of intimal and media layers; we found a reduction of 24% in the area ratio of GLY-treated groups as compared with the balloon-injured control group by GLY at 0.5 g/mL of GLY, respectively (Figure 5B).

Discussion

Earlier studies have shown that pretreatment with antioxidants can significantly reduce balloon injury-induced neointima formation [9]. Our recent data have demonstrated that the MDA levels were significantly reduced by GLY treatment. Also, GLY extracts are effective in scavenging α, α -diphenyl- β -picrylhydrazyl (DPPH) radicals (data not shown). We found GLY extract could induce apoptosis of vascular endothelial cells in a dose-dependent manner [6]. GLY having antioxidation and free radical scavenging activities may be developed as another potential candidate to prevent restenosis. To test whether GLY can be an effective therapeutic intervention for balloon injury, administration of GLY in preventing abnormal cell proliferation were evaluated at various concentrations in vivo. And, we aimed to investigate if GLY contributed to the attenuation of VSMCs migration and therefore, explored its molecular

mechanisms. The present study provides a general insight into the molecular mechanism of GLY in preventing the outgrowth of smooth muscle cells, which is a potential intervention for balloon injury-induced neointima formation.

Recently, Lin et al. mentioned that at least 14 main components of GLY have been identified using liquid chromatography-mass spectrometry (LC-MS) and inductively coupled plasma MS (ICP-MS), and these components included baicalin, baicalein, oroxylin A-7-O-glucuronide, wogonin-7-O-glucuronide, wogonin, and oroxylin A in *Radix Scutellariae*; naringin and neohesperidin in *Aurantii fructus*; and liquiritigenin, liquiritin, and glycyrrhizic acid in *Radix Glycyrrhizae* [10]. In our previous study, the indicator components of GLY extract such as baicalein, chlorogenic acid, and glycyrrhizic acid have been detected by HPLC/ESI-MS analysis [6]. Four other components such as baicalin, naringin, naringenin,







Figure 3 GLY inhibits the release of MMP-2 and MMP-9 in VSMCs. (A). To examine the possible antimigration mechanisms of GLY, we determined the activity of MMP-2 and MMP-9 in culture media of VSMCs cells by zymographic analysis. (B). GLY inhibited MMP-9 and MMP-2 activities in a concentration-dependent manner. Values are means of three separate experiments, with standard errors represented by vertical bars. Mean value was significantly different from that of the control group: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

and vanillin have also been detected from our study (data not shown). It has been mentioned that baicalin and baicalein, two flavonoid compounds, have inhibitory effects on endothelial cell proliferation, migration, and differentiation [11,12]. Endothelial cell proliferation, adhesion, and migration are early essential events for mediating angiogenesis. Hsieh et al. indicated that baicalein significantly up-regulated protein expression of integrins and vinculin in rat endothelial cells, which will result in the increase of focal contact formation and cell adhesion to fibronectin and vitronectin [11]. Coordination and balance between cell adhesion to and detachment from extracellular matrix (ECM) is crucial for cell motility, so unilateral increasing cell adhesion will severely affect endothelial migration and angiogenesis processes. In addition, it has been reported that baicalein and baicalin could inhibit cell proliferation of tumor cells and induce apoptosis of myeloma cells [13,14]. Sun et al. found that oroxylin A suppresses invasion through inhibiting cell migration and down-regulating the expression of matrix metalloproteinase-2/9 in human breast cancer cells [15]. Wogonin has been evidenced to inhibit inflammatory cytokine-induced angiogenesis of HUVEC and suppress the VEGF-stimulated migration and tube formation of HUVEC [16,17]. Wogonin effectively suppresses TNF- α -induced HASMC migration through the selective inhibition of MMP-9 expression [18]. Schindler et al. reported

that naringin could decrease tumor-induced vascular proliferation by reducing the release of VEGF from human tumor cells [19]. The inhibitory effect of liquiritigenin on tube formation of vascular endothelial cells has also been reported [20]. However, glycyrrhizic acid, the main component of licorice, has been investigated for its ability to increase tube formation [20]. Jung et al. mentioned that caffeic acid and its derivative, CAPDE ([3-(3,4-dihydroxyphenyl)-acrylic acid 2-(3,4-dihydroxyphenyl)-ethyl ester]) can suppress tumor angiogenesis by blocking VEGF expression in human renal carcinoma cells [21]. Although chlorogenic acid is also a caffeic acid derivative compound, so far there has not been direct evidence to point out the association with the effect on antiangiogenesis. However, previous literature indicated that it is a new type of strong matrix metalloproteinase-9 inhibitor, which might prevent the invasion and metastasis of malignant cancer cells [22]. According to the above statements, we suppose that these ingredients from GLY might contribute the inhibition of proliferation and migration in VSMCs.

VSMCs proliferation and migration are important contributors to neointima formation after balloon injury. Therefore, modulation of VSMCs growth has critical therapeutic implications [23]. In the present study, for the first time we show that GLY attenuates neointima hyperplasia after balloon angioplasty. To evaluate if GLY



was effective in suppressing neointima formation following balloon angioplasty, an in vivo study using rat carotid artery as a model was conducted in the present study. Sixteen rats were used and divided as two groups including total injury control without GLY, and 0.5 g/mL of GLY-treated groups. A balloon catheter was first surgically inserted into rat carotid arteries to induce injury. At 2 weeks after balloon injury, the arteries were subjected to histological analysis and GLY was found to significantly reduce neointima formation 14d following arterial injury (Figure 5A). The neointima:media ratios

of arterial samples from animals treated with GLY were significantly lower than those of the total injury control tissues (Figure 5B).

GLY was carried out on its mechanisms on serum-induced VSMCs migration behavior. VSMCs migration is important contributors to neointima formation after balloon injury. We first demonstrated that GLY exerted potent inhibitory effects on the growth of VSMCs (Figure 1A). Since the MTT showed that GLY at 1 and 2 mg/mL significantly suppressed cell viability, we postulated that the inhibitory effects of GLY on cell viability




Figure 5 (See legend on next page.)

(See figure on previous page.)

Figure 5 Responses of rat carotid arteries to balloon injury, and the effects of extract of GLY on balloon injury. (A). The left panel represents the low-power (100x) observations from a balloon-injured vessel, a balloon-injured vessel treated with GLY at 0.5 g/ml, and uninjured vessel (sham control). And the right panel represents the high-power (200x). (L, lumen; N, neointima; M, medium). (B). The control group shows a significantly higher neointima to media ratio as compared with the groups treated with extract of GLY at 0.5 g/mL. Values are means of two separate experiments, with standard errors represented by vertical bars. Mean value was significantly different from that of the control group: * $p < 0.05$.

might be mediated by apoptosis. We chose GLY at 0.0625, 0.125, 0.25, 0.5 and 1.0 mg/mL to determine its effects on cell cycle and apoptosis. The results demonstrate that treatment for 24 with GLY at 0.0625, 0.125, 0.25 and 0.5 mg/mL had no effects on the cells apoptotic in the sub- G_1 phase (Figures 1 B and 1C). Therefore, GLY at 0.125, 0.25 and 0.5 mg/mL was conducted on its role in migration analysis. We found that GLY has unique biological properties; it does not affect cell survival (Figure 1A) and apoptosis (Figures 1 B and 1C) at concentrations up to 0.5 mg/mL, whereas it has profound effects on migration at 0.125, 0.25 and 0.5 mg/mL (Figures 2 and 3).

The MAPK pathways play an important role in promoting VSMCs proliferation [24]. The MAPK signaling pathway was activated by vascular injury and the inhibition of activated ERK pathway by drugs or gene therapy can reduce neointima hyperplasia [25]. Our previous reports show that the *ras* gene is involved in the underlying mechanisms for neointima formation by balloon injury [26]. Therefore, several proteins involved in these pathways were examined. FAK is a protein involved in transducing extracellular growth signal from matrix via integrin interaction. Down-regulation of FAK may result in cell-cycle arrest [27]. Activated FAK in cancer cells relays signals through multiple downstream targets. For example, activated FAK binds the Src-homology domain 2 (SH2) of PI3K, thereby transporting the catalytic subunit of PI3K to the membrane, where it catalyzes the phosphorylation of inositol lipids [28]. The residues surrounding Tyr³⁹⁷ can also constitute a sequence that binds to the Ras signaling pathway. The downstream targets of the Ras signaling pathway include ERK1/2 [29]. Indeed, these pathways are activated during integrin binding to the ECM, resulting in the transduction of external stimuli from the ECM to the nucleus [30]. Raf, an important protein in the MAPK pathway, is responsible for signal transduction from Ras to ERK. Along the pathway, signaling of phosphorylated-ERK1/2 is also an essential element for cell proliferation. Therefore, the protein expression levels of FAK and ERK were all evaluated in the present study to explore the mode of preventive action of GLY against neointima formation by balloon injury. In this study, we found that GLY inhibited the activation of FAK, as evidenced by reduced phosphorylation of FAK

(Figure 4B). We also demonstrated in the present study that the level of phosphorylated-ERK was down-regulated by GLY and was in accordance with the well-known function of ERK as a critical signaling molecule leading to cell proliferation and survival (Figure 4B) [31].

Phosphoinositide 3-kinase (PI3K)/AKT signaling pathway plays a major role in the regulation of cellular growth, apoptosis, and metabolism [32]. The PI3K/AKT signaling is reportedly required for VSMCs migration and proliferation, absence of AKT impairs VSMCs proliferation and migration [33]. Level of AKT kinase phosphorylation was increased by serum in our study and GLY inhibited serum-induced phosphorylation of AKT and PI3K, indicating that AKT protein is another potential target for GLY (Figure 4B). AKT mediates cell survival and growth signals by phosphorylating and inactivating pro-apoptotic proteins [34]. Therefore, these results indicate that the ERK1/2-MAPK pathway likely synergizes with other pro-proliferative signaling pathway (s), quite possibly the PI3K/AKT pathway, to regulate serum-induced VSMCs cell proliferation and migration.

The present study demonstrated that balloon injury-induced neointima formation could be markedly reduced by GLY. Its pharmacological mechanism may be associated with the down-regulation of FAK and ERK1/2 phosphorylation proteins levels. Also, downregulation of PI3K and AKT phosphorylation could be involved in the progress of the progress of VSMCs metastasis. The present results detailed the molecular mechanisms of GLY in preventing the smooth muscle cell proliferation in vitro. In the in vivo study, we found that treatments with GLY at 0.5 g/mL significantly reduced the neointima formation in rat carotid arteries after balloon injury. Our findings regarding the inhibitory effects of GLY on smooth muscle cells may shed light onto the conjunctive roles of GLY with some other pharmacological agents in preventing restenosis. These results demonstrate that GLY suppresses the migration of VSMCs via inhibiting MAPK and PI3K/AKT signaling pathways. We proposed that GLY might be a potential agent that may prove clinically useful in patients undergoing PTCA or stenting. A further study on larger animal models or even a clinical evaluation needs to be conducted to confirm the proposed approach in this aspect.

Conclusion

We have demonstrated that GLY inhibits the migration of VSMCs cells. Mechanistically, GLY may occur through inactivation of the ERK1/2 signaling pathway, exerting inhibitory effects on FAK and pFAK protein expressions and inhibiting PI3K, and phospho-AKT levels, thereby decreasing the activities of MMP-2 and MMP-9 leading to inhibition of metastasis of VSMCs (Figure 4B).

Abbreviations

GLY: Gan-Lu-Yin; FBS: Fetal bovine serum; VSMCs: Vascular smooth muscle cells; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; ERK: Extracellular signal-regulated kinase; p-FAK: Phosphorylated focal adhesion kinase; PI: Propidium iodide; PTCA: Percutaneous transluminal coronary angioplasty.

Competing interests

There are no conflicts of interest.

Authors' contribution

YCC prepared ethanol extract of GLY and animal study. MJS, CHW, WHL, YYC, and PLC performed molecular and cell biology studies of VSMCs. YCC, MJS and HCC conceived the ideas, designed the experiments and interpreted the experimental data. All authors contributed to the manuscript preparations and approved the final manuscript.

Acknowledgements

We are grateful to Jeffery Conrad for English writing assistance. The study was supported by grants from the National Science Council (NSC 100-2320-B-039-014-), China Medical University (CMU99-TC-29), and Department of Health Clinical Trial and Research Center of Excellence (DOH101-TD-B-111-004 and DOH101-TD-B-111-005). This research was also supported by the National Science Council and by the Ministry of Education, Taiwan, Republic of China, under the ATU plan (NSC 101-2311-B-005-006-MY3).

Received: 20 March 2012 Accepted: 17 August 2012

Published: 24 August 2012

References

1. Grüntzig AR, Senning A, Siegenthaler WE: **Nonoperative dilatation of coronary-artery stenosis: percutaneous transluminal coronary angioplasty.** *N Engl J Med* 1979, **301**:61–68.
2. Bauters C, Lablanche JM, Mcfadden EP, Leroy F, Bertrand ME: **Clinical characteristics and angiographic follow-up of patients undergoing early or late repeat dilation for a first restenosis.** *J Am Coll Cardiol* 1992, **20**:845–848.
3. Chao JC, Chiang SW, Wang CC, Tsai YH, Wu MS: **Hot water-extracted Lycium barbarum and Rehmannia glutinosa inhibit proliferation and induce apoptosis of hepatocellular carcinoma cells.** *World J Gastroenterol* 2006, **12**:4478–4484.
4. Kumagai T, Müller CI, Desmond JC, Imai Y, Heber D, Koeffler HP: **Scutellaria baicalensis, a herbal medicine: anti-proliferative and apoptotic activity against acute lymphocytic leukemia, lymphoma and myeloma cell lines.** *Leuk Res* 2007, **31**:523–530.
5. Ye F, Jiang S, Volshonok H, Wu J, Zhang DY: **Molecular mechanism of anti-prostate cancer activity of Scutellaria baicalensis extract.** *Nutr Cancer* 2007, **57**:100–110.
6. Pan CH, Hsieh IC, Liu FC, Hsieh WT, Sheu MJ, Koizumi A, Wu CH: **Effects of a Chinese herbal health formula, "Gan-Lu-Yin", on angiogenesis.** *J Agric Food Chem* 2010, **58**:7685–7692.
7. Huang GJ, Yang CM, Chang YS, Amagaya S, Wang HC, Hou WC, Huang SS, Hu ML: **Hispolon suppresses SK-Hep1 human hepatoma cell metastasis by inhibiting matrix metalloproteinase-2/9 and urokinase-plasminogen activator through the PI3K/Akt and ERK signaling pathways.** *J Agric Food Chem* 2010, **58**:9468–9475.
8. Sheu MJ, Cheng HC, Chien YC, Chou PY, Huang GJ, Chen JS, Lin SY, Wu CH: **Molecular mechanism of green microalgae, Dunaliella salina, involved in attenuating balloon injury-induced neointima formation.** *Br J Nutr* 2010, **104**:326–335.
9. Szöcs K, Lassègue B, Sorescu D, Hilenski LL, Valppu L, Couse TL, Wilcox JN, Quinn MT, Lambeth JD, Griendling KK: **Upregulation of Nox-based NAD(P)H oxidases in restenosis after carotid injury.** *Arterioscler Thromb Vasc Biol* 2002, **22**:21–27.
10. Lin IH, Lee MC, Chuang WC: **Application of LC/MS and ICP/MS for establishing the fingerprint spectrum of the traditional Chinese medicinal preparation Gan-Lu-Yin.** *J Sep Sci* 2006, **29**:172–179.
11. Hsieh YC, Hsieh SJ, Chang YS, Hsueh CM, Hsu SL: **The lipoxigenase inhibitor, baicalein, modulates cell adhesion and migration by up-regulation of integrins and vinculin in rat heart endothelial cells.** *Br J Pharmacol* 2007, **151**:1235–1245.
12. Liu JJ, Huang TS, Cheng WF, Lu FJ: **Baicalein and baicalin are potent inhibitors of angiogenesis: Inhibition of endothelial cell proliferation, migration and differentiation.** *Int J Cancer* 2003, **106**:559–565.
13. Ikezoe T, Chen SS, Heber D, Taguchi H, Koeffler HP: **Baicalin is a major component of PC-SPES which inhibits the proliferation of human cancer cells via apoptosis and cell cycle arrest.** *Prostate* 2001, **49**:285–292.
14. Ma Z, Otsuyama K, Liu S, Abroun S, Ishikawa H, Tsuyama N, Obata M, Li FJ, Zheng X, Maki Y, et al: **Baicalein, a component of Scutellaria radix from Huang-Lian-Jie-Du-Tang (HLJDT), leads to suppression of proliferation and induction of apoptosis in human myeloma cells.** *Blood* 2005, **105**:3312–3318.
15. Sun Y, Lu N, Ling Y, Gao Y, Chen Y, Wang L, Hu R, Qi Q, Liu W, Yang Y, et al: **Oroxilin A suppresses invasion through down-regulating the expression of matrix metalloproteinase-2/9 in MDA-MB-435 human breast cancer cells.** *Eur J Pharmacol* 2009, **603**:22–28.
16. Lin CM, Chang H, Chen YH, Wu IH, Chiu JH: **Wogonin inhibits IL-6-induced angiogenesis via down-regulation of VEGF and VEGFR-1, not VEGFR-2.** *Planta Med* 2006, **72**:1305–1310.
17. Lu N, Gao Y, Ling Y, Chen Y, Yang Y, Gu HY, Qi Q, Liu W, Wang XT, You QD, et al: **Wogonin suppresses tumor growth in vivo and VEGF-induced angiogenesis through inhibiting tyrosine phosphorylation of VEGFR2.** *Life Sci* 2008, **82**:956–963.
18. Lee SO, Jeong YJ, Yu MH, Lee JW, Hwangbo MH, Kim CH, Lee IS: **Wogonin suppresses TNF-alpha-induced MMP-9 expression by blocking the NF-kappaB activation via MAPK signaling pathways in human aortic smooth muscle cells.** *Biochem Biophys Res Commun* 2006, **351**:118–125.
19. Schindler R, Mentlein R: **Flavonoids and vitamin E reduce the release of the angiogenic peptide vascular endothelial growth factor from human tumor cells.** *J Nutr* 2006, **136**:1477–1482.
20. Kobayashi S, Miyamoto T, Kimura I, Kimura M: **Inhibitory effect of isoliquiritin, a compound in licorice root, on angiogenesis in vivo and tube formation in vitro.** *Bio Pharm Bull* 1995, **18**:1382–1386.
21. Jung JE, Kim HS, Lee CS, Park DH, Kim YN, Lee MJ, Lee JW, Park JW, Kim MS, Ye SK, et al: **Caffeic acid and its synthetic derivative CADPE suppress tumor angiogenesis by blocking STAT3-mediated VEGF expression in human renal carcinoma cells.** *Carcinogenesis* 2007, **28**:1780–1787.
22. Jin UH, Lee JY, Kang SK, Kim JK, Park WH, Kim JG, Moon SK, Kim CH: **A phenolic compound, 5-caffeoylquinic acid (chlorogenic acid), is a new type and strong matrix metalloproteinase-9 inhibitor: isolation and identification from methanol extract of Euonymus alatus.** *Life Sci* 2005, **77**:2760–2769.
23. Ross R: **Atherosclerosis-an inflammatory disease.** *New Engl J Med* 1999, **340**:115–126.
24. Indolfi C, Coppola C, Torella D, Arcucci O, Chiariello M: **Gene therapy for restenosis after balloon angioplasty and stenting.** *Cardiol Res* 1999, **7**:324–331.
25. Gennaro G, Ménard C, Michaud SE, Deblois D, Rivard A: **Inhibition of vascular smooth muscle cell proliferation and neointima formation in injured arteries by a novel, oral mitogen-activated protein kinase/extracellular signal-regulated kinase inhibitor.** *Circulation* 2004, **110**:3367–3371.
26. Wu CH, Pan JS, Chang WC, Hung JS, Mao SJ: **The molecular mechanism of actinomycin D in preventing neointima formation in rat carotid arteries after balloon injury.** *J Biomed Sci* 2005, **12**:503–512.
27. Zhao JH, Reiske H, Guan JL: **Regulation of the cell cycle by focal adhesion kinase.** *J Cell Biol* 1998, **143**:1997–2008.

28. Meng XN, Jin Y, Yu Y, Bai J, Liu GY, Zhu J, Zhao YZ, Wang Z, Chen F, Lee KY, *et al*: Characterisation of fibronectin-mediated FAK signalling pathways in lung cancer cell migration and invasion. *Br J Cancer* 2009, **101**:327–334.
29. Lai KC, Huang AC, Hsu SC, Kuo CL, Yang JS, Wu SH, Chung JG: Benzyl isothiocyanate (BITC) inhibits migration and invasion of human colon cancer HT29 cells by inhibiting matrix metalloproteinase-2/-9 and urokinase plasminogen (uPA) through PKC and MAPK signaling pathway. *J Agric Food Chem* 2010, **58**:2935–2942.
30. Liao HF, Chen YY, Liu JJ, Hsu ML, Shieh HJ, Liao HJ, Shieh CJ, Shiao MS, Chen YJ: Inhibitory effect of caffeic acid phenethyl ester on angiogenesis, tumor invasion, and metastasis. *J Agric Food Chem* 2003, **51**:7907–7912.
31. Ghiselli G, Chen J, Kaou M, Hallak H, Rubin R: Ethanol inhibits fibroblast growth factor-induced proliferation of aortic smooth muscle cells. *Arterioscler Thromb Vasc Biol* 2003, **23**:1808–1813.
32. Choi KH, Kim JE, Song NR, Son JE, Hwang MK, Byun S, Kim JH, Lee KW, Lee HJ: Phosphoinositide 3-kinase is a novel target of piceatannol for inhibiting PDGF-BB-induced proliferation and migration in human aortic smooth muscle cells. *Cardiovasc Res* 2010, **85**:836–844.
33. Fernández-Hernando C, József L, Jenkins D, Di Lorenzo A, Sessa WC: Absence of Akt1 reduces vascular smooth muscle cell migration and survival and induces features of plaque vulnerability and cardiac dysfunction during atherosclerosis. *Arterioscler Thromb Vasc Biol* 2009, **29**:2033–2040.
34. Seo JM, Kim TJ, Jin YR, Han HJ, Ryu CK, Sheen YY, Kim DW, Yun YP: YSK2821, a newly synthesized indole derivative, inhibits cell proliferation and cell cycle progression via the cell cycle-related proteins by regulating phosphatidylinositol-3 kinase cascade in vascular smooth muscle cells. *Eur J Pharmacol* 2008, **586**:74–81.

doi:10.1186/1472-6882-12-137

Cite this article as: Chien *et al.*: A Chinese herbal formula “Gan-Lu-Yin” suppresses vascular smooth muscle cell migration by inhibiting matrix metalloproteinase-2/9 through the PI3K/AKT and ERK signaling pathways. *BMC Complementary and Alternative Medicine* 2012 **12**:137.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

