

RESEARCH ARTICLE

Open Access

Prolonged A β treatment leads to impairment in the ability of primary cortical neurons to maintain K⁺ and Ca²⁺ homeostasis

Lana Shabala*, Claire Howells, Adrian K West, Roger S Chung

Abstract

Background: Alzheimer's disease (AD) is a progressive neurodegenerative disease, characterised by the formation of insoluble amyloidogenic plaques and neurofibrillary tangles. Beta amyloid (A β) peptide is one of the main constituents in A β plaques, and is thought to be a primary causative agent in AD. Neurons are likely to be exposed to chronic, sublethal doses of A β over an extended time during the pathogenesis of AD, however most studies published to date using *in vitro* models have focussed on acute studies. To experimentally model the progressive pathogenesis of AD, we exposed primary cortical neurons daily to 1 μ M of A β_{1-40} over 7 days and compared their survival with age-similar untreated cells. We also investigated whether chronic A β exposure affects neuronal susceptibility to the subsequent acute excitotoxicity induced by 10 μ M glutamate and assessed how Ca²⁺ and K⁺ homeostasis were affected by either treatment.

Results: We show that continuous exposure to 1 μ M A β_{1-40} for seven days decreased survival of cultured cortical neurons by 20%. This decrease in survival correlated with increased K⁺ efflux from the cells. One day treatment with 1 μ M A β followed by glutamate led to a substantially higher K⁺ efflux than in the age-similar untreated control. This difference further increased with the duration of the treatment. K⁺ efflux also remained higher in A β treated cells 20 min after glutamate application leading to 2.8-fold higher total K⁺ effluxed from the cells compared to controls. Ca²⁺ uptake was significantly higher only after prolonged A β treatment with 2.5-fold increase in total Ca²⁺ uptake over 20 min post glutamate application after six days of A β treatment or longer ($P < 0.05$).

Conclusions: Our data suggest that long term exposure to A β is detrimental because it reduces the ability of cortical neurons to maintain K⁺ and Ca²⁺ homeostasis in response to glutamate challenge, a response that might underlie the early symptoms of AD. The observed inability to maintain K⁺ homeostasis might furthermore be useful in future studies as an early indicator of pathological changes in response to A β .

Background

Alzheimer's disease (AD) is the most common form of dementia within the ageing population and accounts for between 50% and 60% of dementia cases [1]. Sufferers of AD experience progressive loss of memory and cognitive abilities that eventually lead to dementia and death. The pathological hallmarks of the disease include extracellular β -amyloid (A β) plaques, intracellular neurofibrillary tangles (NFTs) and dystrophic neurites (DNs) [2]. The A β peptide is one of the main constituents in A β plaques, and is thought to be a

primary causative agent in AD, significantly contributing towards AD pathogenesis.

AD is a progressive disease which develops over many years, even decades. Therefore the pathogenesis of AD does not entail a sudden insult of A β which causes widespread neuronal death within the brain. Instead, there appears to be a gradual progression of the disease which involves the accumulation of soluble A β within the brain due to the chronic imbalance between production and clearance of A β . This gradual accumulation of A β over extended periods of time leads to the formation of the insoluble A β aggregates which form the characteristic plaques, but it also modifies neuronal function. It is difficult to correlate early stages of AD pathogenesis with the

* Correspondence: L.Shabala@utas.edu.au
NeuroRepair Group, Menzies Research Institute, University of Tasmania,
Private Bag 23, Hobart, Tasmania, 7001, Australia

accumulation of A β , as patients remain asymptomatic. Studies using transgenic mice which express human mutant A β precursor peptide (APP, Tg2576 mice) demonstrated that the increase of A β ₁₋₄₀ and A β ₁₋₄₂ over several months was accompanied by deficits in normal learning and memory [3,4]. When the A β load was minimal, mice demonstrated normal spatial learning and memory, indicating that the gradual build up of A β correlated with the physiological changes associated with AD. Studies have demonstrated that many molecular forms of A β are neurotoxic causing neuronal cell death *in vitro* and neuronal loss *in vivo*. For example, synthetic A β peptides were toxic to hippocampal and cortical neurons in culture [5,6].

Neurons are likely to be exposed to sublethal doses of A β over an extended time during the pathogenesis of AD. However studies to date have not developed an experimental model of this chronic exposure. There have been numerous *in vitro* models used to examine A β neurotoxicity using various forms and concentration of the peptide, over the short term. For instance Deshpande and colleagues examined the toxicity of a single dose of 5 μ M fibrillar A β ₁₋₄₂ and A β -derived diffusible ligands on human cortical neuron cultures for up to 24 hours [6]. At the same time studies examining the chronic exposure of neurons to A β at sublethal doses would provide valuable information relative to the physiological processes which occur in the AD brain.

The accumulation of toxic A β peptide aggregates in AD brain is thought to trigger the extensive synaptic loss and neurodegeneration linked to cognitive decline, an idea that underlies the 'amyloid hypothesis' of AD etiology in both the familial and sporadic forms of the disease [7]. Recent reports strongly suggest that in the initial stages of AD, glutamate receptors are dysregulated by A β accumulation resulting in disruption of glutamatergic synaptic transmission which parallels early cognitive deficits [8]. High concentrations of glutamate have been documented to cause neuronal degeneration in various *in vivo* and *in vitro* models [9,10]. Therefore, it would be of interest whether chronic neuron exposure to A β would sensitise them to excitotoxicity caused by increased glutamate concentration.

Excessive glutamatergic stimulation is associated with an increase in intracellular calcium ([Ca²⁺]_i). Although Ca²⁺ is necessary for a number of physiological processes, excessive amounts may lead to neuronal dysfunction and cell death. Neuronal increases in [Ca²⁺]_i can activate a number of enzymes, such as phospholipases, proteases, endonucleases and nitric oxide synthase (NOS) that are associated with neuronal cell death [11]. Increases in cytosolic Ca²⁺ levels were observed in AD mouse models strongly supporting the theory that Ca²⁺ dysregulation is involved in AD [12-14]. Recent data

also indicate that rodent and human A β modulate K⁺ currents [15]. Voltage-dependent K⁺ (Kv) channels act as potent modulators of diverse excitatory events that are linked to glutamatergic neurotransmission [16] including modulation of Ca²⁺ uptake. Changes in ion fluxes are one of the earliest events in neural responses to A β treatment that precede neuron survival. Therefore, delineation between neuronal ability to maintain Ca²⁺ and K⁺ homeostasis during prolonged A β treatment and their viability might provide inside into mechanisms underlying processes of neurodegeneration.

To experimentally model the progressive pathogenesis of AD, primary cortical neurons were exposed to sublethal doses of A β ₁₋₄₀ over an extended period of time of 7 days. In this study we compared survival of cortical neurons treated for different periods of time with A β ₁₋₄₀ and correlated them to the neuron's ability to maintain homeostasis of Ca²⁺ and K⁺ ions under the same conditions. Finally, we investigated whether prolonged exposure to relatively low A β ₁₋₄₀ (1 μ M) increased neuronal susceptibility to subsequent acute excitotoxicity induced by the neurotransmitter glutamate.


Results

Continuous exposure to Ab₁₋₄₀ is mildly neurotoxic to cultured cortical neurons

We first examined whether chronic exposure of mature neurons to low levels of A β caused cell death. Cortical neurons were maintained for 6 DIV, by which time they had formed a dense meshwork of neuritic processes. They were then treated with soluble monomeric A β ₁₋₄₀ (1 μ M) daily for up to a further seven days. Daily determination of neuronal viability by an Alamar Blue assay revealed that A β ₁₋₄₀ treatment did not reduce cell viability for cells treated for up to six days (Figure 1). However, by seventh day of the treatment neuronal viability decreased by 20% ($P < 0.05$).

Acute exposure to low doses of A β ₁₋₄₀ does not change K⁺ and Ca²⁺ fluxes from neurons


Inability to maintain Ca²⁺ homeostasis has been strongly implicated in AD [12-14]. Therefore, we investigated whether ion homeostasis would be affected by A β . We first examined whether acute exposure of mature neurons to low levels of a soluble A β ₁₋₄₀ affect changes in K⁺ and Ca²⁺ fluxes. We therefore performed simultaneous recordings of K⁺ and Ca²⁺ fluxes from cortical neurons using Microelectrode Ion Flux Estimation (MIFE) after an acute application of 1 μ MA β ₁₋₄₀ to primary 14 DIV neurons (Additional file 1, Figure S1). MIFE enables sensitive, real time measurement of ion fluxes across cell membranes in a variety of cell models and is non-invasive [17,18]. No changes in net K⁺ and Ca²⁺ fluxes were observed within 15 min post treatment



with 1 μM Aβ₁₋₄₀ that served as a control for our further experiments. We also found no changes in net K⁺ and Ca²⁺ fluxes in response to vehicle (Additional file 2, Figure S2) further confirming our findings.


Continuous exposure to Aβ₁₋₄₀ increases K⁺ efflux from neurons

Under physiological conditions neurons are continuously exposed to low doses of Aβ. To simulate those conditions we treated neurons with 1 μM Aβ over a prolonged period of time measuring ion fluxes of interest. Therefore, we investigated whether chronic treatment with soluble Aβ₁₋₄₀ can affect the ability of cortical neurons to maintain Ca²⁺ homeostasis (Figure 2). We also tested whether K⁺ homeostasis would be affected by Aβ treatments since fluxes of this ion are pivotal to neuronal activity. No difference was observed in the level of Ca²⁺ fluxes between untreated cells, and cells treated with Aβ₁₋₄₀ for up to eight days (Figure 2A). At the same time significant differences were observed in K⁺ fluxes (Figure 2B). While the K⁺ flux was similar in control cultures and those treated with Aβ₁₋₄₀ for one and three days, treatment with a soluble monomeric Aβ₁₋₄₀ (1 μM) daily for a period of six days or longer led to a significantly higher K⁺ efflux from the neural cells. Specifically, six and eight day Aβ treatment resulted in four- and three-fold increase in K⁺ efflux from the cells, respectively (P < 0.02) suggesting that chronic treatment with Aβ₁₋₄₀ leads to a reduced capacity of neurons to maintain K⁺ homeostasis.



Chronic exposure to Aβ₁₋₄₀ does not decrease neuronal viability following excitotoxic stress but does affect subsequent K⁺ homeostasis

We next investigated whether neurons exposed to prolonged Aβ₁₋₄₀ treatment had increased susceptibility to subsequent excitotoxicity induced by 200 μM glutamate. While glutamate treatment for 24 hours killed approximately 20% of neurons in the absence of any Aβ₁₋₄₀ treatment (Figure 3A), we found that seven days of pre-treatment with Aβ₁₋₄₀ did not further elevate neuronal susceptibility to glutamate-induced neurotoxicity (Figure 3B). We then examined K⁺ and Ca²⁺ fluxes in neurons which were treated with 1 μM Aβ₁₋₄₀ daily for a period from one to eight days and then challenged with 10 μM



of glutamate. Using MIFE, net fluxes of K⁺ and Ca²⁺ were monitored continuously throughout the experiment starting before the glutamate challenge and up to 30 min afterwards with data acquired every 6 sec. Typical examples of kinetics of net K⁺ and Ca²⁺ fluxes in response to challenge with glutamate are shown in Figures 4A-B and 5A-B, respectively.


Acute treatment of cortical neurons with 10 μM glutamate led to a dramatic K⁺ efflux from neurons that returned to pre-stress conditions within 20 min after the challenge (Figures 4A, B). Notably, the magnitude of K⁺ efflux was higher in cells treated with Aβ for six days or longer (peak values in the graphs). We therefore compared the magnitudes of the peak K⁺ efflux from neurons treated with Aβ with age-similar controls and found that K⁺ efflux was substantially (1.6-fold) higher even after one day of treatment with soluble Aβ₁₋₄₀ (Figure 6A). This difference was more pronounced with increase of duration of the treatment with Aβ leading to more than 3.5-fold increase in the peak K⁺ efflux after six and eight days of treatment with soluble monomeric Aβ₁₋₄₀ (1 μM), (Figure 6A).

We also assessed capacity of the cortical neurons to return K⁺ flux to pre-stress levels after glutamate challenge. For this we compared steady-state values of K⁺ fluxes 20 min after glutamate challenge in Aβ treated and age-similar un-treated control neurons (Figure 6B). While the level of K⁺ flux had returned to basal levels in non Aβ treated cells, this was not the case when neurons had received continual treatment with Aβ, suggesting that Aβ₁₋₄₀ treated neurons have reduced capacity to


maintain K⁺ homeostasis in response to glutamate stimulation. We also calculated total K⁺ efflux over 20 min after acute glutamate application shown as shaded area in Figure 4B. Total amount of K⁺ effluxed from the cells over 20 min post glutamate application was substantially higher after treatment with low concentration of Aβ for six days or longer. For example, it was 2.8-fold higher in cells pre-treated with Aβ for six days than in age-similar controls (22228.1 ± 2800.13 vs 8033.46 ± 3799.76 nmol m⁻², respectively, P < 0.05) further suggesting dysregulation of K⁺ homeostasis in Aβ treated cells.

Chronic Aβ treatment leads to increased Ca²⁺ uptake by cortical neurons in response to glutamate challenge

We similarly used MIFE to examine Ca²⁺ fluxes following glutamate challenge to control and Aβ treated cortical neurons. Neurons which had been exposed to glutamate challenge (10 μM) rapidly took up Ca²⁺ followed by a return to pre-stress levels (Figure 5A, B). In neurons which had been exposed to Aβ for one and three days, glutamate induced an influx of Ca²⁺ that was not statistically different from age-similar control cells (Figure 7A). However, cortical neurons that received soluble Aβ₁₋₄₀ for six consecutive days or longer had a significant impairment in their ability to maintain Ca²⁺ homeostasis, resulting in significantly greater Ca²⁺ uptake following excitotoxic stimulation with glutamate (Figure 7A). The magnitude of the peak Ca²⁺ influx was substantially higher in Aβ-treated cells than in the age-similar control cells (Figure 7A). At the same time




steady-state values of Ca^{2+} fluxes measured 20 min after the glutamate challenge were not statistically different from the relevant values of age-similar control cells even after six and eight days of $A\beta$ treatment, despite an apparent trend towards an increase in Ca^{2+} flux values (Figure 7B). However, a calculation of total Ca^{2+} uptake over 20 min of treatment with glutamate, showed that the amount of Ca^{2+} taken up by $A\beta$ treated cells was 2.5-fold higher than in age-similar un-treated control cells after six days of treatment with $A\beta$ (7964.47 ± 672.84 vs 3156.41 ± 1615.93 nmol m^{-2} , respectively, $P < 0.05$) further confirming Ca^{2+} dysregulation in cortical neurons after prolonged exposure to a soluble $A\beta$.



Discussion

Chronic exposure to $A\beta_{1-40}$ is neurotoxic to cultured cortical neurons


The accumulation of $A\beta$ protein in the brain is considered to be a key factor that causes AD. Most cultured cell models have examined the acute toxicity of $A\beta$, typically following 24 h treatment [6] which may not reflect the situation in vivo. In this study we examined the effect of daily exposure to moderate levels of soluble, monomeric $A\beta_{1-40}$ ($1 \mu\text{M}$) for up to seven days and demonstrated, for the first time, that chronic exposure of primary cortical neurons to levels of $A\beta$ which have no evident effect on viability over the short term, are indeed neurotoxic. A decrease in cell survival was only observed after seven days of exposure to $1 \mu\text{M}$ $A\beta$, with a 20% decrease in cell viability. This finding indicates that $A\beta_{1-40}$ accumulation over time can induce



neurodegenerative changes to neurons and thus contribute to the slowly progressing pathogenesis of AD.

Chronic exposure to Aβ₁₋₄₀ impairs K⁺ homeostasis in cortical neurons

Using MIFE, we observed an increase in K⁺ efflux concurrent with the decline in neuronal viability. While acute treatment with 1 μM Aβ did not affect net fluxes of K⁺ nor Ca²⁺ (Additional file 1: Figure S1) confirming earlier findings regarding intracellular Ca²⁺ changes [19], here we demonstrate that prolonged treatments with low concentrations of Aβ₁₋₄₀ led to dysregulations of K⁺ but not Ca²⁺ homeostasis. This is in agreement



with an earlier demonstration that exposure of neurons to Aβ peptide can enhance voltage gated K⁺ currents [20-22]. For example, preincubation of cells with recombinant human or rat Aβ₁₋₄₀ was shown to significantly increase K⁺ channel current density and levels for Kv4.2 [15]. Studies on brain slices at different stages of AD demonstrated that voltage gated K⁺ currents were already overexpressed in early stages of AD, and in advanced AD stages Kv3.4 was present at high levels in neurodegenerative structures [23]. Other studies also

demonstrated effects of A β on activity of Na⁺/K⁺-ATPase. For instance, it was shown that exposure of cultured neurons to A β ₁₋₄₀ caused selective reduction in Na⁺/K⁺-ATPase activity which preceded loss of Ca²⁺ homeostasis and cell degeneration [24]. This was further confirmed in studies using APP and PS1 deficient mice [25]. Together, enhancements of voltage gated K⁺ channels and decrease in Na⁺/K⁺-ATPase activity would increase overall K⁺ efflux, similar to A β effects observed in our experiments. The decrease in intracellular K⁺ concentration is also consistent with a decrease in cell viability since this is required for apoptosome formation and activation of caspases and endonucleases during apoptosis [26,27].

Exposure to glutamate further decreased impaired neuronal ability to maintain K⁺ homeostasis in cells chronically treated with A β ₁₋₄₀

A growing body of evidence suggests that perturbations in systems using the excitatory amino acid L-glutamate may underlie chronic neurodegenerative disorders such as AD. Therefore, we investigated whether prolonged exposure to 1 μ M A β ₁₋₄₀ increased neuronal susceptibility to subsequent acute excitotoxicity induced by glutamate. Surprisingly, no further decrease in neuronal viability was observed when cortical neurons treated with A β for seven days were exposed to glutamate (Figure 3A, B). Likewise, an earlier study has shown that low concentrations of A β peptide induced early and prolonged activation of proapoptotic markers in neurons without resulting in subsequent cell death in culture [28] most likely due to additional protective mechanisms in place at glutamate concentrations used. However, despite the absence of an overt change in neuronal viability following the combined treatments, we demonstrated that exposure to A β perturbed the ability of neurons to maintain K⁺ homeostasis following glutamate treatment. Treatments with 1 μ M A β for one day were sufficient to lead to glutamate induced K⁺ dyshomeostasis in cortical neurons (Figures 6A, B). Indeed, peak K⁺ efflux after acute exposure to 10 μ M glutamate was substantially higher in A β treated cells than in age-similar controls. This difference increased with time of A β exposure from 1.6 to 4.8-fold for one and six days of the treatment, respectively. Neurons were also unable to return K⁺ to pre-stress level as shown for steady-state K⁺ fluxes measured 20 minutes after the treatment with glutamate (Figure 6B). Another consequence of chronic pretreatment with A β was a higher level of K⁺ efflux even before challenge with glutamate (shown in Figure 2 for initial K⁺ values). Overall, this led to a 2.9-fold higher amount of K⁺ efflux from cells treated with A β ₁₋₄₀ for six days or longer after glutamate application than from age-similar control cells over 20 min of

glutamate exposure. Taken together, our data suggest that chronic A β treatment impairs the ability of cortical neurons to maintain K⁺ homeostasis.

Prolonged A β exposure increased Ca²⁺ uptake by cortical neurons in response to acute glutamate treatment

Ca²⁺ homeostasis undergoes subtle dysregulation during physiological ageing making neurons more vulnerable to additional stress which, in turn, can lead to neuronal degeneration. Recent evidence strongly supports the theory that dysregulation of intracellular Ca²⁺ homeostasis underlies the development of AD [12-14]. Therefore, we investigated whether prolonged exposure to continuous A β ₁₋₄₀ treatment followed by glutamate challenge would affect Ca²⁺ homeostasis in cortical neurons. Acute treatment with 10 μ M glutamate after chronic pre-treatment with 1 μ M A β ₁₋₄₀ in our experiments led to net Ca²⁺ uptake from external environment (Figure 5B). An increase in glutamate concentration leads to activation of ionotropic and metabotropic receptors [29,30] and Ca²⁺ influx through both glutamate-dependent pathways and voltage-dependent Ca²⁺ channels [31-33]. While the observed Ca²⁺ uptake in response to acute glutamate treatment was expected, in the current study we showed that the magnitude of Ca²⁺ uptake was substantially increased in cells treated with 1 μ M A β ₁₋₄₀ for six consecutive days or longer compared to un-treated control cells (Figure 7A). Calculated total Ca²⁺ taken up by A β treated cells over 20 min post glutamate application was 2.5-fold higher than in age-similar untreated control cells ($P < 0.05$) suggesting dysregulation of Ca²⁺ homeostasis in cortical neurons after prolonged exposure to soluble A β . Overall, our data showed that prolonged A β accumulation renders cortical neurons more vulnerable to excitotoxicity leading to their inability to maintain Ca²⁺ homeostasis when glutamate concentration increases.

Conclusions

In this study we developed an *in vitro* model to test the effects of prolonged A β treatment. We demonstrated for the first time that continuous exposure to low levels of A β ₁₋₄₀ induced neurodegeneration of cultured cortical neurons, with decreased viability evident after seven days of treatment. This decrease in viability correlated with increased neuronal K⁺ efflux. We also demonstrated that long term exposure to A β ₁₋₄₀ leads to imbalance in K⁺ and Ca²⁺ homeostasis and contributes to glutamate-induced vulnerability and cell death at later time point. This suggests that chronic accumulation of low levels of A β ₁₋₄₀ may contribute to the slowly progressing pathogenesis of AD. The results also suggest that the inability to maintain ion homeostasis is an early indicator of cell vulnerability that precedes cell death

and might be used as a sensitive tool to cell susceptibility to the excitotoxic treatment.

Methods

Rodent cortical neuron cultures

Cortical tissue was removed from embryonic day 18 Hooded Wistar rat embryos and incubated in sterile 10 mM HEPES buffer (37°C; Sigma). The cortical tissue was trypsinised (0.25%; Sigma) and washed with fresh HEPES buffer. The cell pellet was gently triturated and filtered through 60 µm gauze. Cells were then plated onto glass coverslips (254 mm²) pre-coated overnight with poly-L-lysine (Sigma) at the density of 1 × 10⁵ cells/well and culture medium added to the wells. The culture medium consisted of Neurobasal™ medium (Gibco), supplemented with 10% FCS (Gibco), 2% (f/c) B-27 supplement (Gibco), 5 mM (f/c) L-glutamine (Sigma), and 10 mg/ml gentamycin (Sigma). Cultures were maintained at 37°C in humidified air containing 5% CO₂. The culture medium was replaced with serum-free culture medium after 24 hours, followed by a full media change three times a week.

Aβ treatment and toxicity assay

1 µM Aβ₁₋₄₀ was applied to 6 DIV cortical neuron cultures daily for 7 days. Fresh media was applied a week. A biological reducing agent, 300 µM ascorbate (the physiological concentration within the brain), was freshly applied with each media change for both control and the treatment. At days 2, 3, 4, 6, and 7 the neuronal viability was measured using an Alamar Blue assay, which measures cellular metabolic activity [34]. The degree of cellular metabolic reduction of Alamar Blue was determined by fluorescence (excitation 535 nm, emission 595 nm), and expressed as the percentage of the signal obtained from the vehicle-treated culture. Day zero was the first day of treatment

Neural survival following chronic Aβ treatment

To examine whether Aβ treatment increased neuron susceptibility to excitotoxicity, 6 DIV cultures treated daily with 1 µM of soluble Aβ₁₋₄₀ for seven days were exposed to 200 µM glutamate. Glutamate was prepared as we have reported previously [35]. Briefly, D-glutamic acid was dissolved in appropriate medium (Neurobasal medium for viability assay and aCSF for MIFE experiments). Neuronal viability was measured using the Alamar Blue assay after 24 hours of glutamate treatment.

Ion-selective flux measurements

The technique of Microelectrode ion flux estimation (MIFE) has been reviewed recently [18] and the complete experimental procedure including ion-selective microelectrode fabrication and cell preparation and

immobilisation are given elsewhere [17,18]. Cortical neurons for the MIFE measurements were grown for 6 days at 1 × 10⁵ cells/well on coverslips coated with poly-L-lysine as described above. By 6 DIV a dense monolayer of neurons had developed. Cells were washed in artificial CSF (aCSF) by dipping a coverslip with cells into a beaker with pre-warmed to 37°C aCMF. The coverslip then was placed into a measuring chamber with 2 mL of aCMF at 37°C, and left for adaptation for a further 40 min at room temperature (RT, 23°C) prior to an experiment. The composition of the aCSF was: 150 mM NaCl, 0.5 mM KCl, 0.5 mM CaCl₂, 1.5 mM MgCl₂, 1.25 mM NaH₂PO₄, 2 mM NaHCO₃, 25 mM glucose, pH 7.2. Measurements were taken after one, three, six, and eight days post treatment with Aβ₁₋₄₀. Control experiments used age similar cells that were not pre-treated with Aβ₁₋₄₀. Additional controls such as addition of vehicle (aCMF) and Aβ₁₋₄₀ were also used. For this co-focused K⁺ and Ca²⁺ microelectrodes were positioned ~5 µm above the cells and moved by a computer-driven hydraulic micromanipulator by 50 µm away from the cells and back to the primary position with 0.05 Hz frequency. Data was acquired at a rate of 10 samples/sec and averaged later over 6 second intervals. The ion fluxes were recorded for 5 min prior to the addition of 10 µM of glutamate, and recordings continued for a further 25 min. To achieve the required concentration of glutamate in the measuring chamber, an equal volume of the bath solution with a double concentration of glutamate was added resulting in the final concentrations of glutamate (10 µM). Net ion fluxes (nmol m⁻²s⁻¹) were calculated using planar geometry of diffusion equations.

Statistical analyses of tissue culture experiments

For each experiment unless otherwise stated, a minimum of four wells from at least three separate cultures (derived from different animals), were used for quantification. Statistical analysis was completed using SPSS 16.0 (SPSS). When data was unequally distributed, data was transformed so that the residuals were approximately normally distributed. Statistical significance was calculated using One-Way ANOVA with Tukey's Post Hoc Test and Student's *t*-test.

Additional material

Additional file 1: Figure S1. Effects of acute application of Aβ₁₋₄₀ on net K⁺ and Ca²⁺ fluxes. We tested whether acute application of Aβ₁₋₄₀ to 14 DIV cortical neurons affected magnitudes of K⁺ and Ca²⁺ fluxes. Net fluxes of K⁺ (A) and Ca²⁺ (B) were recorded for 5 min (-5 to 0 min) followed by acute application of 1 µM Aβ₁₋₄₀ (0 to 10 min) and 40 µM Aβ₁₋₄₀ (final concentration, 10 to 25 min) to the bath. Neither concentration caused changes in net ion fluxes measured during the time course tested suggesting that prolonged treatment with Aβ₁₋₄₀ is required to trigger disturbances in ion homeostasis. Error bars are SEM (n = 4).

Additional file 2: Figure S2. Effects of experimental solution application to the measuring chamber on ion fluxes.

An additional control was made to test a possible effect of solution disturbances in the measuring chamber on ion fluxes. Neurons at 14 DIV were used. Net K^+ (A) and Ca^{2+} (B) fluxes were recorded continuously for 5 min (-5 to 0 min) and after vehicle (aCSF) application to the bath (0 to 10 min) with data acquired at a rate of 10 samples/sec and averaged over every 6 sec. Vehicle was applied at zero time as indicated by an arrow. No changes in K^+ and Ca^{2+} fluxes were observed thus validating the approach used. Error bars are SEM (n = 4).

Abbreviations

A β : beta amyloid; AD: Alzheimer's disease; APP: β -amyloid precursor peptide; DIV: days *in vitro*; MIFE: microelectrode ion flux measuring technique.

Acknowledgements

This work was supported by the National Health and Medical Research Council of Australia (#490025, 605528) to AKW and RSC, a Jack and Ethel Goldin Foundation Award (Alzheimer's Australia) to AKW and RSC, a Rising Stars Award from the University of Tasmania to RSC, and a research grant from the Australian Brain Foundation to LS. RSC holds a Research Fellowship from the Australian Research Council.

Authors' contributions

LS carried out MIFE experiments, analysed and interpreted data, and drafted the manuscript. CH carried out the survival experiments and participated in data analyses. AKW participated in the design of the study, participated in its coordination and contributed to the preparation of the manuscript. RSC conceived the study, participated in overall direction of the study and preparation of the manuscript. All authors have read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 6 May 2010 Accepted: 13 August 2010

Published: 13 August 2010

References

1. Blennow K, de Leon MJ, Zetterberg H: **Alzheimer's disease.** *Lancet* 2006, **368**:387-403.
2. Selkoe DJ, Podlisny MB: **Deciphering the genetic basis of Alzheimer's disease.** *Annu Rev Genomics Hum Genet* 2002, **3**:67-99.
3. Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, Yang F, Cole G: **Correlative memory deficits, A β elevation, and amyloid plaques in transgenic mice.** *Science* 1996, **274**:99-102.
4. Koistinaho M, Ort M, Cimadevilla JM, Vondrous R, Cordell B, Koistinaho J, Bures J, Higgins LS: **Specific spatial learning deficits become severe with age in beta -amyloid precursor protein transgenic mice that harbor diffuse beta -amyloid deposits but do not form plaques.** *Proc Natl Acad Sci USA* 2001, **98**:14675-14680.
5. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL: **Diffusible, nonfibrillar ligands derived from A β 1-42 are potent central nervous system neurotoxins.** *Proc Natl Acad Sci USA* 1998, **95**:6448-6453.
6. Deshpande A, Mina E, Glabe C, Busciglio J: **Different conformations of amyloid beta induce neurotoxicity by distinct mechanisms in human cortical neurons.** *J Neurosci* 2006, **26**:6011-6018.
7. Supnet C, Bezprozvanny I: **The dysregulation of intracellular calcium in Alzheimer disease.** *Cell Calcium* 2010, **47**:183-189.
8. Parameshwaran K, Dhanasekaran M, Suppiramaniam V: **Amyloid beta peptides and glutamatergic synaptic dysregulation.** *Exp Neurol* 2008, **210**:7-13.
9. Choi DW: **Glutamate neurotoxicity and diseases of the nervous system.** *Neuron* 1988, **1**:623-634.
10. Smythies J: **The neurotoxicity of glutamate, dopamine, iron and reactive oxygen species: functional interrelationships in health and disease: a review-discussion.** *Neurotox Res* 1999, **1**:27-39.
11. Sattler R, Tymianski M: **Molecular mechanisms of calcium dependent excitotoxicity.** *J Mol Med* 2000, **78**:3-13.
12. Yu JT, Chang RC, Tan L: **Calcium dysregulation in Alzheimer's disease: from mechanisms to therapeutic opportunities.** *Prog Neurobiol* 2009, **89**:240-255.
13. Conley YP, Mukherjee A, Kammerer C, DeKosky ST, Kamboh MI, Finegold DN, Ferrell RE: **Evidence supporting a role for the calcium-sensing receptor in Alzheimer disease.** *Neuron* 2008, **59**:190-194.
14. Bojarski L, Herms J, Kuznicki J: **Calcium dysregulation in Alzheimer's disease.** *Neurochem Int* 2008, **52**:621-633.
15. Kerrigan TL, Atkinson L, Peers C, Pearson HA: **Modulation of 'A'-type K^+ current by rodent and human forms of amyloid beta protein.** *Neuroreport* 2008, **19**:839-843.
16. Mulholland PJ, Carpenter-Hyland EP, Hearing MC, Becker HC, Woodward JJ, Chandler LJ: **Glutamate transporters regulate extrasynaptic NMDA receptor modulation of Kv2.1 potassium channels.** *J Neurosci* 2008, **28**:8801-8809.
17. Shabala L, Ross T, Newman I, McMeekin T, Shabala S: **Measurements of net fluxes and extracellular changes of H^+ , Ca^{2+} , K^+ , and NH_4^+ in *Escherichia coli* using ion-selective microelectrodes.** *J Microbiol Meth* 2001, **46**:119-129.
18. Shabala L, Ross T, McMeekin T, Shabala S: **Non-invasive microelectrode ion flux measurements to study adaptive responses of microorganisms to the environment.** *FEMS Microbiol Rev* 2006, **30**:472-486.
19. Chin JH, Tse FW, Harris K, Jhamandas JH: **Beta-amyloid enhances intracellular calcium rises mediated by repeated activation of intracellular calcium stores and nicotinic receptors in acutely dissociated rat basal forebrain neurons.** *Brain Cell Biol* 2006, **35**:173-86.
20. Pannaccione A, Secondo A, Scorziello A, Cali G, Tagliatalata M, Annunziato L: **Nuclear factor-kappaB activation by reactive oxygen species mediates voltage-gated K^+ current enhancement by neurotoxic beta-amyloid peptides in nerve growth factor-differentiated PC-12 cells and hippocampal neurones.** *J Neurochem* 2005, **94**:572-586.
21. Pannaccione A, Boscia F, Scorziello A, Adornetto A, Castaldo P, Sirabella R, Tagliatalata M, Di Renzo GF, Annunziato L: **Up-regulation and increased activity of KV3.4 channels and their accessory subunit Mink-related peptide 2 induced by amyloid peptide are involved in apoptotic neuronal death.** *Mol Pharmacol* 2007, **72**:499-501.
22. Choi E, Abbott GW: **The MiRP2-Kv3.4 potassium channel: muscling in on Alzheimer's disease.** *Mol Pharmacol* 2007, **72**:499-501.
23. Angulo E, Noé V, Casadó V, Mallol J, Gomez-Isla T, Lluís C, Ferrer I, Ciudad CJ, Franco R: **Up-regulation of the Kv3.4 potassium channel subunit in early stages of Alzheimer's disease.** *J Neurochem* 2004, **91**:547-557.
24. Mark RJ, Hensley K, Butterfield DA, Mattson MP: **Amyloid beta-peptide impairs ion-motive ATPase activities: evidence for a role in loss of neuronal Ca^{2+} homeostasis and cell death.** *J Neurosci* 1995, **15**:6239-6249.
25. Dickey CA, Gordon MN, Wilcock DM, Herber DL, Freeman MJ, Morgan D: **Dysregulation of Na^+/K^+ -ATPase by amyloid in APP+PS1 transgenic mice.** *BMC Neurosci* 2005, **6**:7.
26. Xiao AY, Homma M, Wang XQ, Wang X, Yu SP: **Role of K^+ efflux in apoptosis induced by AMPA and kainate in mouse cortical neurons.** *Neurosci* 2001, **108**:61-67.
27. Zhao YM, Sun LN, Zhou HY, Wang XL: **Voltage-dependent potassium channels are involved in glutamate-induced apoptosis of rat hippocampal neurons.** *Neurosci Lett* 2006, **398**:22-27.
28. White AR, Guirguis R, Brazier MW, Jobling MF, Hill AF, Beyreuther K, Barrow CJ, Masters CL, Collins SJ, Cappai R: **Sublethal concentrations of prion peptide PrP106-126 or the amyloid beta peptide of Alzheimer's disease activates expression of proapoptotic markers in primary cortical neurons.** *Neurobiol Dis* 2001, **8**:299-316.
29. Perrella J, Bhavnani BR: **Protection of cortical cells by equine estrogens against glutamate-induced excitotoxicity is mediated through a calcium independent mechanism.** *BMC Neurosci* 2005, **6**:34.
30. Lei Z, Deng P, Xu ZC: **Regulation of Kv4.2 channels by glutamate in cultured hippocampal neurons.** *J Neurochem* 2008, **106**:182-192.
31. Frick A, Magee J, Johnston D: **LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites.** *Nat Neurosci* 2004, **7**:126-135.

32. Lisman J, Spruston N: Postsynaptic depolarization requirements for LTP and LTD: a critique of spike timing-dependent plasticity. *Nat Neurosci* 2005, **8**:839-841.
33. Magee JC, Johnston D: A synaptically controlled, associative signal for Hebbian plasticity in hippocampal neurons. *Science* 1997, **275**:209-213.
34. White MJ, DiCaprio MJ, Greenberg DA: Assessment of neuronal viability with Alamar Blue in cortical and granule cell cultures. *J Neurosci Methods* 1996, **70**:195-200.
35. Chung RS, McCormack GH, King AE, West AK, Vickers JC: Glutamate induces rapid loss of axonal neurofilament proteins from cortical neurons in vitro. *Exp Neurol* 2005, **193**(2):481-8.

doi:10.1186/1750-1326-5-30

Cite this article as: Shabala *et al.*: Prolonged A β treatment leads to impairment in the ability of primary cortical neurons to maintain K⁺ and Ca²⁺ homeostasis. *Molecular Neurodegeneration* 2010 **5**:30.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

