

CASE REPORT

Open Access

Temporary cognitive impairment related to administration of newly developed anticholinergic medicines for overactive bladder: two case reports

Takako Shiota¹, Kazumasa Torimoto^{2*}, Hitoshi Momose³, Takuya Nakamuro⁴, Hiroshi Mochizuki¹, Hiromi Kumamoto¹, Akihide Hirayama² and Kiyohide Fujimoto²

Abstract

Background: Cognitive impairment is one of the side effects of using anticholinergic medicines for overactive bladder; however, its incidence has not been fully reported. We experienced two elderly Japanese patients with overactive bladder who had temporary cognitive impairment caused by anticholinergic medicines.

Case presentation: The first case was a 79-year-old female patient to whom imidafenacin (0.2 mg) was administered daily to control her frequent micturition and urgency. She was taking the following medicines: etizolam, triazolam, captopril, bisoprolol, and amlodipine besylate. Her Hasegawa dementia rating scale-revised was impaired from 26/30 to 17/30 and recovered to 25/30 after the imidafenacin treatment was stopped. The second case was an 82-year-old female patient to whom imidafenacin (0.2 mg) was administered daily for frequent micturition and urgency. She was taking the following medicines: losartan potassium and clenbuterol. Her Hasegawa dementia rating scale-revised decreased from 28/30 to 19/30 and recovered to 24/30 after the imidafenacin treatment was stopped. In our patients who were taking multiple medicines, there is a possibility that medicines other than anticholinergics may have caused cognitive impairment. We need to keep in mind that many elderly people take multiple medicines because of comorbidity.

Conclusions: Anticholinergic medicines can cause cognitive impairment in elderly people, and attention should be paid to cognition when elderly overactive bladder patients are treated with anticholinergic medicines.

Keywords: Overactive bladder, Anticholinergic medicine, Cognition

Background

Cognitive impairment is one of the side effects of using anticholinergic medicines for overactive bladder (OAB); however, its incidence has not been fully reported. Oxybutynin has been shown to cause cognitive impairment and delirium in elderly people [1]. However, no other report of cognitive impairment caused by anticholinergic medicines for OAB was found. Recently, newly developed anticholinergic medicines for OAB were found to not affect cognition [2,3]. These newly developed anticholinergic medicines for OAB are thought to be safe and not cause cognitive impairment because their movement

into the central nervous system is limited [4]. We experienced two elderly Japanese patients with OAB who had temporary cognitive impairment caused by anticholinergic medicines.

Case presentation

Case 1

A 79-year-old female patient presented to a neurologist with a chief complaint of amnesia six months ago and was found to have normal cognition, as a computed tomography (CT) scan did not detect any organic brain lesions and Hasegawa dementia rating scale-revised (HDS-R) [5] was 26/30. She was taking the following medicines: etizolam, triazolam, captopril, bisoprolol, and amlodipine besylate.

* Correspondence: torimoto@naramed-u.ac.jp

²Department of Urology, Nara Medical University, 840 Shijo-cho, Kashihara-shi, Nara, Japan

Full list of author information is available at the end of the article

Imidafenacin (0.2 mg) was administered daily to control her frequent micturition and urgency. After one month, the imidafenacin treatment was stopped because her cognition was impaired (HDS-R, 17/30), although the lower urinary tract symptoms (LUTS) improved. A CT scan did not detect any brain lesions at that time. After three weeks, her cognition recovered (HDS-R, 25/30). Then, tolterodine (4 mg) was administered daily instead of imidafenacin; however, it was stopped because of cognitive impairment (Figure 1).


Case 2


An 82-year-old female patient presented with a chief complaint of mixed urinary incontinence that had persisted for four months. She was taking losartan potassium. Clenbuterol (40 µg) was administered daily because stress urinary incontinence was predominant over urge urinary incontinence. After one month, her urinary incontinence improved. Three months later, imidafenacin (0.2 mg) was administered daily for frequent micturition and urgency. However, after two weeks of treatment, her HDS-R decreased from 28/30 to 19/30, although the LUTS improved. At that time, she was found to have normal cognition by a neurologist because her mini mental scale examination (MMSE) was 26/30 and a CT scan did not detect any brain lesions. Imidafenacin administration was immediately stopped, and three weeks later, her HDS-R recovered to 24/30, although the LUTS got worse. Then,

mirabegron (50 mg) was administered daily instead of imidafenacin. Her cognition was stable as follows: her HDS-R was 25/30 after one month, and her HDS-R and MMSE were 26/30 and 23/30, respectively, after seven months (Figure 2).

Discussion

Anticholinergics are the first choice for OAB and are often administered to elderly people in whom the prevalence of OAB is high [6]. While it has been reported that anticholinergic medicines for OAB are safe and do not cause cognitive impairment [4], anticholinergics do in fact frequently induce reversible cognitive impairment [7,8]. They were traditionally thought to affect cognition and be high-risk medicines causing cognitive impairment. There is good evidence that the combination of multiple medicines can precipitate or predispose to cognitive impairment. The American College of Surgeons National Surgical Quality Improvement Program (ACS NSQIP®)/American Geriatrics Society (AGS) Best Practices Guidelines list strong anticholinergic medicines for OAB, such as darifenacin, fesoterodine, flavoxate, oxybutynin, solifenacin, tolterodine, and trospium, which may be avoided in elderly people [9]. The adverse effects of anticholinergic medicines on cognition depend on the strength or combination. It has been reported that anticholinergic medicines had a significantly high risk of causing cognitive impairment and dementia after adjusting for other possible





determinants of cognitive impairment [7,10]. In our patients who were taking multiple medicines, there is a possibility that medicines other than anticholinergics may have caused cognitive impairment. We need to keep in mind that many elderly people take multiple medicines because of comorbidity.

Cystometry has become unnecessary to diagnose patients as OAB since the Standardization Subcommittee of the International Continence Society (ICS) recognized OAB as a “symptom syndrome suggestive of lower urinary tract dysfunction” which is specifically defined as “urgency, with or without urge incontinence, usually with frequency and nocturia” in 2002 [11]. Afterwards any medical doctors including primary care doctors can diagnose patients as OAB and treat OAB mainly with anticholinergic medicines. We may have to perform cystometry more aggressively to detect detrusor overactivity in elderly patients who have the risk of drug-induced cognitive impairment. That may result in better selection of elderly patients and allow doctors to avoid side effects of anticholinergic medicines.

While MMSE is the global standard diagnostic method for cognition, HDS-R is the standard method in Japan. In our second case (Case 2), the diagnostic results were different between MMSE and HDS-R. It is still debatable which method is better for the diagnosis of cognitive impairment [5].

Conclusions

Anticholinergic medicines can cause cognitive impairment in elderly people, and attention should be paid to cognition when elderly OAB patients are treated with anticholinergic medicines.

Consent

Written informed consent was obtained from the patients for publication of these Case Reports and any accompanying images. Copies of the written consent are available for review by the Editor-in-Chief of this journal.

Abbreviations

OAB: Overactive bladder; HDS-R: Hasegawa dementia rating scale-revised; CT: Computed tomography; LUTS: Lower urinary tract symptoms; MMSE: Mini mental scale examination.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HM, AH and KF have made substantial contributions to conception and design. TS, TN, HM and HK have made substantial contributions to acquisition of data. TS and KT have been involved in drafting the manuscript and revising it critically for important intellectual content. All authors have given final approval of the version to be published.

Acknowledgements

Nobumichi Tanaka has made contribution to acquisition of data.

Author details

¹Department of Urology, Saiseikai Nara Hospital, 4-643 Hachijo, Nara-shi, Nara, Japan. ²Department of Urology, Nara Medical University, 840 Shijo-cho, Kashihara-shi, Nara, Japan. ³Department of Urology, Hoshigaoka Medical Center, 4-8-1 Hoshigaoka, Hirakata-shi, Osaka, Japan. ⁴Department of Neurology, Saiseikai Nara Hospital, 4-643 Hachijo, Nara-shi, Nara, Japan.

Received: 5 February 2014 Accepted: 20 September 2014

Published: 25 September 2014

References

1. Katz IR, Sands LP, Bilker W, DiFilippo S, Boyce A, D'Angelo K: **Identification of medications that cause cognitive impairment in older people: the case of oxybutynin chloride.** *J Am Geriatr Soc* 1998, **46**(1):8–13.
2. Kessler TM, Bachmann LM, Minder C, Lohrer D, Umbehr M, Schunemann HJ, Kessels AG: **Adverse event assessment of antimuscarinics for treating overactive bladder: a network meta-analytic approach.** *PLoS One* 2011, **6**(2):e16718.
3. Griebling TL: **Re: Anticholinergic medication use and cognitive impairment in the older population: the Medical Research Council Cognitive Function and Ageing Study.** *J Urol* 2012, **187**(4):1357–1358.
4. Pagoria D, O'Connor RC, Guralnick ML: **Antimuscarinic drugs: review of the cognitive impact when used to treat overactive bladder in elderly patients.** *Curr Urol Rep* 2011, **12**(5):351–357.
5. Kim KW, Lee DY, Jhoo JH, Youn JC, Suh YJ, Jun YH, Seo EH, Woo JI: **Diagnostic accuracy of mini-mental status examination and revised hasegawa dementia scale for Alzheimer's disease.** *Dement Geriatr Cogn Disord* 2005, **19**(5–6):324–330.
6. Homma Y, Yamaguchi O, Hayashi K: **An epidemiological survey of overactive bladder symptoms in Japan.** *BJU Int* 2005, **96**(9):1314–1318.
7. Moore AR, O'Keefe ST: **Drug-induced cognitive impairment in the elderly.** *Drugs Aging* 1999, **15**(1):15–28.
8. Weytingh MD, Bossuyt PM, van Crevel H: **Reversible dementia: more than 10% or less than 1%? A quantitative review.** *J Neurol* 1995, **242**(7):466–471.
9. American Geriatrics Society Beers Criteria Update Expert P: **American Geriatrics Society updated Beers Criteria for potentially inappropriate medication use in older adults.** *J Am Geriatr Soc* 2012, **60**(4):616–631.
10. Fox C, Richardson K, Maidment ID, Savva GM, Matthews FE, Smithard D, Coulton S, Katona C, Boustani MA, Brayne C: **Anticholinergic medication use and cognitive impairment in the older population: the medical research council cognitive function and ageing study.** *J Am Geriatr Soc* 2011, **59**(8):1477–1483.
11. Wein AJ, Rovner ES: **Definition and epidemiology of overactive bladder.** *Urology* 2002, **60**(5 Suppl 1):7–12. discussion 12.

doi:10.1186/1756-0500-7-672

Cite this article as: Shiota et al.: Temporary cognitive impairment related to administration of newly developed anticholinergic medicines for overactive bladder: two case reports. *BMC Research Notes* 2014 **7**:672.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

