

RESEARCH

Open Access



Associations of the COVID-19 pandemic with the reported incidence of important endemic infectious disease agents and syndromes in Pakistan

Bayan Missaghi^{1,2*} , Muhammad Wasif Malik³, Waseem Shaukat⁴, Muazam Abbas Ranjha³, Aamer Ikram⁵ and Herman W. Barkema^{4,6,7}

Abstract

Background: Persons in Pakistan have suffered from various infectious diseases over the years, each impacted by various factors including climate change, seasonality, geopolitics, and resource availability. The COVID-19 pandemic is another complicating factor, with changes in the reported incidence of endemic infectious diseases and related syndromes under surveillance.

Methods: We assessed the monthly incidence of eight important infectious diseases/syndromes: acute upper respiratory infection (AURI), viral hepatitis, malaria, pneumonia, diarrhea, typhoid fever, measles, and neonatal tetanus (NNT), before and after the onset of the COVID-19 pandemic. Administrative health data of monthly reported cases of these diseases/syndromes from all five provinces/regions of Pakistan for a 3-year interval (March 2018–February 2021) were analyzed using an interrupted time series approach. Reported monthly incidence for each infectious disease agent or syndrome and COVID-19 were subjected to time series visualization. Spearman's rank correlation coefficient between each infectious disease/syndrome and COVID-19 was calculated and median case numbers of each disease before and after the onset of the COVID-19 pandemic were compared using a Wilcoxon signed-rank test. Subsequently, a generalized linear negative binomial regression model was developed to determine the association between reported cases of each disease and COVID-19.

Results: In late February 2020, concurrent with the start of COVID-19, in all provinces, there were decreases in the reported incidence of the following diseases: AURI, pneumonia, hepatitis, diarrhea, typhoid, and measles. In contrast, the incidence of COVID was negatively associated with the reported incidence of NNT only in Punjab and Sindh, but not in Khyber Pakhtunkhwa (KPK), Balochistan, or Azad Jammu & Kashmir (AJK) & Gilgit Baltistan (GB). Similarly, COVID-19 was associated with a lowered incidence of malaria in Punjab, Sindh, and AJK & GB, but not in KPK and Balochistan.

Conclusions: COVID-19 was associated with a decreased reported incidence of most infectious diseases/syndromes studied in most provinces of Pakistan. However, exceptions included NNT in KPK, Balochistan and AJK & GB, and

*Correspondence: bayan.missaghi@albertahealthservices.ca

¹ Department of Medicine, Cumming School of Medicine, University of Calgary and Alberta Health Services, Office 3683, 3500 – 26 Ave NE, Calgary, AB T1Y 6J4, Canada

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

malaria in KPK and Balochistan. This general trend was attributed to a combination of resource diversion, misdiagnosis, misclassification, misinformation, and seasonal patterns of each disease.

Keywords: COVID-19, SARS-CoV-2, Infectious disease outbreaks, Incidence

Background

Over the years, persons in Pakistan have experienced a wide variety of infectious diseases such as Dengue fever, Crimean-Congo Hemorrhagic fever (CCHF), viral hepatitis, measles, and polio [1]. The diversity of healthcare challenges over time has been exacerbated by many outbreaks of endemic transmissible pathogens, with devastating consequences both from a morbidity and mortality perspective [2], as well as from the standpoint of healthcare costs [3–5]. Unfortunately, as is the case with many developing nations with struggling healthcare sectors, Pakistan has also been quite vulnerable to effects of the novel coronavirus 2019 (COVID-19) pandemic [6], caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Government estimates state that, as of December 1, 2021, the current pandemic had infected more than 1.25 million Pakistanis and taken more than 28,000 lives [7]. Globally, as of the same date, the pandemic had involved 222 countries, infected > 250,000,000 people, and taken more than 5 million lives [8].

The incidence of ongoing non-COVID-19 infectious diseases endemic in Pakistan is influenced by a complex set of evolving circumstances, involving factors such as climate change, global warming, seasonality, loss of biodiversity, changes in vector geography, and many other ecological determinants [1]. Moreover, Pakistan's attempts at control of various outbreaks have suffered from geopolitical factors, misinformation, and a shortage of resources, as exemplified by the ongoing polio outbreak in the tribal-controlled regions of the country [9, 10]. In this context, isolating impacts of any one infectious disease outbreak on another is neither simple nor straightforward.

Nevertheless, despite the inherent complexity of the causes and interplay between determinants, it has been postulated that there have been notable impacts of COVID-19 on the incidence and/or prevalence of several infectious diseases in Pakistan, including arboviral diseases, influenza A virus subtype H1N1, Dengue fever, Crimean-Congo hemorrhagic fever, tuberculosis, malaria, measles, typhoid fever, etc. [11–16]. Some of these impacts have been attributed to under-reporting and a diversion of resources from the diagnosis and management of ongoing endemic pathogens such as measles and polio to the diagnosis and surveillance of COVID-19, whereas others have been proposed to be due to

possible exacerbation of these ongoing other outbreaks by COVID-19 (in a synergistic fashion) and/or over-reporting thereof due to an overlap in symptoms. Early on in the pandemic, there were also efforts made by the National Command and Operations Centre (NCOC) of the Government of Pakistan to develop standard operating procedures that were meant to be strictly followed in public areas, including guidelines on physical distancing, wearing masks, maintaining hand hygiene, and implementing lockdowns [17]. Consequently, it is likely that these efforts had an inadvertent impact on lowering the incidence of other transmissible infections as well, particularly those spread through a respiratory route. Regardless of the specific etiologies, several Pakistani scientists and clinicians have observed anecdotal alterations in the incidence rates of various infectious diseases currently under surveillance throughout the country. However, the directionality of these changes and their degree have not been comprehensively described. Therefore, the objectives were to assess the impact of the COVID-19 pandemic on the reported incidence of endemic infectious diseases/syndromes in Pakistan, and to discuss potential underlying reasons for any such impact.

Methods

We received permission from the Ministry of National Health Services Regulations and Coordination (MoNHSR&C) of the Government of Pakistan to obtain monthly incident infectious disease data for the following pathogens and infectious disease syndromes: acute upper respiratory infection, viral hepatitis (primarily hepatitis A and E), malaria (undifferentiated species), pneumonia (undifferentiated), diarrhea (undifferentiated), typhoid fever, measles, neonatal tetanus and COVID-19 for 3 years (March 2018–February 2021) for all five provinces/regions of Pakistan—namely Punjab, Sindh, Khyber Pakhtunkhwa (KPK), Balochistan, and Azad Jammu & Kashmir (AJK) & Gilgit Baltistan (GB). This interval was from 24 months before to 12 months after the onset of COVID-19 in Pakistan that occurred at the end of February 2020. Data regarding the monthly incidence of COVID-19 were collected from the time of the first case in late February 2020 through February 2021.

Case definitions of the eight diseases discussed in the present study were adopted from the Pakistan List of

Priority Diseases for Implementation of the Integrated Disease Surveillance Response System (IDSR), as follows:

1. Acute upper respiratory infection (AURI)—Acute upper respiratory infection with cough, rhinorrhea or sore throat in the absence of clinical or x-ray evidence of pneumonic involvement.
2. Pneumonia—Cough or difficulty breathing with fever, chills, chest pain and clinical or x-ray evidence of pneumonic consolidation.
3. Viral hepatitis (A & E)—Acute onset of jaundice < 1 month duration and severe illness (dark urine, fatigue, nausea, vomiting, and abdominal pain) with absence of any known precipitating factors (other than contaminated food or water).
4. Diarrhea (Non-Cholera)—An illness characterized by ≥ 3 loose, non-bloody stools over a 24-h interval, with or without dehydration.
5. Typhoid fever—Acute febrile illness, fever of ≥ 38 °C for ≥ 3 days with abdominal discomfort, fatigue and diarrhea/constipation.
6. Measles—Any person in whom a clinician suspects measles infection OR any person with fever AND any 2 of the following signs/symptoms: maculopapular rash (non-vesicular), cough, coryza, or conjunctivitis.
7. Neonatal tetanus (NNT)—Any neonate between 3 and 28 days of age with, in the absence of a more likely diagnosis, an acute illness with muscle spasms or hypertonia and a clinical diagnosis of tetanus by a health care provider, OR death from neonatal tetanus listed on the death certificate as the cause of death or a significant condition contributing to death.
8. Malaria (undifferentiated species)—Any person with fever or history of fever within the past 48 h (with or without other symptoms such as nausea, vomiting and diarrhea, headache, back pain, chills, or myalgia) in whom other obvious causes of fever have been excluded.
9. COVID-19—Acute onset of any 3 or more of the following signs or symptoms: fever, dry cough, general weakness/fatigue, headache, myalgia, sore throat, coryza, dyspnea, anorexia/nausea/vomiting and diarrhea, particularly in the context of having exposure to a known positive case or residing in a high-transmission area within 14 d prior to the onset of symptoms.

The entire population of Pakistan, including persons of all age groups, sexes, educational status, socio-economic status, and residence, was included in this study. Non-residents of Pakistan were excluded. A standardized tool used by the MoNHSR&C for the collection of aggregated data for their Demographic Health Information System (DHIS) was adopted, specifically targeting

eight priority infectious diseases/syndromes in the country in addition to COVID-19. Of note, the DHIS works at the national level in Pakistan under the administrative control of the MoNHSR&C, Government of Pakistan. All districts across Pakistan report data monthly to the DHIS for various diseases. Specifically, data flowed from 17,003 data reporting sites, including health facilities, up to the district level, and then it was subsequently combined at a provincial level and transmitted to the National Health Ministry database.

Data Analyses

Statistical analyses were performed using Stata/SE 17.0 for Mac (StataCorp. 2021. Stata Statistical Software: Release 17. College Station, TX: StataCorp LLC). The reported incidence of all eight infectious diseases before and after the onset of the COVID-19 pandemic in Pakistan was analyzed using an interrupted time series approach. Reported monthly incidence of each infectious disease pathogen or syndrome was assessed for normality using histograms and the Shapiro–Wilk test. These data were non-normally distributed, and, therefore, case counts were reported as medians along with interquartile range (IQR). Time series line plots were created for each disease and province, respectively, representing monthly case numbers along with COVID-19 case numbers to visually compare the cases in each of the 3 years in the dataset, including 2 years before (March 2018–February 2020) and 1 year after the onset of COVID-19 (March 2020–February 2021). Subsequently, the Spearman's correlation coefficient (ρ) between each separate infectious disease pathogen or syndrome and COVID-19 was calculated using a two-tailed $\alpha < 0.05$ defined as meeting significance. To determine whether the number of monthly cases for any disease differed between the pre-COVID and COVID period, a Wilcoxon signed-rank test was used to compare monthly averages of the pre-COVID period with the COVID period, considering the seasonal variations between months in a year. Boxplots were created to visually assess the distribution of cases between pre-COVID and COVID periods for each of the infectious diseases/syndromes in each province/region.

Thereafter, in view of overdispersion in these count data, a generalized linear negative binomial regression model with log link was developed, which allowed adequate handling of over-dispersed count data to determine associations between COVID-19 and each of the infectious disease pathogens or syndromes separately for each province. To account for seasonal trends in the data, monthly indicator variables were included in the final models. The final fitted model that we report in this

study, separately for each disease and province, was this equation:

$$\log(E(Disease)) = \beta_0 + \beta_1 \cdot (COVID\ cases\ in\ Thousands) + \sum_{i=1}^{12} \beta_i \cdot I_{Month_i}$$

where E(Disease) is the expected monthly number of the cases of disease for which the model is fitted, β_0 is the log of the expected count of disease cases when COVID-19 cases were 0 for the month of January as the base, β_1 is the difference in the log of the expected count of disease cases with every 1000 increase in COVID-19 cases at any given month, and β_i is the difference in the log of expected count of disease cases between January (as a base) and the respective month.

In view of autocorrelation in this time series data, we applied heteroskedasticity and autocorrelation-consistent (HAC) variance estimator using Newey–West as a kernel to compute the HAC standard errors for point estimates from the models. Finally, model residuals for autocorrelation were validated using Breusch–Godfrey and Cumby–Huizinga tests [18]. Model fit was assessed by visualization of residuals and approximation of fitted values. Finally, we exponentiated the model coefficients to report incidence rate ratios (IRRs) along with 95% confidence intervals (CIs) to reflect associations of COVID-19 cases with reporting of various infectious diseases.

Due to logistical challenges in consistent data collection throughout the study period, there were intervals for specific diseases in certain provinces with missing data, including data for all diseases in KPK province for

October 2019 and for malaria in Sindh province for August 2018 to March 2019. All missing points pertained to the pre-pandemic period and were not imputed.

Results

Reported monthly incidence of each of the eight infectious disease pathogens/syndromes decreased in all provinces of Pakistan with the advent of COVID-19 in late February 2020 (Fig. 1). Moreover, the incidence of all these diseases remained lower in 2020–2021 than in the pre-pandemic period, especially when COVID-19 cases peaked (Figs. 1, 2, 3, 4, 5, 6, 7, 8, 9). Reported incidence of COVID-19 and incidence of pneumonia, hepatitis and diarrhea were negatively correlated in all five provinces ($p < 0.05$), whereas a negative correlation was observed for measles and typhoid in all provinces except Balochistan (Table 1). Notably, a negative correlation between AURI and COVID-19 occurred in all provinces except for Sindh and KPK. Similarly, the reported incidence of malaria was negatively correlated in all provinces except KPK and Balochistan. However, the incidence of NNT was only negatively correlated in Sindh province. Median monthly reported cases of all diseases along with inter-quartile range (Q1–Q3) in the pre-pandemic and pandemic periods in various provinces/regions are presented in Table 2 and visualized in Additional file 1: Fig. S1.

Respiratory infections

Median number of reported AURI cases decreased in the pandemic period compared to the pre-pandemic period in all provinces/regions (Table 2). As the cases of COVID-19 began in late February 2020 and peaked




Fig. 1 Monthly number of reported cases of various infectious diseases or syndromes in Pakistan from March 2018 to February 2021 and cases of COVID-19 from March 2020 to February 2021



Fig. 2 Monthly number of reported cases of acute upper respiratory infection (AURI) and COVID-19 in five provinces/regions of Pakistan from March 2018 to February 2021








Fig. 3 Monthly reported cases of pneumonia and COVID-19 in five provinces/regions of Pakistan from March 2018 to February 2021








Fig. 8 Monthly reported cases of NNT and COVID-19 in five provinces/regions of Pakistan from March 2018 to February 2021




Fig. 9 Monthly reported cases of malaria and COVID-19 in five provinces/regions of Pakistan from March 2018 to February 2021

Table 1 Spearman correlation of COVID-19 cases with various infectious diseases/syndromes in various provinces of Pakistan

	Punjab		Sindh		Khyber Pakhtunkhwa		Balochistan		AJK & GB ^a	
	r_s	p value	r_s	p value	r_s	p value	r_s	p value	r_s	p value
Acute upper respiratory infection	-0.752	<0.001	-0.266	0.118	-0.227	0.189	-0.510	0.002	-0.799	<0.001
Pneumonia	-0.721	<0.001	-0.529	0.001	-0.339	0.046	-0.445	0.006	-0.772	<0.001
Hepatitis	-0.819	<0.001	-0.592	<0.001	-0.562	<0.001	-0.735	<0.001	-0.742	<0.001
Diarrhea	-0.701	<0.001	-0.514	0.001	-0.375	0.026	-0.680	<0.001	-0.759	<0.001
Typhoid	-0.679	<0.001	-0.363	0.029	-0.562	<0.001	-0.138	0.426	-0.761	<0.001
Malaria	-0.820	<0.001	-0.554	0.002	0.029	0.867	-0.045	0.794	-0.758	<0.001
Measles	-0.666	<0.001	-0.742	<0.001	-0.558	<0.001	-0.118	0.494	-0.728	<0.001
Neonatal tetanus	-0.012	0.943	-0.394	0.017	-0.085	0.626	-0.077	0.654	-0.036	0.834

^a Azad Jammu & Kashmir & Gilgit Baltistan

in June 2020 in Punjab and in AJK & GB, the reported number of AURI cases decreased compared to the preceding years' cases (Fig. 2). The lowest incidence of AURI occurred between May and July 2020 and approached the preceding years' reported incidence from August onwards until November 2020 when another dip concurrent with the second wave of COVID-19 in these provinces. Similar trends were observed for Sindh, KPK and Balochistan in which a reduced incidence of AURI occurred during the first wave of COVID-19, though the second wave did not seem to be associated with as much a decline in incidence of AURI. The regression model revealed that when seasonal trends are controlled for, an increase in COVID-19 cases was associated with a reduction in the IRR of AURI in all provinces/regions, particularly in AJK & GB (Table 3).

A similar pattern was observed with pneumonia (Table 2). Reported pneumonia cases decreased with the advent of increasing reported COVID-19 cases in 2020–2021 compared to the reported pneumonia cases in the same months in the preceding years in Punjab and Sindh (Fig. 3). A similar pattern was observed in KPK and Balochistan, but only for the first wave of the pandemic. In AJK & GB, there was a sharp decrease in reported pneumonia cases with the onset of the pandemic and the lowest numbers of reported cases were observed in November 2020 when COVID-19 cases peaked in these regions (Fig. 3). The regression model demonstrated that the reported pneumonia cases decreased in all provinces/regions, with an increase of reported COVID-19 cases, when seasonality was included in the model (Table 3).

Food and waterborne diseases

Reported incidence of viral hepatitis in all provinces/regions decreased at the start of the pandemic (Table 2, Fig. 4) and was consistently lower in the pandemic period compared to the preceding years in Punjab, especially

during both waves of COVID-19. Viral hepatitis incidence decreased compared to preceding years in Sindh, but only during the first wave of COVID-19. Similar trends were observed in other provinces. The regression model adjusted for seasonality revealed that an increase of reported COVID-19 cases was associated with reduced IRRs of viral hepatitis in all provinces/regions (Table 3).

A similar pattern was observed with diarrhea in all five provinces/regions (Table 2). Our regression model revealed that the increase of COVID-19 cases was associated with lowering the incidence of reported cases of diarrhea by 1.6% in Punjab, 0.8% in Sindh, 2.5% in KPK, 9.7% in Balochistan, and 47.9% in AJK & GB (Table 3). In addition, median reported cases of diarrhea during the pandemic period (2020–2021) were lower compared to the corresponding pre-pandemic period (2018–2020) in all provinces/regions (Table 2). The incidence of diarrhea cases did not increase from March to July in 2020 as it did for the 2 preceding years in Punjab (Fig. 5). As the incidence of COVID-19 declined after July 2020, the incidence of diarrhea approached the preceding years' levels during the fall of 2020, and, with the second wave of COVID-19, another decline from the previous years' levels was observed in Punjab. However, in Sindh, KPK, Balochistan and AJK & GB, the reported number of diarrhea cases was lower than the previous years' only during the first wave of COVID-19 but not during the second wave (Fig. 5).

Reported incidence of typhoid was also negatively associated with increasing cases of COVID-19 in our regression model that depicted a significant association between typhoid and COVID-19 in all provinces/regions (Table 3). Median reported cases of typhoid fever were higher during the corresponding pre-pandemic compared to the pandemic period in Punjab, Sindh, KPK, and AJK & GB, but not Balochistan (Table 2). There was a sharp reduction in the number of cases of typhoid in

Table 2 Monthly number of cases of various infectious diseases/syndromes in various provinces of Pakistan before and after the onset of COVID-19 pandemic expressed as median (Q1–Q3)

	Monthly cases Pre-COVID period (March 2018–February 2020)	Monthly cases COVID period (March 2020–February 2021)	p value
Acute upper respiratory infection			
Punjab	3,069,346 (2,735,657–3,218,780)	2,006,049 (1,474,394–2,385,241)	< 0.001
Sindh	1,391,410 (1,169,758–1,620,949)	1,287,438 (826,934–1,451,322)	0.003
Khyber Pakhtunkhwa	549,015 (478,170–665,718)	465,123 (339,966–671,673)	0.005
Balochistan	187,799 (161,662–215,482)	163,602 (109,895–177,966)	< 0.001
AJK & GB ^a	116,112 (102,663–127,583)	42,116 (30,764–58,881)	< 0.001
Pneumonia			
Punjab	135,383 (104,878–158,489)	85,240 (63,440–98,109)	< 0.001
Sindh	121,051 (105,231–132,694)	90,485 (53,528–101,750)	< 0.001
Khyber Pakhtunkhwa	51,385 (47,393–59,902)	42,099 (32,135–54,206)	0.002
Balochistan	29,717 (25,860–32,946)	24,649 (15,922–27,566)	0.002
AJK & GB	29,795 (21,942–36,362)	9871 (7582–12,321)	0.001
Hepatitis			
Punjab	121,129 (105,419–134,753)	67,946 (50,033–75,602)	< 0.001
Sindh	53,351 (47,236–58,453)	40,325 (29,190–44,709)	0.001
Khyber Pakhtunkhwa	11,772 (11,061–13,648)	8943 (8060–11,423)	0.005
Balochistan	1951 (1699–2538)	720 (512–1140)	0.001
AJK & GB	2431 (2212–3144)	210 (140–537)	< 0.001
Diarrhea			
Punjab	980,016 (743,933–1,108,243)	608,436 (596,977–679,985)	< 0.001
Sindh	538,899 (417,356–613,838)	396,529 (357,571–448,058)	0.009
Khyber Pakhtunkhwa	347,301 (244,320–397,148)	251,935 (179,117–330,954)	0.024
Balochistan	137,060 (118,872–142,008)	99,075 (85,077–105,934)	< 0.001
AJK & GB	57,419 (45,728–69,461)	22,335 (17,320–34,625)	0.001
Typhoid			
Punjab	114,277 (107,429–121,366)	77,258 (67,090–94,348)	< 0.001
Sindh	40,952 (37,368–45,210)	32,629 (29,515–40,423)	0.016
Khyber Pakhtunkhwa	41,403 (34,867–45,410)	30,171 (20,901–36,690)	0.003
Balochistan	12,797 (12,461–13,982)	12,732 (11,021–13,422)	0.266
AJK & GB	11,171 (8862–13,178)	2710 (2097–3436)	0.001
Malaria			
Punjab	124,713 (114,852–130,712)	68,394 (59,933–85,812)	< 0.001
Sindh	247,570 (232,257–262,272)	197,501 (169,485–235,045)	0.125
Khyber Pakhtunkhwa	92,535 (78,840–112,447)	97,759 (71,764–108,433)	0.700
Balochistan	49,232 (47,222–57,941)	45,421 (42,173–59,299)	0.970
AJK & GB	11,525 (9145–13,089)	1681 (1069–3175)	0.001
Measles			
Punjab	2223 (1396–3072)	570 (424–746)	0.002
Sindh	796 (617–1718)	227 (64–309)	< 0.001
Khyber Pakhtunkhwa	3900 (2080–4378)	1814 (1525–2085)	0.007
Balochistan	951 (860–1129)	918 (639–1279)	0.970
AJK & GB	917 (679–1110)	86 (51–211)	0.001
Neonatal tetanus			
Punjab	100 (47–226)	113 (9–143)	0.733
Sindh	15 (12–49)	6 (2–17)	0.258
Khyber Pakhtunkhwa	274 (160–520)	176 (140–423)	0.413
Balochistan	17 (11–48)	8 (5–68)	0.910
AJK & GB	0 (0–2)	0 (0–0)	0.750

^a Azad Jammu & Kashmir & Gilgit Baltistan

Punjab, compared to previous years, as the incidence of COVID-19 increased between March and July 2020, as well as in November 2020. A similar reduction in cases of typhoid occurred in this province as did in KPK—with a bimodal reduction of cases during the two waves of the COVID-19 pandemic in 2020. Of note, Sindh province had a reduction in typhoid incidence during the first wave of the pandemic only, whereas aside from a small reduction of typhoid cases in April and May 2020 during the first wave of the pandemic in Balochistan, there was no notable change in reported incidence in that province. In contrast, in AJK & GB, typhoid incidence dropped precipitously after the onset of the pandemic and remained significantly lower than that seen in previous years throughout the study period (Fig. 6).

Vaccine preventable diseases

Median recorded cases of measles differed in the pre-pandemic vs. the pandemic period all provinces/regions but Balochistan (Table 2, Fig. 7). However, when seasonality was incorporated, reported cases of measles were negatively associated with pandemic cases in all provinces/regions (Table 3). Reductions in measles incidence during the first wave of the pandemic occurred in all five provinces/regions, whereas reductions in measles incidence during the second wave primarily occurred in Punjab, Sindh, and AJK & GB and not in KPK or Balochistan (Fig. 7).

Incidence of NNT did not differ in the pre-pandemic vs. the pandemic period in any province: (Table 2). However, the regression model incorporating seasonality revealed that reported cases of NNT were negatively associated with pandemic cases in Punjab and Sindh, but not in KPK, Balochistan and AJK & GB (Table 3). The incidence of NNT was lower in Punjab in the first wave of the pandemic but not during the second wave, whereas the reverse was true for Sindh (Fig. 8). Furthermore, there was no difference in the reported incidence of NNT in KPK, Balochistan and AJK & GB between the pre-pandemic and the pandemic periods (Fig. 8).

Vector-borne diseases

Reported incidence of malaria differed in the pre-pandemic vs the pandemic period only in Punjab and AJK & GB. Interestingly, our regression model demonstrated that reported cases of malaria were negatively associated with pandemic cases in Punjab, Sindh, and AJK & GB, in contrast with KPK and Balochistan, where no significant difference was detected (Table 3). The incidence of malaria was lower during both the first and second waves of the pandemic in Punjab, Sindh and AJK & GB. Conversely, there was no difference in malaria case numbers between the pre-pandemic and pandemic

periods in Balochistan, although a possible trend towards a decreased number of cases occurred during the first wave of the COVID-19 pandemic in KPK (Fig. 9).

Discussion

The reported incidence of several infectious disease pathogens/syndromes decreased after the first case of COVID-19 was reported in Pakistan in late February 2020. The decrease occurred for all eight infectious disease pathogens/syndromes in most of the provinces/regions included in this study: AURI, pneumonia, viral hepatitis, diarrhea, typhoid fever, measles, malaria and NNT. The only exceptions were malaria in KPK and Balochistan, as well as NNT in KPK, Balochistan and AJK & GB, which demonstrated no difference in IRRs between the pre-pandemic and pandemic periods—after seasonality was accounted for. A potential reason that these two particular infectious diseases did not demonstrate a significant difference in all five provinces/regions is that the risk factors, clinical manifestations, and diagnosis of these two diseases are, generally, different enough from that of COVID-19 so as not to cause confusion and lead to frequent misclassification or misdiagnosis [19, 20]. Presumably, there was sufficient priority and interest in the control and prevention of malaria and NNT in Pakistan, to ensure that resources were not diverted from them to combat the COVID-19 pandemic [21, 22] in every province. Moreover, in respect to NNT specifically, case numbers in every province were relatively limited at the outset (i.e., in the tens to hundreds), likely as a result of vaccination efforts over the years. As an illustration, there were no cases of NNT in AJK & GB during the study period.

The incidence of several of the diseases such as malaria are very affected by seasonal fluctuations, likely related to the life cycle of its mosquito vectors [23] and other weather-related factors. It is therefore important to include seasonality in the model when analyzing the association between reported COVID-19 cases and incidence of these diseases. This is illustrated by the fact that when using the Wilcoxon signed-rank test (which does not take the varying trend of COVID-19 cases into account), the reported incidence of the 8 diseases was associated with the reported incidence of COVID-19 in less provinces/regions than in the regression models that included seasonality.

Monthly incidence of most of the infectious disease pathogens/syndromes (AURI, pneumonia, viral hepatitis, diarrhea, typhoid fever, and measles) decreased with the advent of COVID-19 in late February 2020 overall in the entire country (Fig. 1) as well as in all five provinces/regions separately (Figs. 2, 3, 4, 5, 6, 7, 8, 9). A potential explanation is that after the first COVID-19 cases were

Table 3 Overall interrupted time series analysis using generalized linear negative binomial regression models adjusted for seasonality for various infectious diseases/syndromes as dependent variables and COVID-19 cases in thousands as an independent variable in various provinces/regions of Pakistan

	Punjab		Sindh		Khyber Pakhtunkhwa		Balochistan		AJK & GB ^b	
	IRR ^a (95% CI)	p value	IRR (95% CI)	p value	IRR (95% CI)	p value	IRR (95% CI)	p value	IRR (95% CI)	p value
Acute upper respiratory infection	0.981 (0.979, 0.983)	< 0.001	0.993 (0.992, 0.994)	< 0.001	0.976 (0.970, 0.980)	< 0.001	0.906 (0.878, 0.935)	< 0.001	0.527 (0.509, 0.546)	< 0.001
Pneumonia	0.979 (0.975, 0.983)	< 0.001	0.987 (0.985, 0.988)	< 0.001	0.962 (0.955, 0.968)	< 0.001	0.895 (0.864, 0.928)	< 0.001	0.474 (0.462, 0.487)	< 0.001
Hepatitis	0.976 (0.972, 0.980)	< 0.001	0.989 (0.987, 0.991)	< 0.001	0.959 (0.952, 0.967)	< 0.001	0.695 (0.636, 0.760)	< 0.001	0.250 (0.217, 0.289)	< 0.001
Diarrhea	0.984 (0.983, 0.986)	< 0.001	0.992 (0.990, 0.993)	< 0.001	0.975 (0.968, 0.982)	< 0.001	0.903 (0.874, 0.934)	< 0.001	0.521 (0.507, 0.536)	< 0.001
Typhoid	0.985 (0.981, 0.988)	< 0.001	0.994 (0.991, 0.996)	< 0.001	0.959 (0.952, 0.965)	< 0.001	0.956 (0.922, 0.990)	0.012	0.386 (0.365, 0.407)	< 0.001
Malaria	0.979 (0.975, 0.983)	< 0.001	0.991 (0.990, 0.992)	< 0.001	1.000 (0.991, 1.010)	0.946	0.981 (0.940, 1.023)	0.369	0.268 (0.242, 0.298)	< 0.001
Measles	0.955 (0.947, 0.963)	< 0.001	0.951 (0.945, 0.958)	< 0.001	0.941 (0.923, 0.959)	< 0.001	0.875 (0.854, 0.896)	< 0.001	0.278 (0.248, 0.311)	< 0.001
Neonatal tetanus	0.943 (0.903, 0.985)	0.008	0.953 (0.917, 0.990)	0.014	0.980 (0.933, 1.030)	0.427	0.790 (0.597, 1.046)	0.099	0.851 (0.555, 1.304)	0.457

^a Incidence rate ratio

^b Azad Jammu & Kashmir & Gilgit Baltistan

reported in late February 2020, precautionary measures such as the closure of schools, work from home policies, public health restrictions and lockdowns were implemented [24], which adversely affected surveillance and data reporting of these diseases. In addition, the outpatient treatment departments (OPDs) of all major hospitals across the country were closed for patients during the initial few weeks of the pandemic [24, 25] leading to a substantial effect on the inflow of patients to the hospitals (potential data points for surveillance) for reasons other than COVID-19. This explains a sudden drop in reported cases of almost all major ongoing infectious diseases in all provinces. Subsequently, different provinces adopted different strategies to re-open under their respective administrative controls. With the onset of pandemic, a sudden increase in the “fear factor” associated with forced isolation and perceived stigma around a novel disease, largely spread by hyper-reactive and sensational media reports [26], also negatively affected the public’s willingness to seek medical attention through qualified doctors and hospitals, replacing this instead with a surge in self-prescription and home remedies for various health problems [27, 28]. This trend is likely one of the main reasons for a sudden decline of reported incidence of various diseases with the escalating pandemic in many developing countries such as Pakistan. Alternatively, or perhaps additionally, several of these infectious disease pathogens/syndromes have presentations which overlap with those of COVID-19, particularly considering the purely clinically based case definitions used by the Pakistan IDSR, and may have been mistakenly diagnosed as such [16].

Some interesting findings became clear when modeling these infectious disease pathogens/syndromes. It is not surprising to see an inverse relationship between the number of COVID-19 cases and the number of AURI cases (Fig. 2), given the overlap in symptoms between these 2 conditions. The diagnosis of AURI is made clinically and is essentially defined as an upper respiratory tract infection (URTI), whereas upper respiratory symptoms, including sore throat, cough and rhinorrhea, are well documented symptoms of COVID-19 [29]. As such, given the overlap in symptoms and lack of clear diagnostic testing for AURI, it is highly likely that COVID-19 could have been mistaken for AURI and vice versa, especially in a developing setting with less access to diagnostic testing. A similar argument could be made for pneumonia (Fig. 3), given the overlap in symptoms between these two conditions. Pneumonia due to COVID-19 is well described [30] and, thus, there is ample opportunity for misclassification between COVID-19 pneumonia and pneumonia due to other pathogens, particularly in a

developing setting with less access to distinct and differential diagnostic testing.

There was a clear inverse relationship between COVID-19 cases and the number of reported cases of viral hepatitis (Fig. 4), believed to be mainly food and waterborne hepatitis due to hepatitis A and E. As the number of cases of COVID-19 in Pakistan increased, peaked in June 2020 and then declined, the number of reported cases of viral hepatitis steadily decreased, plateaued and then began to increase and approach the expected incidence again by September 2020 in most province/regions. We believe that the inverse relationship in this case may be primarily due to a diversion of resources, including diagnostic testing capacity, from viral hepatitis towards COVID-19 testing and management, although it may also be due to an overlap in symptoms during the pre-icteric phase of viral hepatitis, resulting in misdiagnosis and misclassification.

Similarly, the reported incidence of diarrhea in Pakistan declined with the onset of COVID-19 cases in late February 2020 (Fig. 5), perhaps because diarrhea is a common manifestation of COVID-19 [16, 29]. As expected, as COVID-19 case numbers decreased in July and August 2020, reported cases of diarrheal illness continued to increase and peaked in September 2020 (Fig. 6). Notably, typhoid fever is mainly transmitted through exposure to contaminated food and water, the latter of which is more likely to be consumed in larger quantities during the hot summer months. Regardless, there was a clear negative impact of the onset of COVID-19 cases in late February 2020 on the reported incidence of typhoid fever, and a similar phenomenon occurred in October through December 2020. The general inverse correlation between typhoid fever and COVID-19 was likely due to both a diversion of resources from the former to the latter as well as an overlap in symptoms such as fever, weakness, diarrhea and cough—resulting in misdiagnosis and misclassification [31].

The incidence of reported measles cases also decreased after the advent of COVID-19 cases in late February 2020, whereas reductions in COVID-19 cases since that time were associated with increased reporting of measles cases (Fig. 7). This was attributed to a diversion of resources during the COVID-19 pandemic previously put towards measles diagnostic testing, tracing and management. Of interest, the impact of the pandemic on the reporting of measles cases was significant during the first wave of the pandemic, where all provinces had a notable decline in case numbers, whereas in Balochistan and AJK & GB the same reduction in incidence did not occur during the second wave. This may have been due to some level of adjustment and improvement in measles diagnostic testing and tracing that were in place by this point in time in those provinces/regions.

The NNT case numbers were low at the outset (Fig. 8), and apparently least affected by the pandemic of all diseases/syndromes that we explored, demonstrating a small reduction during the first wave of the pandemic in Punjab and during the second wave of the pandemic in Sindh, but not otherwise. Malaria case numbers were impacted by the onset of the pandemic in some provinces/regions more than others, with a notable decrease in case numbers in Punjab, Sindh and AJK & GB (Fig. 9). As alluded to, this infectious disease was the most impacted by seasonal variations, as demonstrated by the difference between our Wilcoxon signed-rank test and our negative binomial regression model.

Despite a growing body of literature attempting to speculate, define and describe the impact of the COVID-19 pandemic on other ongoing infectious diseases outbreaks in many developing settings, to our knowledge our study is the first original research paper to do this using infectious diseases surveillance data from the MoNHSR&C. This had several advantages: we were able to utilize a set of national level data categorized by province/region and our interrupted time series study design allowed us to implement data that were objectively defined, easy to collect, simple to present graphically and straightforward to interpret.

Our study also had several weaknesses, the most important of which is that we were only able to determine correlations and associations between the onset of the COVID-19 pandemic and each of these other infectious disease pathogens/syndromes. Based on our study design, we were unable to ascertain causality. Moreover, several of these infectious disease pathogens/syndromes overlap to varying degrees with symptoms and manifestations of COVID-19 and could easily have led to misclassification bias during data collection. Due to efforts made early on by the Pakistani Government, especially under NCOC, to control COVID transmission using various non-pharmaceutical containment measures inclusive of lockdowns, implementation of masking, sanitization and social distancing policies and educational campaigns [17], there was likely an early reduction in the incidence of influenza and other respiratory infections [32] which could certainly have impacted our findings. Notably, however, many regions of the country have had poor access to masks and hand sanitizers, particularly the rural parts of the country [33], and the countrywide lockdowns were short-lived for financial reasons, ending in May 2020.

There also may have been differences in data collection resources in various parts of the country at some time points, as well as national changes in data collection methodologies over time, as the Pakistani infectious disease surveillance system has evolved and somewhat improved over the past few years from completely paper-based to

primarily electronic [34–36]. Nevertheless, despite some improvements in technology, it is notable that several infections that are preventable through appropriate surveillance and coupled public health efforts, including tuberculosis, meningitis, malaria, typhoid and hepatitis B and C, are still responsible for nearly half of the deaths in the country [37]. Overcoming major financial and logistic barriers, through collaboration with international partners, will be critical in developing a more vigorous and reliable national surveillance system—with laboratory integration—to properly serve the needs of the country in a sustainable fashion [38, 39]. As a result of the existing system, our surveillance data was, therefore, not consistently robust and comprehensive throughout the entire reporting interval, potentially resulting in imperfect capture of monthly incidence for some pathogens/syndromes. Fortunately, there were no missing data during the pandemic period and, thus, no need was ascertained to impute data. Importantly, however, the disease definitions we used, though based on Pakistani national surveillance protocols, are by their nature non-specific and somewhat difficult to compare with one another—some representing specific pathogens and others representing syndromes. Moreover, we worked on aggregated data and did not have access to case-based data; therefore, were unable to cater for confounder and effect modifiers at study design or analysis levels.

It is evident that the incidence, surveillance, and reporting of the studied diseases were affected considerably with progression of the pandemic. Whatever the situation was, it can be assumed that it resulted in adverse consequences with respect to assessing true infectious disease burden and capturing all outbreaks. As a result, many cases of these endemic infectious disease pathogens/syndromes have likely been present in Pakistan over the past 24 months, but not captured and mitigated by the health system, resulting in uncontained transmission to susceptible populations. Notably, there is a growing body of literature that corroborates the fact that several endemic infectious diseases in Pakistan have actually become more prevalent as a result of a diversion of scarce resources towards combatting the COVID-19 pandemic. Weakened and ineffective approaches towards the prevention and control of HIV disease, poliomyelitis, Dengue fever, tuberculosis and mucormycosis infections with a resulting increasing burden of disease have recently been well-documented and highlighted in Pakistan, particularly over the past couple years [40–44]. We believe that only time will demonstrate the full impact of COVID-19 on these and other endemic diseases, and the challenges that will need to be overcome just to achieve the same level of control as had been acquired prior to the pandemic. Ultimately, such a situation is likely to

result in further outbreaks, hyper-endemic pathogens, or even other epidemics.

Conclusions

Despite the presence of surveillance systems for several infectious disease pathogens/syndromes in Pakistan, eradication has been difficult, given the number of concurrent endemic pathogens, resource limitations, and the vastness of the population. The COVID-19 pandemic has affected surveillance of other important ongoing endemic infectious diseases, including AURI, viral hepatitis, pneumonia, diarrhea, typhoid fever, measles, NNT, and malaria. Though the exact reasons for this cannot be determined by our study, we confidently postulate that this was in some degree due to a lack of sufficient resources, and diversion of existing resources towards addressing the COVID-19 pandemic. In particular, there is a lack of laboratory diagnostic capability to differentiate between COVID-19 and several other infectious disease pathogens endemic to Pakistan [16].

It is recommended that surveillance and reporting systems of diseases should be strengthened and coordinated so that any change in disease trends may be detected and responded to at the earliest opportunity to contain disease spread locally. There is also a need to develop public health emergency preparedness plans to cater to outbreaks and health emergencies more efficiently and effectively. Countrywide awareness and risk communication regarding personal protective measures, access to safe food and water supply, and education and access to vaccination are among interventions that are desperately needed in Pakistan's battle against these various infectious diseases. It was recognized early on in the COVID-19 pandemic that vaccination would offer the global community the best chance at controlling it. Though many countries were early adopters [45], Pakistan has suffered from the twin perilous ideologies of vaccine hesitancy [46] and vaccine nationalism [47, 48]. The former has been a longstanding problem in the country and is the major reason for which it has been one of the last bastions of poliomyelitis in the world in the years leading up to the pandemic, whereas the latter has been deleterious not just for low and middle income countries such as Pakistan but for the developed world as well (despite each nation believing that it is acting in its best self-interest) [47, 48], resulting in the development of new variant strains and lengthening the duration of the pandemic. Regrettably, if these factors persist, the consequences will be much worse for Pakistan's dilapidated health-care sector, which is in dire need of development and implementation of concrete and achievable policies for the country's largest-ever mass vaccination campaign to achieve set targets within planned timelines. In addition,

educational campaigns must be carried out to address vaccine misconception and hesitancy issues through various local and mass-media resources such as television, radio, newspapers, websites and social media [47]. If we delay or fail to do this appropriately, then control of the pandemic for Pakistan could become an unfulfilled dream, and more precious lives will be lost.

Despite some limitations, our study was a starting point for characterizing impacts of the COVID-19 pandemic on other endemic infectious diseases in Pakistan as well as in other developing countries. We are convinced that such work will be pivotal in making the case that more resources are required to ensure that we do not lose sight of other control and eradication efforts, while our attention is captured by this all-consuming pandemic. If we are to act otherwise, many of the great efforts that have been made over the previous decades may be consigned to oblivion.

Abbreviations

CCHF: Crimean-Congo Hemorrhagic fever; COVID-19: Novel coronavirus 2019; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2; NCOC: National Command and Operations Center; MoNHSR&C: Ministry of National Health Services Regulations and Coordination; KPK: Khyber Pakhtunkhwa; AJK: Azad Jammu & Kashmir; GB: Gilgit Baltistan; IDSR: Integrated Disease Surveillance Response System; AURI: Acute upper respiratory infection; NNT: Neonatal tetanus; DHIS: Demographic Health Information System; IQR: Interquartile range; HAC: Heteroskedasticity and autocorrelation-consistent; IRRs: Incidence rate ratios; CIs: Confidence intervals; OPDs: Outpatient treatment departments; URTI: Upper respiratory tract infection.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12879-022-07869-3>.

Additional file 1: Figure S1. Boxplots of distributions of monthly reported cases of various diseases pre-COVID and after the onset of the COVID-19 pandemic in five provinces/regions of Pakistan.

Acknowledgements

We acknowledge the Pakistan National Institute of Health and the MoNHSR&C of the Government of Pakistan for coordinating closely to provide us with access to the data used to complete this project. In particular, we offer our sincere thanks to Drs. Rana Safdar and Jamil Ansari, as well as their staff, for their help. Finally, the authors thank Dr. John Kastelic for his assistance in editing the manuscript.

Author contributions

BM contributed substantially to the conception and design of this project and was deeply involved in the interpretation of its results and in drafting and editing the manuscript. MWM contributed to the design of the project, assisted in acquiring the necessary data to complete it, and was involved in drafting the manuscript. WS carried out the statistical analyses, assisted in their interpretation, and helped revise the manuscript. MAR assisted in the original design of the project, as well as in data acquisition and statistical analysis. AI approved the original study design, and substantively revised the manuscript. HB supervised the project, guided its completion, assisted in determining the approach to the analysis of the data as well as its presentation, and was centrally involved in manuscript revision. All authors read and approved the final manuscript.

Authors' information

Bayan Missaghi is a Clinical Associate Professor of Medicine at the University of Calgary and an infectious disease physician consultant with Alberta Health Services. Muhammad Wasif Malik and Muazam Abbas Ranjha are Senior Scientific Officers working in the Division of Field Epidemiology & Disease Surveillance at the Pakistan National Institute of Health. Waseem Shaukat is a Doctor of Veterinary Medicine and PhD scholar at the University of Calgary, with a special interest in biostatistics and epidemiology of infectious diseases. Aamer Ikram is the Executive Director of the Pakistan National Institute of Health and Director of the Pakistan Field Epidemiology and Laboratory Training Program; he has been awarded FRCP by the Royal College of Edinburgh, FRCPath by the Royal College of Pathologists in London, and a Fellowship in Public Health from Royal Colleges UK. Herman W. Barkema is a Professor in Epidemiology of Infectious Diseases and Director of One Health at UCalgary at the University of Calgary, Scientific Director of Antimicrobial Resistance—One Health Consortium, and an NSERC Industrial Research Chair in Infectious Diseases of Dairy Cattle.

Funding

No funding was secured or used for this study.

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available and were acquired with permission from the Government of Pakistan MoNHSR&C DHIS database. These data will not be provided as a supplement to this manuscript but may be made available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

As all data were completely de-identified and de-coupled from individual patient-level data, neither ethical approval nor patient consent was required. Permission was obtained from the Government of Pakistan MoNHSR&C prior to conducting the study. As per the Pakistan National Bioethics Committee—Research Ethics Committee policy, 'studies that do not require direct human subject participants (e.g. chart review, already collected data for routine public health reasons, etc.) may be exempted from review' as was the case for our study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests with respect to this study.

Author details

¹Department of Medicine, Cumming School of Medicine, University of Calgary and Alberta Health Services, Office 3683, 3500 – 26 Ave NE, Calgary, AB T1Y 6J4, Canada. ²Infection Prevention and Control, Calgary Zone, Alberta Health Services, Calgary, AB, Canada. ³Field Epidemiology & Disease Surveillance Division, National Institute of Health (NIH), Islamabad, Pakistan. ⁴Department of Production Animal Health, Faculty of Veterinary Medicine, University of Calgary, 3330 Hospital Drive NW, HSC 2561, Calgary, AB, Canada. ⁵Executive Office, National Institute of Health (NIH), Islamabad, Pakistan. ⁶Department of Community Health Sciences, Cumming School of Medicine, University of Calgary, Office HS2521, 3330 Hospital Dr. NW, Calgary, AB T2N 4N1, Canada. ⁷One Health at UCalgary, University of Calgary, Calgary, AB, Canada.

Received: 31 March 2022 Accepted: 10 November 2022

Published online: 26 November 2022

References

- Khalil AT, Ali M, Tanveer F, Ovais M, Idrees M, Shinwari ZK, et al. Emerging viral infections in Pakistan: issues, concerns, and future prospects. *Health Secur.* 2017;15(3):268–81.

- Sultan F, Khan A. Infectious diseases in Pakistan: a clear and present danger. *Lancet.* 2013;381(9884):2138–40.
- Shah S, Abbas G, Riaz N, Anees Ur R, Hanif M, Rasool MF. Burden of communicable diseases and cost of illness: Asia pacific region. *Expert Rev Pharmacoecon Outcomes Res.* 2020;20(4):343–54.
- Salahuddin N, Khalid M, Baig-Ansari N, Iftikhar S. Five-year audit of infectious diseases at a tertiary care hospital in Karachi, Pakistan. *Cureus.* 2018;10(11):e3551.
- Mejia N, Qamar F, Yousafzai MT, Raza J, Garrett DO, Date K, et al. Typhoid and paratyphoid cost of illness in Pakistan: patient and health facility costs from the surveillance for enteric fever in Asia Project II. *Clin Infect Dis.* 2020;71(Supplement_3):S319–35.
- Atif M, Malik I. Why is Pakistan vulnerable to COVID-19 associated morbidity and mortality? A scoping review. *Int J Health Plan Manag.* 2020;35(5):1041–54.
- Government of Pakistan COVID-19 Situation. <https://covid.gov.pk/>. Accessed 1 Dec 2021.
- Worldometer Countries where COVID-19 has spread. <https://www.worldometers.info/coronavirus/countries-wherecoronavirus-has-spread/>. Accessed 1 Dec 2021.
- Ali M, Ahmad N, Khan H, Ali S, Akbar F, Hussain Z. Polio vaccination controversy in Pakistan. *Lancet (London, England).* 2019;394(10202):915–6.
- Khan YH, Mallhi TH, Alotaibi NH, Alzarea AI, Alanazi AS, Tanveer N, et al. Threat of COVID-19 vaccine hesitancy in Pakistan: the need for measures to neutralize misleading narratives. *Am J Trop Med Hyg.* 2020;103(2):603–4.
- Awan UA, Zahoor S, Ayub A, Ahmed H, Aftab N, Afzal MS. COVID-19 and arboviral diseases: another challenge for Pakistan's dilapidated healthcare system. *J Med Virol.* 2021;93(7):4065–7.
- Awan UA, Zahoor S, Rehman K, Khattak AA, Ahmed H, Aftab N, et al. COVID-19 and influenza H1N1: a dangerous combination for Pakistan in the upcoming winter season. *J Med Virol.* 2021;93(4):1875–7.
- Haqqi A, Awan UA, Ali M, Saqib MAN, Ahmed H, Afzal MS. COVID-19 and dengue virus coepidemics in Pakistan: a dangerous combination for an overburdened healthcare system. *J Med Virol.* 2021;93(1):80–2.
- Ahmed W, Malik MW, Wattoo MM, Nadeem F, Ahmed Chaudhry MS, Afzal MS, et al. Impact of COVID-19 pandemic on surveillance of Crimean-Congo haemorrhagic fever (CCHF) in Pakistan. *Travel Med Infect Dis.* 2021;41: 102011.
- Awan UA, Khattak AA, Afzal MS, Khan HI, Saqib MAN, Ahmed H. COVID-19 and Tuberculosis overlapping epidemics: a holistic review from Pakistan. *J Med Virol.* 2021;93(5):2573–5.
- Rana MS, Usman M, Alam MM, Ikram A, Salman M. Overlapping clinical manifestations of COVID-19 with endemic infectious diseases in Pakistan: a looming threat of multiple lethal combinations. *Infect Ecol Epidemiol.* 2021;11(1):1873494.
- Akhtar H, Afridi M, Akhtar S, Ahmad H, Ali S, Khalid S, et al. Pakistan's response to COVID-19: overcoming national and international hypes to fight the pandemic. *JMIR Public Health Surveill.* 2021;7(5): e28517.
- Baum CF, Schaffer ME. ACTEST: Stata module to perform Cumby-Huizinga general test for autocorrelation in time series. *Statistical Software Components, Boston College Department of Economics.* 2013;5457668.
- Bae C, Bourget D. Tetanus [updated 2021 Dec 1]. In: *StatPearls [Internet] Treasure Island: StatPearls Publishing; 2022.*
- Di Gennaro F, Marotta C, Locantore P, Pizzolo D, Putoto G. Malaria and COVID-19: common and different findings. *Trop Med Infect Dis.* 2020;5(3):141.
- Lambo JA, Nagulesapillai T. Neonatal tetanus elimination in Pakistan: progress and challenges. *Int J Infect Dis.* 2012;16(12):e833–42.
- Nasir SMI, Amarasekara S, Wickremasinghe R, Fernando D, Udagama P. Prevention of re-establishment of malaria: historical perspective and future prospects. *Malar J.* 2020;19(1):452.
- Selvaraj P, Wenger EA, Gerardin J. Seasonality and heterogeneity of malaria transmission determine success of interventions in high-endemic settings: a modeling study. *BMC Infect Dis.* 2018;18(1):413.
- Akhtar H, Afridi M, Akhtar S, Ahmad H, Ali S, Khalid S, et al. Pakistan's response to COVID-19: overcoming national and international hypes to fight the pandemic. *JMIR Public Health Surveill.* 2021;7(5):e28517.
- Jabeen A, Ansari JA, Ikram A, Abbasi SH, Khan MA, Rathore TR, et al. Comparison of actions taken by Pakistan, United Arab Emirates and Vietnam

- for COVID-19 prevention and control. *Glob Biosecur.* 2020. <https://doi.org/10.31646/gbio.79>.
26. Su Z, McDonnell D, Wen J, Kozak M, Abbas J, Šegalo S, et al. Mental health consequences of COVID-19 media coverage: the need for effective crisis communication practices. *Glob Health.* 2021;17(1):4.
 27. Yasmin F, Asghar MS, Naeem U, Najeeb H, Nauman H, Ahsan MN, et al. Self-medication practices in medical students during the COVID-19 pandemic: a cross-sectional analysis. *Front Public Health.* 2022. <https://doi.org/10.3389/fpubh.2022.803937>.
 28. Iqbal Arain M, Shahnaz S, Anwar R, Anwar K. Assessment of self-medication practices during COVID-19 pandemic in Hyderabad and Karachi. *Pak Sudan J Med Sci.* 2021;16(3):347–54.
 29. Johnson KD, Harris C, Cain JK, Hummer C, Goyal H, Perisetti A. Pulmonary and extra-pulmonary clinical manifestations of COVID-19. *Front Med (Lausanne).* 2020;7:526.
 30. Ginsburg AS, Klugman KP. COVID-19 pneumonia and the appropriate use of antibiotics. *Lancet Glob Health.* 2020;8(12):e1453–4.
 31. Haqqi A, Khurram M, Din MSU, Aftab MN, Ali M, Ahmed H, et al. COVID-19 and Salmonella Typhi co-epidemics in Pakistan: a real problem. *J Med Virol.* 2021;93(1):184–6.
 32. Dadras O, Alinaghi SAS, Karimi A, MohsseniPour M, Barzegary A, Vahedi F, et al. Effects of COVID-19 prevention procedures on other common infections: a systematic review. *Eur J Med Res.* 2021;26(1):67.
 33. Haq SU, Shahbaz P, Boz I. Knowledge, behavior and precautionary measures related to COVID-19 pandemic among the general public of Punjab province, Pakistan. *J Infect Dev Ctries.* 2020;14(8):823–35.
 34. Pakistan Go. Ministry of National Health Services Regulations and Coordination. <https://nhsrc.gov.pk/>. Accessed 18 Feb 2022.
 35. Agency JIC. Activities in Pakistan: The District Health Information System (DHIS) project for evidence-based decision making and management. 2022.
 36. Comprehensive assessment of Pakistan's health information system 2017. Cairo: WHO Regional Office for the Eastern Mediterranean; 2019. Licence: CC BY-NC-SA 3.0 IGO.
 37. Baig MA, Shaikh BT. Disease surveillance system: a mandatory conduit for effective control of infectious diseases in Pakistan. *Asia Pac J Public Health.* 2012;24(4):586–94.
 38. Shaikh TG, Waseem S, Ahmed SH, Swed S, Hasan MM. Infectious disease surveillance system in Pakistan: challenges and way forward. *Trop Med Health.* 2022;50(1):46.
 39. Aborode AT, Hasan MM, Jain S, Okereke M, Adediji OJ, Karra-Aly A, et al. Impact of poor disease surveillance system on COVID-19 response in africa: time to rethink and rebuilt. *Clin Epidemiol Glob Health.* 2021;12: 100841.
 40. Khatri G, Sahito AM, Siddiqui MS, Farooq M, Fatima R, Mehmood Q, et al. HIV's imminent menace in a COVID-ridden country: the case of Pakistan. *Ann Med Surg.* 2012;2022(78): 103869.
 41. Sahito AM, Saleem A, Javed SO, Farooq M, Ullah I, Hasan MM. Polio amidst COVID-19 in Pakistan: ongoing efforts, challenges, and recommendations. *Int J Health Plan Manag.* 2022;37(4):1907–11.
 42. Khatri G, Hasan MM, Shaikh S, Mir SL, Sahito AM, Priya, et al. The simultaneous crises of dengue and COVID-19 in Pakistan: a double hazard for the country's debilitated healthcare system. *Trop Med Health.* 2022;50(1):18.
 43. Awan HA, Sahito AM, Sukaina M, Khatri G, Waheed S, Sohail F, et al. Tuberculosis amidst COVID-19 in Pakistan: a massive threat of overlapping crises for the fragile healthcare systems. *Epidemiol Infect.* 2022;150: e41.
 44. Ghazi BK, Rackimuthu S, Wara UU, Mohan A, Khawaja UA, Ahmad S, et al. Rampant increase in cases of mucormycosis in India and Pakistan: a serious cause for concern during the ongoing COVID-19 pandemic. *Am J Trop Med Hyg.* 2021;105(5):1144–7.
 45. Sallam M, Al-Sanafi M, Sallam M. A global map of COVID-19 vaccine acceptance rates per country: an updated concise narrative review. *J Multidiscip Healthc.* 2022;15:21–45.
 46. Mehmood Q, Ullah I, Hasan MM, Kazmi SK, Ahmadi A, Lucero-Prisno DE 3rd. COVID-19 vaccine hesitancy: Pakistan struggles to vaccinate its way out of the pandemic. *Ther Adv Vaccines Immunother.* 2022;10:25151355221077656.
 47. Riaz MMA, Ahmad U, Mohan A, Dos Santos Costa AC, Khan H, Babar MS, et al. Global impact of vaccine nationalism during COVID-19 pandemic. *Trop Med Health.* 2021;49(1):101.
 48. Prasad S, Shahid A, Co ELF, Khatri G, Cheema HA, Rocha ICN, et al. Vaccine apartheid: the separation of the world's poorest and most vulnerable and the birth of Omicron. *Ther Adv Vaccines Immunother.* 2022;10:25151355221107976.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

