

RESEARCH ARTICLE

Open Access



# Potential importance of protease activated receptor (PAR)-1 expression in the tumor stroma of non-small-cell lung cancer

Cong Lin<sup>1\*</sup>, Christof J. Majoor<sup>2†</sup>, Joris J. T. H. Roelofs<sup>3†</sup>, Martijn D. de Kruif<sup>2,4</sup>, Hugo M. Horlings<sup>5</sup>, Keren Borensztajn<sup>1,6,7</sup> and C. Arnold Spek<sup>1</sup>

## Abstract

**Background:** Protease activated receptor (PAR)-1 expression is increased in a variety of tumor cells. In preclinical models, tumor cell PAR-1 appeared to be involved in the regulation of lung tumor growth and metastasis; however the role of PAR-1 in the lung tumor microenvironment, which is emerging as a key compartment in driving cancer progression, remained to be explored.

**Methods:** In the present study, PAR-1 gene expression was determined in lung tissue from patients with non-small-cell lung cancer (NSCLC) using a combination of publicly available RNA microarray datasets and in house-made tissue microarrays including tumor biopsies of 94 patients with NSCLC (40 cases of adenocarcinoma, 42 cases of squamous cell carcinoma and 12 cases of other type of NSCLC at different stages).

**Results:** PAR-1 gene expression strongly correlated with tumor stromal markers (i.e. macrophage, endothelial cells and (myo) fibroblast markers) but not with epithelial cell markers. Immunohistochemical analysis confirmed the presence of PAR-1 in the tumor stroma and showed that PAR-1 expression was significantly upregulated in malignant tissue compared with normal lung tissue. The overexpression of PAR-1 in tumor stroma of NSCLC appeared to be independent from tumor type, tumor stage, histopathological differentiation status, disease progression and patient survival.

**Conclusion:** Overall, our data provide evidence that PAR-1 in NSCLC is mainly expressed on cells that constitute the pulmonary tumor microenvironment, including vascular endothelial cells, macrophages and stromal fibroblasts.

**Keywords:** Protease activated receptor, NSCLC and tumor stroma

## Background

Lung cancer is the leading cause of cancer related death, with around 1.6 million deaths worldwide and the mortality rates for lung cancer are still increasing annually [1, 2]. Non-small-cell lung cancer (NSCLC), the most common type of lung cancer, has a devastating survival outcome. Traditional chemotherapy, including predominantly platinum-based regimens, as first-line standard treatment for NSCLC only shows a modest prolongation of median and overall survival. Despite

aggressive multimodality therapy, 5-year survival rate for patients with stage IV NSCLC at diagnosis is only approximately 2% [3]. More recently, targeted therapies showed efficacy in patients with advanced NSCLC who have specific genetic alterations, like mutations of the anaplastic lymphoma kinase gene or of the epidermal growth factor receptor [1]. However, these available molecular therapies can only be applied to selective patients and the observed benefits are small, suggesting that more in-depth studies of molecules that relate to the pathogenesis of NSCLC is required.

Protease-activated receptor (PAR)-1 is a cell surface seven-transmembrane G protein coupled receptor that is activated by proteolytic cleavage. Removal of the N-terminal extracellular domain of PAR-1 reveals a new

\* Correspondence: c.lin@amc.uva.nl

†Equal contributors

<sup>1</sup>Center for Experimental and Molecular Medicine, Academic Medical Center, Amsterdam 1105 AZ, The Netherlands

Full list of author information is available at the end of the article



tethered ligand that binds to the body of PAR-1 and activates transmembrane signaling to intracellular G proteins, thereby leading to multiple pathophysiological responses [4, 5]. Overexpression of PAR-1 has been detected in various types of cancers, including ovarian, breast, lung, prostate cancer and melanoma [6–10]. Importantly, elevated PAR-1 expression is closely associated with diseases progression and overall survival in breast, prostate, gastric cancer and melanoma [6, 8, 9, 11]. Moreover, tumor cell PAR-1 is recently identified as a promising target to decrease lung cancer progression. Indeed, PAR-1 peptidic inhibitors not only block the migration of both primary and established lung cancer cell lines, but also significantly limit lung tumor growth in nude mice [10]. Moreover, melanoma growth and metastasis were significantly decreased in mice treated with PAR-1 small interfering RNA (siRNA) [12].

During the last decade, the paradigm that tumor growth solely relies on the malignant cells has shifted to a more comprehensive view that tumor growth is dependent on interactions between cancer cells and their adjacent microenvironment, also known as the stroma [13]. The tumor stroma, predominately composed of basement membrane, fibroblasts, vasculature with endothelial cells, inflammatory cells and extra cellular matrix proteins such as collagen and fibronectin [14], is indeed emerging as a key player in promoting carcinogenesis by modulating tumor growth, angiogenesis, invasion and metastasis [15, 16]. Targeting the tumor stroma is consequently under intense investigation as novel treatment strategy in cancer.

Interestingly, PAR-1 expression is not tumor cell specific and PAR-1 is also expressed on key cell types that constitute the tumor stroma such as endothelial cells, fibroblasts and macrophages. Activation of PAR-1 on these stromal cells leads to increased vascular permeability, fibroblast activation, extracellular matrix production and cytokine secretion, thereby potentially driving tumor growth and metastasis [13]. In line with these observations, colonic adenocarcinoma growth was limited in PAR-1-deficient mice, suggesting the importance of PAR-1 in the tumor microenvironment [17]. In addition, pancreatic tumors in PAR-1 deficient animals were significantly smaller compared with tumors in wild type mice. Moreover, the same study also showed that stromal cells drive tumor growth and induce chemoresistance of pancreatic cancer in a PAR-1 dependent manner [18]. Overall these data point to an important role of stromal cell-associated PAR-1 in tumor progression. However, the role of stromal PAR-1 in lung cancer has not been explored yet. In the present study, we examined PAR-1 expression in NSCLC stroma and assessed its correlation with disease progression.

## Methods

### Patients

Tissue microarrays (TMAs, triplicate cores per case) were prepared with tumor sections obtained from NSCLC patients during surgery according to the guidelines of the Medical Ethical Committee of the Academic Medical Center of Amsterdam. The TMAs consist of samples from 94 patients with NSCLC, including 40 cases of adenocarcinoma (ADC), 42 cases of squamous cell carcinoma (SCC) and 12 cases of other type of NSCLC at different stages (Table 1). On each TMA, 3 cases of healthy lung tissue (i.e. adjacent normal tissue) were also included.

### Mining of publicly available RNA microarray dataset

The datasets were derived from Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/gds>) using the R2 microarray analysis and visualization platform (<http://r2.amc.nl>). Correlation of gene expression between PAR-1 and markers of different stromal cell types in NSCLC cancer patients were derived by the R2 program from five different datasets, including Bild ( $n = 114$ , GSE3141), Peitsch ( $n = 150$ , GSE43580), EXPO ( $n = 121$ , GSE2109), Mao ( $n = 124$ , GSE 31852) and Hou ( $n = 156$ , GSE 19188).

**Table 1** Patient characteristics

Characteristic	Patient	
	N	%
Male	63	67
Median Age (Range)	66 (30–86)	
Progression	26	36
Tumor type:		
Adenocarcinoma	40	42.5
Squamous cell carcinoma	42	44.7
Other type*	12	12.8
Tumor differentiation:		
Well differentiated	6	10.4
Less differentiated	30	51.7
Little differentiated	14	24.1
Poorly differentiated	8	13.8
NSCLC stage:		
I	53	57.6
II	28	30.4
III	10	10.9
IV	1	1.1
Lymph node metastasis	14	15.2

\*This group includes 2 large cell carcinoma patients, 10 patients with mixed tumor types (for instance adenocarcinoma/bronchioloalveolar carcinoma)

### Immunohistological analysis

Four- $\mu\text{m}$  sections were first deparaffinized and rehydrated. Endogenous peroxidase activity was quenched with 0.3% H<sub>2</sub>O<sub>2</sub> in methanol. PAR-1 staining was performed with a primary antibody specific for PAR-1 (ATAP-2 ;1:200; SC-13503, 24 h at 4 °C, Santa Cruz, San Diego, CA) [19, 20]. A horseradish peroxidase-conjugated polymer detection system (ImmunoLogic, Duiven, the Netherlands) was applied for visualization, using an appropriate secondary antibody and diaminobenzidine staining. Specimens with PAR-1 immunostaining were reviewed jointly at a multi-head microscope by 2 investigators blinded to the patients' clinical status. To evaluate immunohistochemical expression of PAR-1, the intensity of PAR-1 staining was graded by consensus on a scale from 0 to 3 (0 = negative staining; 1 = weakly positive; 2 = moderately positive; 3 = strongly positive). Slides were photographed with a microscope equipped with a digital camera (Leica CTR500).

### Statistics

Statistical analyses were conducted using GraphPad Prism (GraphPad software, San Diego). Comparisons between conditions were analyzed using two tailed unpaired t-tests when the data were normally distributed; otherwise Mann–Whitney analysis was performed. Results are expressed as mean  $\pm$  SEM, *P* values < 0.05 are considered significant.

## Results


### PAR-1 gene expression is correlated with lung tumor stroma activation

To explore the association of PAR-1 expression with the NSCLC stroma, we correlated PAR-1 gene expression levels with specific markers of different stromal cell types, including macrophages, endothelial cells, epithelial cells and (myo) fibroblasts in resected tumor specimens using publicly available microarray datasets. To this end, 3 markers were selected for each stromal cell type, except for (myo) fibroblasts for which we included markers of differentiated fibroblasts and markers for extracellular matrix (ECM) produced by myofibroblasts. Interestingly, tumors with higher PAR-1 levels also displayed elevated expression levels of markers for macrophages, endothelial cells and (myo) fibroblasts on the microarrays. Using the GSE3141 dataset (Fig. 1), PAR-1 gene expression was correlated with all three markers for human monocytes and macrophages, i.e. *CD68* ( $p < 0.01$ ), *CD163* ( $p < 0.001$ ) and *CD14* ( $p < 0.0001$ ) [21]. Correlations with specific vascular endothelial cell markers (e.g. Platelet endothelial cell adhesion molecule (*PECAM*)-1) and fibroblasts markers (e.g. Vimentin (*VIM*) and fibroblast activation protein alpha (*FAP*)) were also significant ( $p < 0.0001$ ), with *r*-values ranging from 0.2 to 0.7. The commonly used differentiation marker for fibroblasts *ACTA2* (gene encoding

for alpha-smooth muscle actin,  $\alpha$ -SMA [22]) and markers for prominent constituents of ECM deposition Collagen, type I, alpha (*COL1A1*) and Fibronectin (*FNI*) were also all correlated with PAR-1 gene expression in the NSCLC specimens (all  $p < 0.01$ ). Intriguingly, PAR-1 expression did not correlate to epithelial (tumor) cell markers Epithelial cell adhesion molecule (*EpCAM*), Cadherin 1 (*CDH1*) and Mucin 1 (*MUC1*). These observed correlations (and lack of correlation in epithelial cells) were confirmed in four additional independent microarray datasets from NSCLC (Table 2). However, no correlation between PAR-1 and stromal markers was observed in the healthy control group included in the Hou *et al.* set (GSE19188), suggesting the correlation between PAR-1 gene expression and stroma activity specifically exists in tumor micro-environment. To confirm the identity of the stromal cell types expressing PAR-1, we performed immunohistochemistry with different cell type markers on consecutive lung cancer slides. As shown in Additional file 1: Figure S1, PAR-1 positive areas are also positive for CD31 (endothelial marker), CD68 (macrophage marker) and  $\alpha$ SMA (myofibroblast marker).

### PAR-1 is overexpressed in stroma of primary pulmonary tumors on TMAs

To confirm the presence of PAR-1 in NSCLC stroma, we next analyzed PAR-1 protein expression in tumor sections using immunohistochemistry. Ninety-four patients with pathologically confirmed diagnosis of NSCLC were included into this study. The median age at diagnosis was 66 years (range 30 to 86 years), and the majority of patients had NSCLC stage I disease ( $n = 53$ , 57.6%). Six cases were well differentiated (2 ADC, 1 SCC, 3 other types), 30 cases were moderately differentiated (12 ADC, 18 SCC) and 22 cases were poorly differentiated (10 ADC, 11 SCC, 1 other types) (Table 1). Overall, strong PAR-1 expression was seen in stroma of all different types of NSCLC (ADC, SCC and large-cell carcinoma) as opposed to weak PAR-1 staining on control sections (Fig. 2). In line with our observations in the tumor microarray datasets, the stromal cells (fibroblast-like cells, inflammatory cells and endothelial cells) were all intensively stained for PAR-1, while cancer cells were negative for PAR-1 or showed only weak PAR-1 staining. Subsequent quantifications showed that 93 out of the 94 cases had PAR-1 expression in the stroma, with an average score of 2, while 1 SSC patient was PAR-1 negative. Importantly, the average PAR-1 score in control lungs was significantly lower as in NSCLC stroma (average score of 1; Fig. 3a). As shown in Fig. 3b, PAR-1 levels were similar in different subtype of NSCLC (average scores of 2.11, 2.01 and 2.08 for ADC, SCC and other type of NSCLCs respectively). Stromal PAR-1 expression levels did not correlate with clinical variables like stage of NSCLC (Fig. 3c),



**Fig. 1** PAR-1 expression correlates with stromal markers in NSCLC patients. Scatter plot of PAR-1 gene (F2R) expression versus the expression of specific macrophage (a), endothelial (b), epithelial (c) and (myo) fibroblast (d) markers in tumors derived from NSCLC patients (Bild microarray dataset; GSE3141,  $n = 114$ ). Linear regression analysis was used to determine the correlation coefficient, and  $p$ -values of significant correlations are indicated in red

differentiation status (Fig. 3d), disease progression (Fig. 3e) and overall survival (Fig. 3f).

### Discussion

One of the anticipated future treatment options for NSCLC is to target the interactions between tumor and stromal

cells, since stromal cells provide additional signals that support tumor growth and invasion [1, 16]. In the present study, we determined PAR-1 expression in NSCLC patients and found high PAR-1 expression predominantly in the tumor stroma compartment during early stage cancer. This was reflected by the correlation of PAR-1 gene expression

**Table 2** Correlation of gene expression between PAR-1 and markers of different stromal cell types

Stroma		Database	Bild (n=114)		Peitsch (n=150)		EXPO (n=121)		Mao (n=124)		Hou (n=91)		(nc=65)	
		Gene	P-value	R-value	P-value	R-value	P-value	R-value	P-value	R-value	P <sub>n</sub> -value	R <sub>n</sub> -value	P <sub>nc</sub> -value	R <sub>nc</sub> -value
Macrophage	CD68	6.4e-03	0.254	8.3e-04	0.270	0.03	0.198	0.03	0.191	2.5e-04	0.375	0.06	-0.238	
	CD14	2.9e-05	0.381	3.3e-07	0.402	0.02	0.213	0.82	0.020	6.7e-03	0.282	0.21	-0.157	
	CD163	2.0e-04	0.342	2.6e-05	0.336	0.05	0.175	1.5e-07	0.450	2.3e-03	0.316	0.24	-0.149	
Endothelium	PECAM-1	1.8e-06	0.430	8.3e-05	0.316	2.0e-06	0.417	3.2e-06	0.404	0.36	0.098	0.45	-0.095	
	CDH5	3.3e-04	0.331	7.9e-03	0.216	5.1e-04	0.311	6.4e-07	0.429	0.02	0.239	0.15	0.183	
	vWF	1.8e-03	0.289	9.8e-05	0.313	5.7e-03	0.250	1.1e-05	0.383	1.5e-03	0.328	0.53	0.080	
Epithelium	EpCAM	0.10	-0.154	2.0e-04	-0.299	0.04	-0.188	0.09	-0.151	1.3e-03	-0.333	0.19	0.166	
	CDH1	0.11	-0.149	1.8e-07	-0.410	0.62	-0.046	0.04	-0.182	0.05	-0.207	0.61	0.064	
	MUC1	0.99	0.001	0.33	0.079	0.96	0.005	0.07	-0.165	0.39	-0.091	0.15	-0.182	
(Myo)Fibroblasts	VIM	4.6e-13	0.612	2.6e-11	0.510	1.8e-03	0.281	4.9e-07	0.433	4.4e-08	0.536	0.70	0.048	
	PDGFRB	9.4e-09	0.506	8.1e-12	0.521	3.3e-05	0.368	1.6e-17	0.670	2.2e-05	0.429	0.02	0.286	
	FAP	2.4e-05	0.384	7.0e-12	0.522	9.2e-04	0.297	7.5e-08	0.460	2.1e-05	0.430	0.02	0.299	
	ACTA2	6.0e-08	0.481	4.2e-08	0.429	8.7e-06	0.392	8.9e-16	0.642	2.6e-03	0.312	0.20	0.159	
	COL1A1	5.7e-03	0.257	9.6e-07	0.388	1.0e-05	0.389	1.0e-12	0.585	0.24	0.124	0.05	0.249	
	FN1	6.4e-09	0.511	3.3e-07	0.402	7.8e-07	0.431	1.6e-10	0.535	1.9e-05	0.432	0.23	0.150	

Selection of the datasets is based on the patient group size  $\geq 50$ . Bild ( $n = 114$ , GSE3141), Peitsch ( $n = 150$ , GSE43580), EXPO ( $n = 121$ , GSE2109), Mao ( $n = 124$ , GSE 31852), Hou ( $n = 91$ , control group nc = 65, GSE 19188). Linear regression analysis was used to determine  $p$ -value of correlation. Significant correlations are indicated in red


with stroma markers like CD163, CD31 and vimentin, and ECM proteins like collagen and fibronectin, as well as by a significant increase in the intensity of PAR-1 staining in stromal cells of tumor tissue compared with normal lung tissue. Although it has been documented that upregulation of PAR-1 expression appears in a variety of invasive cancers of epithelial origin, our data do show that increased PAR-1 expression in NSCLC patients arises mainly in the tumor stroma rather than in the epithelial cancer cells.

The observed PAR-1 expression pattern in NSCLC resembles that seen in other malignancies. In breast cancer, PAR-1 expression, as shown by immunohistochemistry and *in situ* hybridization, is observed in mast cells, macrophages, endothelial cells, and vascular smooth muscle cells of the metastatic tumor microenvironment. Interestingly however, PAR-1 expression is particularly increased in stromal fibroblasts surrounding breast carcinoma cells as opposed to low/negative expression in fibroblasts of healthy or benign conditions [23]. Moreover, in prostate cancer PAR-1 is predominantly expressed in peritumoral stroma. In particular, PAR-1 is mainly expressed in myofibroblasts and to a lower level in endothelial cells in isolated capillaries around the malignant glands [24, 25].

The enrichment of PAR-1 expression in the stroma surrounding the tumor may actually be clinically relevant. Indeed, in the setting of pancreatic cancer, PAR-1 also

coincides with the expression pattern of the stromal markers, such as vimentin, collagen I and  $\alpha$ -SMA [18]. More importantly, PAR-1 promoted monocyte recruitment due to fibroblast dependent chemokine production, thereby driving pancreatic tumor growth and chemoresistance [18]. In the context of lung cancers, the expression of PAR-1 mRNA in alveolar walls with surface spreading of neoplastic cells was shown to increase by 10-fold compared with alveolar walls without surface spreading of neoplastic cells, and stimulation of PAR-1 led to the proliferation of alveolar capillary endothelial cells, pointing to PAR-1 as a potential regulator in alveolar angiogenesis [26]. Interestingly, accumulating evidence indicates that PAR-1 also exerts pro-inflammatory and pro-fibrotic functions through macrophages and fibroblasts during pulmonary fibroproliferative disease progression [27–29], which may also benefit tumor progression and metastasis.

Previous studies about PAR-1 in NSCLC focused on its function in cancer cells. Indeed, multiple reports showed that PAR-1 modulates lung cancer cell proliferation and migration, thereby supporting tumor growth and invasion [10, 30]. Hence, targeting PAR-1 to inhibit progression of lung cancer cells seems to be an option for cancer therapy. Recently, emphasis has shifted toward the tumor stroma for novel therapeutic strategies and several approaches targeting the stromal tissue in different types of cancers




**Fig. 2** Stromal PAR-1 expression is upregulated in NSCLC patients. Representative PAR-1 staining of normal lung tissue and tumor sections of NSCLC patients (Pictures were taken with 100x magnification; Enlarged pictures were taken with 200x magnification). ADC indicates adenocarcinoma, SCC indicates squamous cell carcinoma and LCC indicates large cell carcinoma. Tumor cells are indicated by (i) whereas inflammatory cells are indicated by *solid arrowheads*, vascular endothelial cells are indicated by *stars* and fibroblasts-like cells and ECM are indicated by *crosses*

have been proved to be effective [31–33]. Our data showing high stromal PAR-1 expression in NSCLC may thus indicate stromal PAR-1 may be the main target of the treatment for NSCLC. However, before drawing conclusions on potential clinical implications of stromal PAR-1 in NSCLC, it is important to elucidate the functional consequence of PAR-1 activation on stromal cells with respect to lung cancer development.

In the present study, we observed that PAR-1 expression is highly upregulated in the tumor stroma but not in normal lung tissue, suggesting that PAR-1 may have a diagnostic value in NSCLC. However, the increased PAR-1 expression does not seem to correlate with diseases

progression, which indicates that stromal PAR-1 in lung cancer is crucial for carcinogenesis but may not be a determinant factor for cancer progression. These results are in line with a recent study by Erturk and colleagues, who determined serum PAR-1 levels in 80 patients with lung cancer [34]. Serum PAR-1 concentrations of lung cancer patients were significantly increased as compared to controls (i.e. median values of 26.45 ng/mL and 0.07 ng/mL, respectively), but serum PAR-1 levels did not correlate with clinical variables and failed to predict prognosis of the lung cancer patients. In apparent disagreement, other studies using immunohistochemistry analysis showed that PAR-1 may be a prognostic factor for poor prognosis in



**Fig. 3** Association of stromal PAR-1 expression with clinical parameters in NSCLC patients. **a** Stromal PAR-1 expression in healthy lung tissue and in NSCLC. **b** Stromal PAR-1 expression in healthy lung tissue and in different types of NSCLC. **c** Stromal PAR-1 expression in healthy lung tissue and in different stages of NSCLC. **d** Stromal PAR-1 expression according to the differentiation status of NSCLC, including well differentiated, moderately differentiated and poorly differentiated. **e** Stromal PAR-1 expression in NSCLC patients with disease progression and in patients with stable disease (no-progression). **f** Stromal PAR-1 expression of survivors and non-survivors of NSCLC. All data are expressed as mean  $\pm$  SEM, \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$

both early-stage and advanced stages (III and IV) of NSCLC [35, 36]. Importantly however, these studies analyzed tumor cell PAR-1 expression and did not address PAR-1 expression in the stromal compartment.

## Conclusion

In summary, our data show PAR-1 is overexpressed in the tumor stroma of NSCLC, but stromal PAR-1 expression levels do not correlate with disease progression and/or overall survival.

## Additional file

**Additional file 1: Figure S1.** Correlation of PAR-1 expression and specific markers for endothelial cells, macrophages and myofibroblasts. Consecutive lung cancer slides stained for PAR-1 (left panels), CD31 (endothelial marker), CD68 (macrophage marker) and  $\alpha$ SMA (myofibroblast marker). Please note that due to the use of consecutive slides, the structure of the tissue in the PAR-1 stained slide is somewhat different from the CD31, CD68 and  $\alpha$ SMA stained slides. Pictures were taken with 100x magnification. (TIF 7026 kb)

## Abbreviations

ADC: Adenocarcinoma; CDH1: Cadherin-1; COL1A1: Collagen, type I, alpha; ECM: Extracellular matrix; EpCAM: Epithelial cell adhesion molecule; FAP: Fibroblast activation protein alpha; FN1: Fibronectin 1; MUC1: Mucin 1; NSCLC: Non-small-cell lung cancer; PAR-1: Protease activated receptor-1; PECAM-1: Platelet endothelial cell adhesion molecule-1; SCC: Squamous cell carcinoma; siRNA: Small interfering RNA; TMA: Tissue microarray; VIM: Vimentin;  $\alpha$ -SMA: Alpha-smooth muscle actin

## Acknowledgements

Not applicable.

## Funding

This study was supported by grant from the Netherlands Organization for Scientific Research (016.136.167). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Availability of data and materials

The microarray datasets analyzed during the current study were derived from the Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/gds>) using the R2 microarray analysis and visualization platform (<http://r2.amc.nl>). The five microarray datasets include Bild ( $n = 114$ , GSE3141), Peitsch ( $n = 150$ , GSE43580), EXPO ( $n = 121$ , GSE2109), Mao ( $n = 124$ , GSE 31852) and Hou ( $n = 156$ , GSE 19188). The data obtained from TMAs and biopsies are available upon reasonable request from the corresponding author.

## Authors' contributions

CL conceived and designed the experiments, performed the experiments, analyzed the data and wrote the manuscript; CJM performed part of the experiments and analyzed the data; JJTH performed part of the experiments and analyzed the data; MDK analyzed the data; HMH performed part of the experiments; KB analyzed the data and wrote the manuscript; CAS conceived and designed the experiments, and was a major contributor in writing the manuscript. All authors have read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

This research project used anonymized human tissue (both NSCLC tumorous and adjacent healthy tissue) that was removed from a patient during the normal course of treatment and which was later made available for scientific research (so-called 'further use' of human tissue). According to the Code of Conduct for dealing responsibly with human tissue in the context of health research (Human Tissue and Medical Research: Code of conduct for responsible use drawn up by the Federation of Dutch Medical Scientific Societies in

collaboration with the Dutch Patient Consumer federation, the Federation of Parent and Patient Organisations and the Biobanking and Biomolecular Resources Research Infrastructure; [https://www.federa.org/sites/default/files/digital\\_version\\_first\\_part\\_code\\_of\\_conduct\\_in\\_uk\\_2011\\_12092012.pdf](https://www.federa.org/sites/default/files/digital_version_first_part_code_of_conduct_in_uk_2011_12092012.pdf)) these biological materials are as such not subject to any requirement for ethical review or consent from patients.

#### Author details

<sup>1</sup>Center for Experimental and Molecular Medicine, Academic Medical Center, Amsterdam 1105 AZ, The Netherlands. <sup>2</sup>Department of Respiratory Medicine, Academic Medical Center, Amsterdam 1105 AZ, The Netherlands. <sup>3</sup>Department of Pathology, Academic Medical Center, Amsterdam 1105 AZ, The Netherlands. <sup>4</sup>Department of Pulmonology, Zuyderland Hospital, Henri Dunantstraat 5, 6419 PC Heerlen, The Netherlands. <sup>5</sup>Department of Pathology, The Antonie van Leeuwenhoek hospital, Amsterdam 1066 CX, The Netherlands. <sup>6</sup>Inserm UMR1152, Medical School Xavier Bichat, 16 rue Henri Huchard, 75018 Paris, France. <sup>7</sup>Département Hospitalo-universitaire FIRE (Fibrosis, Inflammation and Remodeling) and LabEx Inflammex, Paris, France.

Received: 3 August 2016 Accepted: 23 January 2017

Published online: 07 February 2017

#### References

- Chen Z, Fillmore CM, Hammerman PS, Kim CF, Wong KK. Non-small-cell lung cancers: a heterogeneous set of diseases. *Nat Rev Cancer*. 2014;14:535–46.
- Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136:E359–86.
- Ettinger DS, et al. NCCN guidelines insights: Non-small cell lung cancer, version 4.2016. *J Natl Compr Canc Netw*. 2016;14:255–64.
- Coughlin SR. Thrombin signalling and protease-activated receptors. *Nature*. 2000;407:258–64.
- Vu TK, Hung DT, Wheaton VI, Coughlin SR. Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell*. 1991;64:1057–68.
- Boire A, Covic L, Agarwal A, Jacques S, Sherifi S, Kuliopulos A. PAR1 is a matrix metalloprotease-1 receptor that promotes invasion and tumorigenesis of breast cancer cells. *Cell*. 2005;120:303–13.
- Grisaru-Granovsky S, Salah Z, Maoz M, Pruss D, Beller U, Bar-Shavit R. Differential expression of protease activated receptor 1 (Par1) and pY397FAK in benign and malignant human ovarian tissue samples. *Int J Cancer*. 2005;113:372–8.
- Massi D, Naldini A, Ardinghi C, Carraro F, Franchi A, Paglierani M, Tarantini F, Ketabchi S, Cirino G, Hollenberg MD, Geppetti P, Santucci M. Expression of protease-activated receptors 1 and 2 in melanocytic nevi and malignant melanoma. *Hum Pathol*. 2005;36:676–85.
- Black PC, Mize GJ, Karlin P, Greenberg DL, Hawley SJ, True LD, Vessella RL, Takayama TK. Overexpression of protease-activated receptors-1, 2, and 4 (PAR-1, -2, and -4) in prostate cancer. *Prostate*. 2007;67:743–56.
- Cisowski J, O'Callaghan K, Kuliopulos A, Yang J, Nguyen N, Deng Q, Yang E, Fogel M, Tressell S, Foley C, Agarwal A, Hunt 3rd SW, McMurry T, Brinckerhoff L, Covic L. Targeting protease-activated receptor-1 with cell-penetrating pepsiducins in lung cancer. *Am J Pathol*. 2011;179:513–23.
- Fujimoto D, Hirono Y, Goi T, Katayama K, Yamaguchi A. Prognostic value of protease-activated receptor-1 (PAR-1) and matrix metalloproteinase-1 (MMP-1) in gastric cancer. *Anticancer Res*. 2008;28:847–54.
- Villares GJ, Zigler M, Wang H, Melnikova VO, Wu H, Friedman R, Leslie MC, Vivas-Mejia PE, Lopez-Berestein G, Sood AK, Bar-Eli M. Targeting melanoma growth and metastasis with systemic delivery of liposome-incorporated protease-activated receptor-1 small interfering RNA. *Cancer Res*. 2008;68:9078–86.
- Zigler M, Kamiya T, Brantley EC, Villares GJ, Bar-Eli M. PAR-1 and thrombin: the ties that bind the microenvironment to melanoma metastasis. *Cancer Res*. 2011;71:6561–6.
- Reck M, Heigener DF, Mok T, Soria JC, Rabe KF. Management of non-small-cell lung cancer: recent developments. *Lancet*. 2013;382:709–19.
- El-Nikhely N, Larzabal L, Seeger W, Calvo A, Savai R. Tumor-stromal interactions in lung cancer: novel candidate targets for therapeutic intervention. *Expert Opin Investig Drugs*. 2012;21:1107–22.
- Quail DF, Joyce JA. Microenvironmental regulation of tumor progression and metastasis. *Nat Med*. 2013;19:1423–37.
- Adams GN, Rosenfeldt L, Frederick M, Miller W, Waltz D, Kombrinck K, McElhinney KE, Flick MJ, Monia BP, Revenko AS, Palumbo JS. Colon cancer growth and dissemination relies upon thrombin, stromal PAR-1, and fibrinogen. *Cancer Res*. 2015;75:4235–43.
- Queiroz KC, Shi K, Duitman J, Aberson HL, Wilmink JW, van Noesel CJ, Richel DJ, Spek CA. Protease-activated receptor-1 drives pancreatic cancer progression and chemoresistance. *Int J Cancer*. 2014;135:2294–304.
- Weinstein JR, Lau AL, Brass LF, Cunningham DD. Injury-related factors and conditions down-regulate the thrombin receptor (PAR-1) in a human neuronal cell line. *J Neurochem*. 1998;71:1034–50.
- O'Brien PJ, Prevost N, Molino M, Hollinger MK, Woolkalis MJ, Woulfe DS, Brass LF. Thrombin responses in human endothelial cells. Contributions from receptors other than PAR1 include the transactivation of PAR2 by thrombin-cleaved PAR1. *J Biol Chem*. 2000;275:13502–9.
- Lau SK, Chu PG, Weiss LM. CD163: a specific marker of macrophages in paraffin-embedded tissue samples. *Am J Clin Pathol*. 2004;122:794–801.
- Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol*. 2002;3:349–63.
- D'Andrea MR, Derian CK, Santulli RJ, Andrade-Gordon P. Differential expression of protease-activated receptors-1 and -2 in stromal fibroblasts of normal, benign, and malignant human tissues. *Am J Pathol*. 2001;158:2031–41.
- Zhang X, Wang W, True LD, Vessella RL, Takayama TK. Protease-activated receptor-1 is upregulated in reactive stroma of primary prostate cancer and bone metastasis. *Prostate*. 2009;69:727–36.
- Wang W, Mize GJ, Zhang X, Takayama TK. Kallikrein-related peptidase-4 initiates tumor-stroma interactions in prostate cancer through protease-activated receptor-1. *Int J Cancer*. 2010;126:599–610.
- Lin E, Fujiwara M, Pan X, Ghazizadeh M, Arai S, Ohaki Y, Kajiwara K, Takemura T, Kawanami O. Protease-activated receptor (PAR)-1 and PAR-2 participate in the cell growth of alveolar capillary endothelium in primary lung adenocarcinomas. *Cancer*. 2003;97:703–13.
- Lin C, Rezaee F, Waasdorp M, Shi K, van der Poll T, Borensztajn K, Spek CA. Protease activated receptor-1 regulates macrophage-mediated cellular senescence: a risk for idiopathic pulmonary fibrosis. *Oncotarget*. 2015;6:35304–14.
- Howell DC, Johns RH, Lasky JA, Shan B, Scotton CJ, Laurent GJ, Chambers RC. Absence of proteinase-activated receptor-1 signalling affords protection from bleomycin-induced lung inflammation and fibrosis. *Am J Pathol*. 2005;166:1353–65.
- Lin C, Duitman J, Daalhuisen J, Ten Brink M, von der Thüsen J, van der Poll T, Borensztajn K, Spek CA. Targeting protease activated receptor-1 with P1pal-12 limits bleomycin-induced pulmonary fibrosis. *Thorax*. 2014;69:152–60.
- Wu Z, Zeng Y, Zhong M, Wang B. Targeting A549 lung adenocarcinoma cell growth and invasion with protease-activated receptor-1 siRNA. *Mol Med Rep*. 2014;9:1787–93.
- Tchou J, Conejo-Garcia J. Targeting the tumor stroma as a novel treatment strategy for breast cancer: shifting from the neoplastic cell-centric to a stroma-centric paradigm. *Adv Pharmacol*. 2012;65:45–61.
- Olive KP, Jacobetz MA, Davidson CJ, Gopinathan A, McIntyre D, Honess D, Madhu B, Goldgraben MA, Caldwell ME, Allard D, Frese KK, Denicola G, Feig C, Combs C, Winter SP, Ireland-Zecchini H, Reichelt S, Howat WJ, Chang A, Dhara M, Wang L, Rückert F, Grützmann R, Pilarsky C, Izeradjene K, Hingorani SR, Huang P, Davies SE, Plunkett W, Egorin M, Hruban RH, Whitebread N, McGovern K, Adams J, Iacobuzio-Donahue C, Griffiths J, Tuveson DA. Inhibition of hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. *Science*. 2009;324:1457–61.
- Lopez MV, Rivera AA, Viale DL, Benedetti L, Cuneo N, Kimball KJ, Wang M, Douglas JT, Zhu ZB, Bravo AI, Gidekel M, Alvarez RD, Curiel DT, Podhajcer OL. A tumor-stroma targeted oncolytic adenovirus replicated in human ovary cancer samples and inhibited growth of disseminated solid tumors in mice. *Mol Ther*. 2012;20:2222–33.
- Erturk K, Tastekin D, Bilgin E, Tas F, Disci R, Duranyildiz D. Clinical significance of serum protease activated receptor1 levels in patients with lung cancer. *Eur Rev Med Pharmacol Sci*. 2016;20:243–9.
- Ghio P, Cappia S, Selvaggi G, Novello S, Lausi P, Zecchina G, Papotti M, Borasio P, Scagliotti GV. Prognostic role of protease-activated receptors 1 and 4 in resected stage IB non-small-cell lung cancer. *Clin Lung Cancer*. 2006;7:395–400.
- de Meis E, Azambuja D, Ayres-Silva JP, Zamboni M, Pinheiro VR, Levy RA, Monteiro RQ. Increased expression of tissue factor and protease-activated receptor-1 does not correlate with thrombosis in human lung adenocarcinoma. *Braz J Med Biol Res*. 2010;43:403–8.