




# Loss of NFBD1/MDC1 disrupts homologous recombination repair and sensitizes nasopharyngeal carcinoma cells to PARP inhibitors

Zhihai Wang<sup>1</sup>, Wenqi Zuo<sup>1</sup>, Quan Zeng<sup>1</sup>, Yi Qian<sup>1</sup>, Yanshi Li<sup>1</sup>, Chuan Liu<sup>1</sup>, Jue Wang<sup>1</sup>, Shixun Zhong<sup>1</sup>, Youquan Bu<sup>2</sup> and Guohua Hu<sup>1\*</sup> 

## Abstract

**Background:** Nasopharyngeal carcinoma (NPC), a highly invasive tumor, exhibits a distinctive racial and geographic distribution. As options of agents for effective combination chemoradiotherapy for advanced NPC are limited, novel therapeutic approaches are desperately needed. Here the potential of silencing NFBD1 in combination with PARP inhibition as a novel therapeutic strategy for NPC was investigated.

**Methods:** To investigate the function of NFBD1, we created NFBD1-depleted NPC cell lines via lentivirus mediated shRNA, and the colony formation, MTS assay, comet assay and apoptosis analysis were used to evaluate the sensitivity of NFBD1 knockdown on PARP inhibition. The signaling change was assessed by western blot, Immunofluorescence and flow cytometry. Furthermore, Xenografts model was used to evaluate the role of silencing NFBD1 in combination with PARP inhibition.

**Results:** We find that silencing NFBD1 in combination with PARP inhibition significantly inhibits the cell proliferation and cell cycle checkpoint activity, and increases the apoptosis and DNA damage. Mechanistic studies reveal that NFBD1 loss blocks olaparib-induced homologous recombination repair by decreasing the formation of BRCA1, BRCA2 and RAD51 foci. Furthermore, the xenograft tumor model demonstrated significantly increases sensitivity towards PARP inhibition under NFBD1 deficiency.

**Conclusions:** We show that NFBD1 depletion may possess sensitizing effects of PARP inhibitor, and consequently offers novel therapeutic options for a significant subset of patients.

**Keywords:** Nasopharyngeal carcinoma, PARP inhibitor; homologous recombination, NFBD1/MDC1, DNA damage response

## Background

Nasopharyngeal carcinoma (NPC), a highly invasive cancer, is a common highly malignant head and neck cancer derived from the epithelium of nasopharynx. It is prevalent in Southern China, Malaysia, and Singapore [1, 2]. Although technical improvements in diagnostic technology and clinical treatment, including radiotherapy and chemotherapy, local recurrences and distant metastasis

often occur in 30–40% of NPC patients at advanced staged, and majority of patients will also ultimately die of their disease [3].

Poly (ADP-ribose) polymerase (PARP) is a nuclear enzyme that senses DNA single strand breaks (SSBs). When PARP is inhibited, SSBs are converted into double-strand DNA breaks (DSBs) through collapse of the replication fork. DSBs can be repaired by homologous recombination (HR) which is a high fidelity, error-free form of DNA repair [4]. BRCA1 and BRCA2 proteins are critical components in the process of homologous recombination repair (HRR) for the repair of

\* Correspondence: [hghcq@sina.com](mailto:hghcq@sina.com); [wzhcqm@163.com](mailto:wzhcqm@163.com)

<sup>1</sup>Department of Otorhinolaryngology, the First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China  
Full list of author information is available at the end of the article



DSBs, in BRCA-deficient tumors, HRR is not functional, and therefore the cell is hypersensitive to PARP inhibitors [5–7]. However, PARP inhibitors could also potentially be used as agents that enhance chemo- or radiotherapy-induced DNA damage in patients without defined gene mutations [8]. Therefore, the other mutations/deletions in DNA damage repair genes which have been used to enhance the sensitivity of PARP inhibitors have being widely investigated.

NFBD1 (also known as KIAA01770 or MDC1) is an identified nuclear protein that regulates many aspects of the DNA damage-response pathway, such as intra-S phase checkpoint, G2/M checkpoint, and spindle assembly checkpoint [9–11]. Human NFBD1 comprises 2089 amino acid residues, has a predicted molecular weight of ~220 kDa, and contains an FHA (Forkhead Associated) domain two BRCT (BRCA1 carboxy terminal) domains [12]. These are important structures shared by many DNA damage response proteins, such as Chk2, NBS1 and the tumor suppressor BRCA1. Recent studies have shown that NFBD1 is a participant in the early response to DNA damage and its subsequent signaling within cells. NFBD1 exists in a complex with Chk2 and BRCA1 [9, 13], which are proteins involved in the pathway of homologous recombination. Furthermore, the observed nuclear colocalization of NFBD1 with BRCA1 is further suggestive of a role for NFBD1 in homologous recombination. We focused on NFBD1 in this study and showed that NPC cells with NFBD1-deficient are hypersensitive to the PARP inhibitors olaparib. Thus, PARP inhibitors have therapeutic potential in the treatment of NFBD1-deficient NPC, and our results might extend the concept of synthetic lethality to tumors bearing alterations in NFBD1.

## Methods

### Cell lines and reagents

CNE1, CNE2 and HNE1 were obtained from the Molecular Medicine and Cancer Research Center, Chongqing Medical University. The cells were grown in RPMI-1640 medium (HyClone, Logan City, Utah, USA) with 10% fetal bovine serum (HyClone, Logan City, Utah, USA) at 37 °C with 5% CO<sub>2</sub>. The lentivirus-mediated shNFBD1 and shControl were purchased from Genechem, Shanghai,

China. PARP inhibitor Olaparib (AZD2281) was obtained from MedChemExpress (Princeton, NJ, USA). Hoechst 33342 were purchased from Beyotime Institute of Biotechnology (Nantong, China). The antibodies used in this study were anti-NFBD1 (Abcam, UK); anti-RAD51, anti-BRCA1, anti-BRCA2, and anti-PARP1 (Santa Cruz Biotechnology, USA); anti- $\gamma$ -H2AX (Cell Signaling Technology, Danvers, MA, USA).

### Lentivirus infection

The lentiviral transduction was performed as previously described [11, 14–16]. Cells were transferred into six-well plates, and then viral supernatants were added. The transfected cells of stable expression shNFBD1 and shControl were obtained under puromycin (1  $\mu$ g/ml).

### RNA extraction and real-time quantitative RT-PCR (qRT-PCR)

Total cellular RNA was extracted using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. One microgram of total RNA was used to synthesize cDNA using the One-Step SYBR PrimeScript™ RT-PCR Kit II (TaKaRa Biotechnology, Dalian, China). The qRT-PCR was performed using SYBR Premix Ex Taq in a LightCycler 480 qRT-PCR system (Bio-Rad, Hercules, CA, USA). The qRT-PCR primers of NFBD1 were NFBD1-F (AGCAACCCAGTTGTCATT C) and NFBD1-R (TCCACCACCCTGTTGCTGTA). The  $2^{-\Delta\Delta C_t}$  method was used to determine the relative quantification of NFBD1 expression.

### Clonogenic survival and cytotoxicity assays

For clonogenic survival assay, 500 cells were seeded in 6-well plates in triplicate. After 24 h, the cells were treated continuously with various concentrations of olaparib for another 24 h. After 10–14 days, colonies were fixed with methanol, and stained with crystal violet solution. Colonies containing 50 or more cells were counted as survivors. For cytotoxicity assays, cells were seeded in 96 well plates, and subjected to olaparib treatment as indicated for 48 h. Cell growth was examined using the Promega MTS assay [14–16].

### Flow cytometry

Cells were exposed to olaparib or vehicle for 24 h. For cell cycle analysis, cells were washed in PBS, fixed in 70% ice-cold ethanol at 4 °C overnight and stained with 50  $\mu$ g/ml propidium iodide (PI) solution containing 0.2% Triton X-100 and 100  $\mu$ g/ml DNase-free RNase A. For apoptosis analysis, cells were harvested and stained using Annexin V-FITC Apoptosis Detection kit (Beyotime Institute of Biotechnology, Nantong, China) and according to manufacturer's recommendation. For  $\gamma$ -H2AX, RAD51 and G2/M checkpoint analysis, cells were fixed with ethanol, re-suspended in PBS containing 0.25% (vol/vol) Triton X-100, incubated on ice for 15 min and then incubated in  $\gamma$ -H2AX, RAD51 or phospho-histone H3 (Ser10) antibody for 1 h at room temperature. Samples were then incubated for 30 min at room temperature with secondary antibodies and were determined analyzed by a FACSVantage SE system (BD Biosciences).

### Hoechst 33342 staining

Cells were cultured, exposed to olaparib for 24 h, and then fixed with 4% paraformaldehyde for 1 h at room temperature, and stained with Hoechst 33342 in the dark for 30 min. Images were acquired under a Leica MD2700M fluorescence microscope (German).

### Comet assay

Comet assays were performed as described elsewhere [11]. The Comet Assay kit (Trevigen Inc., Gaithersburg, MD, USA) was used under alkalic conditions according to the manufacturer's specifications. Comets were visualized using a Leica MD2700M fluorescence microscope (German). The tail moments (TMs) of comets were scored using CASP software.

### DR-GFP for HR assay

Cells were transfected the direct-repeat (DR)-GFP plasmid (Addgene, Watertown, MA, USA) by TransIn EL Transfection Reagent (TransGen Biotech, Beijing, China), and the stably expressing the DR-GFP construct were required by G418 screening, and subsequently transiently transfected with pCBASce expression vector (Addgene, Watertown, MA, USA) using TransIn EL Transfection Reagent. GFP signal was assayed 48-h post-transfection on a FACSVantage SE system (BD Biosciences).

### Immunofluorescence

Immunofluorescence were performed as described elsewhere [11, 15]. Briefly, Cells were fixed, permeabilized, washed, blocked, and then primary antibodies were applied overnight at 4 °C, and secondary antibodies were applied for 60 min at room temperature. Finally, the cells were washed three times in PBS, and the DNA was stained using DAPI (Sigma-Aldrich, St. Louis, MO, USA) at 5 ng/ml. The slides were observed under a Leica MD2700M fluorescence microscope (German).

### Western blotting and coimmunoprecipitation

Total protein extracts from cells were prepared using RIPA buffer (Beyotime Institute of Biotechnology, Nantong, China). Proteins were fractionated in SDS-polyacrylamide gels, transferred to polyvinylidene fluoride (Millipore, Billerica, MA, USA), and western blotting were performed by using the appropriate antibody. Antibody/antigen complexes were detected using ECL (Western Bright Sirius; Advansta, Inc., Menlo Park, CA, USA) and images were acquired using an enhanced chemifluorescence detection system (Amersham Biosciences, Piscataway, NJ, USA) under the room temperature.

For coimmunoprecipitation, Total protein lysates (0.5 mg) were incubated with 4 µg of specific anti-NFBD1 antibody overnight at 4 °C. NFBD1 immunocomplexes

were captured with SureBeads™ Magnetic Beads (Bio-Rad, Hercules, CA, USA) for 1 h at 4 °C. The resulting immunocomplexes were collected by centrifugation, were washed, boiled in SDS sample buffer, loaded on an SDS-polyacrylamide gel. Proteins were analysed by western blotting using standard methods and detected as described above.

### Xenograft experiments

All animal husbandry and experiments were performed under a protocol approved by Institutional Animal Care Committee at Chongqing Medical University. CNE1 cells ( $6.0 \times 10^6$ ) in 0.2 ml of growth medium were subcutaneously injected into the axilla of the Balb/c nude mice. Two days after transplantation success (the diameter of tumor was approximately 5 mm), mice were treated daily with either vehicle or 50 mg/kg bodyweight of olaparib intraperitoneally five days a week. Tumor size was measured every week and tumor volume =  $1/2 \times \text{length} \times \text{width}^2$ . Mice were killed after 42 days after cell injection, and then the tumors in the left and right axillary region were excised and weighed.

### Statistics

Statistical comparison of mean values was performed using ANOVA, rank sum test (non-parametric statistics) or chi-square ( $\chi^2$ ) test. Differences with a *P*-value of < 0.05 were considered statistically significant.


## Results

### Lentivirus-mediated shRNA inhibited NFBD1 mRNA and protein expression in NPC cells

The lentiviral expressing NFBD1 shRNA and control shRNA were transfected into NPC cells. The transfected cells of stable expression shNFBD1 and shControl were obtained under puromycin (1 µg/ml). NFBD1 downregulation was confirmed by qRT-PCR and western blotting analysis (Fig. 1a and b). In addition, immunofluorescence also revealed corresponding decrease in the protein levels (Fig. 1c). These results demonstrated that the lentivirus-mediated shRNA targeting NFBD1 effectively knocked down NFBD1 expression at both mRNA and protein levels in the NPC cells.

### Loss of NFBD1 enhances NPC cells to be sensitive to PARP inhibitor olaparib

To evaluate the role of NFBD1 in the response of NPC cells to PARP inhibitor-induced DSBs, clonogenic assays were also performed to determine the effects of the NFBD1 knockdown on the sensitivity of NPC cells to olaparib. NPC cells with stable NFBD1 knockdown were markedly sensitive to olaparib compared to control cells (Fig. 2a). In proliferation assays, silencing NFBD1 significantly increased the sensitivity upon olaparib treatment



compared to controls, furthermore, NFBD1 knockdown resulted in a marked decrease of IC<sub>50</sub> values (Fig. 2b).


#### NFBD1 participates in the regulation of olaparib-induced cell cycle distribution and G<sub>2</sub>/M checkpoint activity

The sensitivity to olaparib suggests an important role in responding to DNA damage; therefore we next tested whether NFBD1 loss could change olaparib-induced cell cycle distribution. shControl and shNFBD1 cells were exposed to olaparib and the cell cycle distribution was determined by flow cytometry (FCM). Control cells exhibited a decrease in the rate of G<sub>1</sub> phase, whereas an increase in the rate of S phase compared with shNFBD1 cells (Fig. 3a). This demonstrates a role for NFBD1 in regulating cell cycle progression after DNA damage. We next examined the integrity of the G<sub>2</sub>/M checkpoint. Cells were exposed to olaparib and labelled with an anti-phospho-histone H3 antibody as a marker of mitotic cells. A clear reduction in phospho-H3-positive cells was observed in the shControl cells after olaparib treatment, whereas a significant number of the cells lacking NFBD1


entered mitosis (Fig. 3b), indicative of a defect in the ability to arrest the cell cycle in G<sub>2</sub> phase.

#### Deficiency in NFBD1 causes defective DSB repair and confers enhanced apoptosis to the PARP inhibitor olaparib

To determine whether NFBD1 knockdown affects DNA repair post olaparib treatment, we measured the persistence of DNA damage in olaparib-treated NPC cells utilizing comet assay (Fig. 4a). Olaparib treatment induced DSBs, visibly by increased DNA mobility or comet tail. The cells lacking NFBD1 revealed significantly increased the intensity and length of comet tails after olaparib treatment compared with control cells. Considering the notion that cells can undergo apoptosis when DNA damage is irreparable, therefore, to examine whether downregulation of NFBD1 following olaparib treatment can induce apoptosis, NPC cells were exposed to olaparib, and determined the apoptosis rate using FCM (Fig. 4b). Olaparib potently induced apoptosis in CNE1 and CNE2 cells in a dose-dependent manner,



**Fig. 2** Knockdown of NFB1 enhances NPC cells to be sensitive to PARP inhibitor olaparib. **(a)** Cells were plated at low density, irradiated and colonies counted after 10–14 days. Results are normalized for effects of NFB1 and fitted to a standard linear quadratic model. **(b)** NFB1 knockdown enhanced the cytotoxicity to olaparib in a dose-dependent manner. shControl and shNFB1 cells were treated with various concentrations of olaparib for 24 h. Cell viability was measured using MTS assays. The data are presented as the mean ± SD of triplicate determinations. IC50 values are presented as the mean ± SD of triplicate determinations. \*\**P* < 0.01 and \*\*\**P* < 0.001




**Fig. 3** The effects of olaparib on cell cycle distribution. NPC cells were treated with various concentrations of olaparib for 24 h, **(a)** the cell cycle distribution was determined by FCM, and **(b)** mitotic cells were determined by phospho-histone H3 staining and FCM. Data shown are mean  $\pm$  SD from three independent experiments. \* $P < 0.05$ \*\*,  $P < 0.01$  and \*\*\* $P < 0.001$

furthermore, NFBD1 knockdown significantly enhanced olaparib-induced apoptosis. To further confirm the above result, we performed Hoechst 33342 staining. Consistent with our FCM results, the shRNA group demonstrated a greater number of fragmented nuclei and nuclear shrinkage compared with the shControl group following olaparib treatment (Fig. 4c).

#### Silencing NFBD1 abrogates olaparib-induced formation of $\gamma$ -H2AX and Rad51 foci

To assess whether NFBD1 could localize to sites of damage, cells were treated with olaparib and stained with

anti-NFBD1 antibody. At the level of single nuclei, olaparib induced punctated and distinct NFBD1 foci (Fig. 5a). A proportion of untreated cells also contained NFBD1 foci, indicating that NFBD1 may be responding to endogenous damage or replication stress. As NFBD1 controls the phosphorylation of several checkpoint-responsive proteins, we sought to examine whether it might have a role in H2AX phosphorylation. Silencing NFBD1 significantly affected the phosphorylation of H2AX after olaparib treatment and the formation of  $\gamma$ -H2AX foci (Fig. 5b), the same result was observed in FCM and western blotting analysis (Fig. 5d and g). Furthermore, NFBD1 foci



also significantly co-localized with  $\gamma$ -H2AX foci (Fig. 5b and c). Thus NFB1 was necessary for H2AX phosphorylation and foci formation after olaparib treatment. Because loss of NFB1 enhances NPC cells to be sensitive to PARP inhibitor olaparib, and the formation of nuclear Rad51 foci is known as the key step of HRR, we ascertained whether NFB1 had effects on Rad51-mediated repair of DSBs. As hypothesized, immunofluorescence analysis showed that RAD51 foci formation in response to olaparib was greatly inhibited in NFB1 loss cells (Fig. 5e), and FCM analysis indicated that more RAD51-positive cells were consistently observed after olaparib treatment in the shControl group than that in shNFB1 group (Fig. 5f). The same result was observed in western blotting analysis (Fig. 5g).


#### Loss of NFB1 disrupts olaparib-induced homologous recombination repair

HRR is a high fidelity, error-free form of DNA repair, and BRCA1 and BRCA2 proteins are critical components in the process of HRR. Cancer cells with deleterious BRCA1/2 mutations are defective in HRR and therefore are hypersensitive to PARP inhibitors [17, 18]. We used direct-repeat (DR)-GFP plasmid to assay the

HR pathway, and the HR recombination assay relies on the DR-GFP transfected cells to express GFP, if HR activity is functional. The results clearly show significant decrease in percentage GFP-positive cells with NFB1 inhibition compared to untreated cells (Fig. 6a). Furthermore, we assess whether silencing NFB1 could affect olaparib-induced the formation of BRCA1 and BRCA2 foci in nuclei. Olaparib induced a marked increase in BRCA1 and BRCA2 foci in shControl cells but not in shNFB1 cells (Fig. 6b and c), the same results were observed in western blotting analysis (Fig. 6e), and NFB1 foci also significantly co-localized with BRCA1 foci (Fig. 6c). In addition, BRCA1 was recruited to the DNA damage sites, and co-localized with  $\gamma$ -H2AX foci following olaparib treatment (Fig. 6d).

#### NFB1 loss results in increased PARP inhibitor sensitivity in vivo


To assess the therapeutic effect of olaparib on NFB1 depleted cells in vivo, we investigated the ability of olaparib to suppress the growth of NFB1-depleted CNE1 cells derived xenograft tumor. shControl or shNFB1 cells were subcutaneously grafted into balb/c nude mice. Two days after transplantation success, mice were




treated with olaparib or vehicle. Six weeks post-treatment, mice were sacrificed and tumor volume measured and quantified. Our results showed that either downregulation of NFBFD1 or olaparib alone resulted in significantly smaller tumor than untreated xenografts, however, the combination of downregulated NFBFD1 and

olaparib resulted in significantly smaller tumors as compared with untreated controls or to tumors treated with downregulated NFBFD1 or olaparib alone (Fig. 7a–c). Thus, the results suggested that silencing NFBFD1 enhanced the response of NPC cells to olaparib and resulted in tumor growth inhibition in vivo.





**Fig. 6** Loss of NFB1 causes HRR deficiency **(a)** Cells stably transfected with a DR-GFP reporter construct express GFP were transfected with I-SceI endonuclease expression vector. Extent of HR was analyzed by FCM 48 h after transfection. **(b and c)** Inhibition of NFB1 resulted in defective BRCA1 and BRCA2 foci formation after olaparib exposure. **(d)** Silencing NFB1 disrupts olaparib-induced BRCA1 interaction with γ-H2AX. **(e)** Cells were treated 4 μg/ml olaparib for 24 h, and subjected to western blotting analysis with indicated antibodies. Representative blots were shown with β-actin as loading control



**Fig. 7** Combination therapy inhibits tumor growth of nasopharyngeal cancer in vivo. **(a and b)** Loss of NFBD1 significantly reduces xenograft growth after PARP inhibitor olaparib treatment. The tumors were measured every week using a digital caliper. Tumor volume =  $1/2 \times \text{length} \times \text{width}^2$ . **(c)** Tumor weight at 42 days after treatments,  $n = 6$  mice per condition. \* $P < 0.05$  and \*\* $P < 0.01$

### Discussion

Cells are continuously exposed to exogenous and endogenous agents that lead to DNA damage, which, if not

repaired effectively and efficiently, can result to genomic instability or cell death [19]. The HRR pathway is an especially important DNA repair pathway in this respect.

HR-deficiency is observed in some cancers, such as breast cancer and ovarian cancer. Interestingly, tumors with defective HR, including BRCA1/2 mutation cancers, exhibit particular sensitivity to PARP inhibitors [20–23]. However, the use of PARP inhibitors is limited to cancer patients with BRCA1/2 mutations, which represent only a small subset of cases. Therefore, new therapeutic strategies and approaches are urgently needed to make broader use of PARP inhibitors for the majority of other tumor cases.

Conceptually, PARP inhibitors can selectively augment DNA damage in tumor with HR-deficiency compared with normal tissues. As implicated by previous studies, NFBBD1 has important roles in DDR, involving a complex network of signaling pathways that regulates cell cycle checkpoints, DNA repair and cell death [11, 15, 24–27]. Our studies showed that NFBBD1 protein is highly expressed in nasopharyngeal carcinoma tissues, and shRNA targeting NFBBD1 enhanced the radiosensitivity of CNE1 cells and impaired DNA damage-induced RAD51 foci formation [11, 15, 16]. This suggested that the rationale for the utility of olaparib is based upon the notion that silencing NFBBD1 disturbs the HRR pathway required for DNA damage repair following olaparib treatment. In this study, silencing NFBBD1 significantly enhanced olaparib-induced apoptosis and growth inhibition of NPC cells. More importantly, NFBBD1 knock-down can delay olaparib-induced DNA damage repair, impair cell cycle checkpoint activity, and enhance the sensitivity of NPC cells to olaparib. Xenografts models showed that silencing NFBBD1 significantly enhanced the anti-tumor activity of olaparib, leading to tumor growth inhibition of the combination therapy. Mechanistic studies revealed that silencing NFBBD1 disturbed the HRR pathway through impairing the foci formation of olaparib-induced BRCA1, BRCA2 and RAD51. These findings provide a molecular basis of using PARP inhibitors to potentiate treatment of nasopharyngeal carcinoma with NFBBD1-deficiency.

It has been argued that cell cycle checkpoint arrest, although important for maintaining genomic stability post DNA damage, makes a less significant contribution to survival. However, when cell cycle checkpoint inactivation is combined with defective DSBs repair, the impact is more than additive, consistent with the notion that cell cycle checkpoint arrest enhances the opportunity for DSBs repair [28]. In our studies, olaparib treatment led to a significant increase in G1 and M phase cell population and a corresponding decrease in S phase for NFBBD1-deficiency NPC cell lines, indicative of a defect in the G1/S and G2/ M checkpoint. Furthermore, the comet assay showed that silencing NFBBD1 significantly increased olaparib-inducing DNA damage. From these data, silencing NFBBD1 enhanced the sensitivity of NPC

cells to olaparib through impairing cell cycle checkpoint activity and improving DNA damage.

On the basis of our findings, we propose that the enhanced sensitivity of NFBBD1-deficiency cells to olaparib is largely due to defect of HRR pathway. In the case of DNA damage,  $\gamma$ -H2AX is ATM-dependent, which is involved in the amplification step required for optimal checkpoint response in the DDR [29, 30]. It is evident that a network of interactions is initiated around  $\gamma$ -H2AX, which recruits and maintains many DDR proteins at sites of DSBs [31–35]. We found that olaparib induced punctated and distinct NFBBD1 and  $\gamma$ -H2AX foci, and BRCA1 was also recruited to the DNA damage sites and colocalized with  $\gamma$ -H2AX. However, knock-down of NFBBD1 greatly reduced olaparib-induced formation of  $\gamma$ -H2AX foci, moreover, our colocalization studies indicated NFBBD1 foci extensively overlap with  $\gamma$ -H2AX foci, suggesting that NFBBD1 was required for the formation of olaparib-induced  $\gamma$ -H2AX foci. In addition, NFBBD1-deficient cells revealed a decrease BRCA1, BRCA2 and RAD51 foci relative to controls, which indicated the use of NFBBD1 loss combination with olaparib may directly inhibit HR-mediated DNA repair through impairing the foci formation of BRCA1, BRCA2 and RAD51. The impairment of DDR proteins foci formation could be due to overall protein level is decreased, and/or HR proteins cannot be recruited to DSB sites. Clearly, the western blotting shows that the depletion of NFBBD1 significantly reduces the protein level of BRCA1, BRCA2, and RAD51. Therefore, these results indicated that downregulation of NFBBD1 can inhibited the amplification of the olaparib-induced DNA damage signal, decreased the levels of DDR proteins, and failed to accumulate and retain DDR proteins at the sites of DNA damage, which led to defective HRR activation following olaparib treatment.

## Conclusions

We report here that NFBBD1 knockdown combination with olaparib can enhance the sensitivity of olaparib in NPC cells. Silencing NFBBD1 also inhibit the amplification of the olaparib-induced DNA damage signal, and fail to accumulation and retain DDR proteins at the sites of DNA damage, which leads to defective checkpoint activation following DNA damage. Furthermore, silencing NFBBD1 inhibits the HRR pathway activity by impairing the foci formation of olaparib inducing BRCA1, BRCA2 and RAD51. Our studies provide compelling evidence that combining depletion of NFBBD1 and olaparib represents a rational strategy for the treatment of patients with NPC.

## Abbreviations

BER: Base excision repair; DDR: DNA damage response; DSBs: DNA double-strand breaks; FCM: Flow cytometry; HR: Homologous recombination; HRR: Homologous recombination repair; NHEJ: Non-homologous end-

joining; NPC: Nasopharyngeal carcinoma; PARP: Poly (ADP-ribose) polymerase; qRT-PCR: Real-time quantitative RT-PCR; SSB: Single-strand break

#### Acknowledgements

We thank Guojun Chen for his technical support in the signaling Pathway experiments.

#### Funding

The work was supported by the National Natural Science Foundation of China (81470676 and 81271061) and Nurture Fund of The First Affiliated Hospital of Chongqing Medical University (2018PYJJ-01).

#### Availability of data and materials

The datasets during and/or analyzed during the current study available from the corresponding author on reasonable request.

#### Authors' contributions

G Hu and Z Wang conceived the study design and analysis plan. Z Wang, G Hu, W Zuo, and Q Zeng wrote the first draft of the report. Y Qian, Z Wang, Y Li, C Liu and J Wang performed the statistical analysis. Y Bu, Z Wang, S Zhong and G Hu jointly directed the study. All authors reviewed the manuscript and provided comments or suggestions. All authors read and approved the final manuscript.

#### Ethics approval and consent to participate

All procedures followed were in accordance with the ethical standards of the responsible committee on animal experimentation and with the Helsinki Declaration of 1975 and later versions. All animal husbandry and experiments were performed under a protocol approved by Institutional Animal Care Committee at Chongqing Medical University.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### Author details

<sup>1</sup>Department of Otorhinolaryngology, the First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China. <sup>2</sup>Department of Biochemistry and Molecular Biology, Molecular Medicine and Cancer Research Center, Chongqing Medical University, Chongqing 400016, China.

Received: 20 August 2018 Accepted: 22 January 2019

Published online: 04 February 2019

#### References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin*. 2011;61:69–90.
- Xu YF, Li YQ, Liu N, He QM, Tang XR, Wen X, Yang XJ, Sun Y, Ma J, Tang LL. Differential genome-wide profiling of alternative polyadenylation sites in nasopharyngeal carcinoma by high-throughput sequencing. *J Biomed Sci*. 2018;25:74.
- Bensouda Y, Kaikani W, Ahbeddou N, Rahhali R, Jabri M, Mrabti H, Boussem H, Errihani H. Treatment for metastatic nasopharyngeal carcinoma. *Eur Ann Otorhinolaryngol Head Neck Dis*. 2011;128:79–85.
- Xu G, Chapman JR, Brandsma I, Yuan J, Mistrik M, Bouwman P, Bartkova J, Gogola E, Warmerdam D, Barazas M, Jaspers JE, Watanabe K, Pieterse M, Kersbergen A, Sol W, Celie PHN, Schouten PC, van den Broek B, Salman A, Nieuwland M, de Rink I, de Ronde J, Jalink K, Boulton SJ, Chen J, van Gent DC, Bartek J, Jonkers J, Borst P, Rottenberg S. REV7 counteracts DNA double-strand break resection and affects PARP inhibition. *Nature*. 2015;521:541–4.
- Labidi-Galy SI, Olivier T, Rodrigues M, Ferraioli D, Bodmer A, Petignat P, Rak B, Chopin N, Tredan O, Heudel PE, Stuckelberger SJ, Meeus P, Meraldi P, Viassolo V, Ayme A, Chappuis PO, Stern MH, Houdayer C, Stoppa-Lyonnet D, Buisson A, Golmard L, Bonadona V, Ray-Coquard I. Location of mutation in BRCA2 gene and survival in patients with ovarian cancer. *Clin Cancer Res*. 2017.
- Maxwell KN, Wubbenhorst B, Wenz BM, De Sloover D, Pluta J, Emery L, Barrett A, Kraya AA, Anastopoulos IN, Yu S, Jiang Y, Chen H, Zhang NR, Hackman N, D'Andrea K, Daber R, Morrisette JJD, Mitra N, Feldman M, Domchek SM and Nathanson KL. BRCA locus-specific loss of heterozygosity in germline BRCA1 and BRCA2 carriers. 2017; 8: 319.
- Zhao W, Steinfeld JB, Liang F, Chen X, Maranon DG, Jian Ma C, Kwon Y, Rao T, Wang W, Sheng C, Song X, Deng Y, Jimenez-Sainz J, Lu L, Jensen RB, Xiong Y, Kupfer GM, Wiese C, Greene EC, Sung P. BRCA1-BARD1 promotes RAD51-mediated homologous DNA pairing. *Nature*. 2017;550:360–5.
- Pignochino Y, Capozzi F, D'Ambrosio L, Dell'Aglio C, Basiricò M, Canta M, Lorenzato A, Vignolo LF, Aliberti S, Palesandro E. PARP1 expression drives the synergistic antitumor activity of trabectedin and PARP1 inhibitors in sarcoma preclinical models. *Mol Cancer*. 2017;16:86.
- Lou Z, Minter-Dykhouse K, Wu X, Chen J. MDC1 is coupled to activated CHK2 in mammalian DNA damage response pathways. *Nature*. 2003;421:957–61.
- Goldberg M, Stucki M, Falck J, D'Amours D, Rahman D, Pappin D, Bartek J, Jackson SP. MDC1 is required for the intra-S-phase DNA damage checkpoint. *Nature*. 2003;421:952–6.
- Wang Z, Zeng Q, Chen T, Liao K, Bu Y, Hong S, Hu G. Silencing NFB1/MDC1 enhances the radiosensitivity of human nasopharyngeal cancer CNE1 cells and results in tumor growth inhibition. *Cell Death Dis*. 2015;6:e1849.
- Ozaki T, Nagase T, Ichimiya S, Seki N, Ohiri M, Nomura N, Takada N, Sakiyama S, Weber BL, Nakagawara A. NFB1/KIAA0170 is a novel nuclear transcriptional transactivator with BRCT domain. *DNA Cell Biol*. 2000;19:475–85.
- Stewart GS, Wang B, Bignell CR, Taylor AMR, Elledge SJ. MDC1 is a mediator of the mammalian DNA damage checkpoint. *Nature*. 2003;421:961–6.
- Liu X, Qiu Z, Wang Z, Zuo W, Gong Z, Liu C, Zeng Q, Qian Y, Jiang L, Li Y, Bu Y and Hu G. NFB1/MDC1 participates in the regulation of proliferation and apoptosis in human laryngeal squamous cell carcinoma. 2017;
- Zeng Q, Wang Z, Liu C, Gong Z, Yang L, Jiang L, Ma Z, Qian Y, Yang Y, Kang H, Hong S, Bu Y, Hu G. Knockdown of NFB1/MDC1 enhances chemosensitivity to cisplatin or 5-fluorouracil in nasopharyngeal carcinoma CNE1 cells. *Mol Cell Biochem*. 2016;418:137–46.
- Wang Z, Liao K, Zuo W, Liu X, Qiu Z, Gong Z, Liu C, Zeng Q, Qian Y, Jiang L, Bu Y, Hong S, Hu G. Depletion of NFB1/MDC1 induces apoptosis in nasopharyngeal carcinoma cells through the p53-ROS-mitochondrial pathway. *Oncol Res*. 2017;25:123–36.
- Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB, Santaros M, Dillon KJ, Hickson I, Knights C, Martin NM, Jackson SP, Smith GC, Ashworth A. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*. 2005;434:917–21.
- Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E, Kyle S, Meuth M, Curtin NJ, Helleday T. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature*. 2005;434:913–7.
- Hoeijmakers JH. Genome maintenance mechanisms for preventing cancer. *Nature*. 2001;411:366–74.
- Helleday T, Petermann E, Lundin C, Hodgson B, Sharma RA. DNA repair pathways as targets for cancer therapy. *Nat Rev Cancer*. 2008;8:193–204.
- Lord CJ, Ashworth A. Targeted therapy for cancer using PARP inhibitors. *Curr Opin Pharmacol*. 2008;8:363–9.
- Kummar S, Chen A, Parchment RE, Kinders RJ, Ji J, Tomaszewski JE, Doroshow JH. Advances in using PARP inhibitors to treat cancer. *BMC Med*. 2012;10:25.
- Livraghi L, Garber JE. PARP inhibitors in the management of breast cancer: current data and future prospects. *BMC Med*. 2015;13:188.
- Bartek J, Lukas J. DNA damage checkpoints: from initiation to recovery or adaptation. *Curr Opin Cell Biol*. 2007;19:238–45.
- Kastan MB, Bartek J. Cell-cycle checkpoints and cancer. *Nature*. 2004;432:316–23.
- Burma S, Chen BP, Murphy M, Kurimasa A, Chen DJ. ATM phosphorylates histone H2AX in response to DNA double-strand breaks. *J Biol Chem*. 2001; 276:42462–7.
- Nakanishi M, Ozaki T, Yamamoto H, Hanamoto T, Kikuchi H, Furuya K, Asaka M, Delia D, Nakagawara A. NFB1/MDC1 associates with p53 and regulates its function at the crossroad between cell survival and death in response to DNA damage. *J Biol Chem*. 2007;282:22993–3004.
- Lobrich M, Jeggo PA. The impact of a negligent G2/M checkpoint on genomic instability and cancer induction. *Nat Rev Cancer*. 2007;7:861–9.

29. Stucki M, Clapperton JA, Mohammad D, Yaffe MB, Smerdon SJ, Jackson SP. MDC1 directly binds phosphorylated histone H2AX to regulate cellular responses to DNA double-strand breaks. *Cell*. 2005;123:1213–26.
30. Wang J, Gong Z, Chen J. MDC1 collaborates with TopBP1 in DNA replication checkpoint control. *J Cell Biol*. 2011;193:267–73.
31. van Attikum H, Fritsch O, Hohn B, Gasser SM. Recruitment of the INO80 complex by H2A phosphorylation links ATP-dependent chromatin remodeling with DNA double-strand break repair. *Cell*. 2004;119:777–88.
32. Morrison AJ, Highland J, Krogan NJ, Arbel-Eden A, Greenblatt JF, Haber JE, Shen X. INO80 and gamma-H2AX interaction links ATP-dependent chromatin remodeling to DNA damage repair. *Cell*. 2004;119:767–75.
33. Downs JA, Allard S, Jobin-Robitaille O, Javaheri A, Auger A, Bouchard N, Kron SJ, Jackson SP, Cote J. Binding of chromatin-modifying activities to phosphorylated histone H2A at DNA damage sites. *Mol Cell*. 2004;16:979–90.
34. Park JH, Park EJ, Lee HS, Kim SJ, Hur SK, Imbalzano AN, Kwon J. Mammalian SWI/SNF complexes facilitate DNA double-strand break repair by promoting gamma-H2AX induction. *EMBO J*. 2006;25:3986–97.
35. Kobayashi J, Fujimoto H, Sato J, Hayashi I, Burma S, Matsuura S, Chen DJ, Komatsu K. Nucleolin participates in DNA double-strand break-induced damage response through MDC1-dependent pathway. *PLoS One*. 2012;7:e49245.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

