

RESEARCH

Open Access



# Homotopy perturbation transform method for nonlinear differential equations involving to fractional operator with exponential kernel

JF Gómez-Aguilar<sup>1\*</sup>, H Yépez-Martínez<sup>2</sup>, J Torres-Jiménez<sup>3</sup>, T Córdova-Fraga<sup>4</sup>, RF Escobar-Jiménez<sup>5</sup> and VH Olivares-Peregrino<sup>5</sup>

\*Correspondence:

jgomez@cenidet.edu.mx

<sup>1</sup>CONACyT-Centro Nacional de Investigación y Desarrollo Tecnológico, Tecnológico Nacional de México, Interior Internado Palmira S/N, Col. Palmira, Cuernavaca, Morelos 62490, México  
Full list of author information is available at the end of the article

## Abstract

This work presents the homotopy perturbation transform method for nonlinear fractional partial differential equations of the Caputo-Fabrizio fractional operator. Perturbative expansion polynomials are considered to obtain an infinite series solution. The effectiveness of this method is demonstrated by finding the exact solutions of the fractional equations proposed, for the special case when the limit of the integral order of the time derivative is considered.

**PACS Codes:** 02.30.Jr; 02.60.Cb; 02.60.Lj

**Keywords:** fractional calculus; nonlinear fractional differential equations; Caputo-Fabrizio fractional operator; homotopy perturbation transform method; approximate solution

## 1 Introduction

Fractional calculus (FC) has become an alternative mathematical method to describe models with nonlocal behavior. In the last decade, considerable interest in fractional differential equations has been stimulated due to their numerous applications in the areas of physics and engineering [1–15]. Several numerical and analytical methods have been developed to study the solutions of nonlinear fractional partial differential equations, fractional sub-equation methods [16–18], the homotopy perturbation methods [19–21], the variational iteration methods [22–27], homotopy perturbation transform methods [28, 29], Adomian decomposition methods [30–33], and other analytical approaches that could be of interest for the reader are presented in [34–38]. It worth noting that there exist only two main definitions of the fractional derivative; the first was proposed by Riemann and Liouville and is the derivative of the convolution of a given function and a power law kernel, the second one was suggested by Caputo and it is the convolution of the local derivative of a given function with power law function [39].

Due to the fact that the power law cannot be used to model all the physical problems, Caputo and Fabrizio [40] have suggested an alternative concept of differentiation using the exponential decay as kernel instead of the power law. This new differentiation has

also attracted attention of many scholars but was also disqualified for classification as a fractional derivative due to the fact that the kernel was not nonlocal; however, it is clear that many problems in nature also follow the exponential decay law which indeed has no singularity; therefore this derivative is significantly useful in modeling such real world problems [41–44].

The homotopy analysis method (HAM), proposed by Liao, has been successfully applied to solving many problems in physics and science [45–48], this method transforms a problem into an infinite number of linear problems without using the perturbation techniques. The Laplace homotopy analysis method (LHAM) is a combination of HAM and Laplace transform [49, 50]. The homotopy perturbation method is also combined with the well-known Laplace transformation method and the variational iteration method to produce a highly effective technique (homotopy perturbation transform method) for handling many nonlinear problems [51].

In this paper, we use the homotopy perturbation transform method (HPTM) to solve nonlinear fractional partial differential equations using the fractional operator of Caputo-Fabrizio type. The basic definitions of fractional calculus are given in Section 2, several test problems that show the effectiveness of the proposed method are given in Section 3, and finally the conclusion is given in Section 4.

## 2 Basic tools

The Liouville-Caputo fractional derivative is defined for  $(\gamma > 0)$  as

$${}_0^C D_t^\mu f(t) = \frac{1}{\Gamma(n - \mu)} \int_0^t (t - s)^{n-\mu-1} f^{(n)}(s) ds, \tag{1}$$

where  ${}_0^C D_t^\mu$  is a Liouville-Caputo fractional derivative with respect to  $t$ ,  $f^{(n)}$  is the derivative of integer  $n$ th order of  $f$ ,  $n = 1, 2, \dots \in \mathbb{N}$ , and  $n - 1 < \mu \leq n$ .

Now, if the kernel  $(t - s)^{n-\mu-1}$  is changed for the function  $\exp(-\mu(t - s)/(1 - \mu))$ , and  $\frac{1}{\Gamma(n-\mu)}$  for  $\frac{(2-\alpha)M(\alpha)}{2(1-\alpha)}$  in equation (1), we can show the new definition of fractional operator proposed by Caputo and Fabrizio (CF), which is expressed as follows [40, 41]:

$${}_0^{CF} D_t^\mu f(t) = \frac{(2 - \mu)M(\mu)}{2(1 - \mu)} \int_0^t \exp\left(\frac{-\mu}{1 - \mu}(t - s)\right) f^{(n)}(s) ds, \tag{2}$$

where  $M(\alpha)$  is a normalization function such that  $M(0) = M(1) = 1$ . This new definition does not have singularities at  $t = s$ .

If  $0 < \mu \leq 1$  and  $n \in \mathbb{N}$ , then we define the Laplace transform for the CF fractional operator as follows [40]:

$$\begin{aligned} \mathcal{L} [ {}_0^{CF} D_t^{(n+\mu)} f(t) ](s) &= \frac{1}{1 - \mu} \mathcal{L} [ f^{(n+1)}(t) ] \mathcal{L} \left[ \exp\left(-\frac{\mu}{\mu - 1} t\right) \right] \\ &= \frac{s^{n+1} \mathcal{L} [ f(t) ] - s^n f(0) - s^{n-1} f'(0) \dots - f^{(n)}(0)}{s + \mu(1 - s)}. \end{aligned} \tag{3}$$

### 3 General description of the method using the operator of Caputo-Fabrizio type

Consider the following nonlinear partial differential equation in the Caputo-Fabrizio sense:

$${}_{0^+}^{CF} \mathcal{D}_t^{(n+\mu)} u(x, t) + \beta u(x, t) + \varphi u(x, t) = \kappa(x, t), \quad m - 1 < \mu + n \leq m, \tag{4}$$

with initial conditions

$$\frac{\partial^h u(x, 0)}{\partial t^h} = f_h(x), \quad h = 0, 1, \dots, m - 1. \tag{5}$$

Applying the Laplace transform (3) in (4) yields

$$\mathcal{L}[u(x, t)] = \Theta(x, s) - \left( \frac{s + \mu(1 - s)}{s^{n+1}} \right) \mathcal{L}[\beta u(x, t) + \varphi u(x, t)], \tag{6}$$

where

$$\Theta(x, s) = \frac{1}{s^{n+1}} [s^n f_0(x) + s^{n-1} f_1(x) + \dots + f_n(x)] + \frac{s + \mu(1 - s)}{s^{n+1}} \tilde{\kappa}(x, s). \tag{7}$$

Applying the Laplace inverse operator on both sides of (6) yields

$$u(x, t) = \Theta(x, t) - \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1 - s)}{s^{n+1}} \right) \mathcal{L}[\beta u(x, t) + \varphi u(x, t)] \right], \tag{8}$$

where  $\Theta(x, t)$  represents the term arising from the source term. Now, we apply the HPTM to obtain the solution of equation (8) starting by the hypothesis that  $u(x, t)$  is a solution of this equation, which we express as

$$u(x, t) = \sum_{n=0}^{\infty} z^n u_n(x, t), \tag{9}$$

where  $u_n(x, t)$  are known functions, the nonlinear term can be decomposed as

$$\varphi u(x, t) = \sum_{n=0}^{\infty} z^n H_n(x, t), \tag{10}$$

the polynomials  $H_n(x, t)$  are given by [52]

$$H_n(u_0, u_1, \dots, u_n) = \frac{1}{n!} \frac{\partial^n}{\partial z^n} \left[ \varphi \left( \sum_{i=0}^{\infty} z^i u_i \right) \right]_{z=0}, \quad n = 0, 1, 2, \dots; \tag{11}$$

substituting (9) and (10) into (8) we get

$$\sum_{n=0}^{\infty} u_n(x, t) = \Theta(x, t) - z \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1 - s)}{s^{n+1}} \right) \mathcal{L} \left[ \beta \sum_{n=0}^{\infty} z^n u_n(x, t) + \sum_{i=0}^{\infty} z^i H_n \right] \right]. \tag{12}$$

Comparing the coefficients of like powers of  $z$  yields

$$\begin{aligned}
 z^0 : u_0(x, t) &= \Theta(x, t), \\
 z^1 : u_1(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s^{n+1}} \right) \mathcal{L} [\beta u_0(x, t) + H_0(u)] \right], \\
 z^2 : u_2(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s^{n+1}} \right) \mathcal{L} [\beta u_1(x, t) + H_1(u)] \right], \\
 z^3 : u_3(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s^{n+1}} \right) \mathcal{L} [\beta u_2(x, t) + H_2(u)] \right], \\
 &\vdots \\
 z^{n+1} : u_{n+1}(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s^{n+1}} \right) \mathcal{L} [\beta u_n(x, t) + H_n(u)] \right].
 \end{aligned}
 \tag{13}$$

### 4 Applications

We present the solutions obtained by the application of the HPTM with Caputo-Fabrizio fractional operator for some NFPDEs.

#### Example 1

Regarding the following nonlinear KdV equation in the Caputo-Fabrizio sense:

$${}_0^{\text{CF}}\mathcal{D}_t^\mu u(x, t) = -u \frac{\partial u}{\partial x} - u \frac{\partial^3 u}{\partial x^3}, \quad 0 < \mu \leq 1,
 \tag{14}$$

with the initial condition

$$u(x, 0) = x,
 \tag{15}$$

applying the Laplace transform to (14) and considering the condition (15), we have

$$\mathcal{L} [u(x, t)] = \frac{1}{s} u(x, 0) - \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} + u \frac{\partial^3 u}{\partial x^3} \right].
 \tag{16}$$

Applying the inverse of the Laplace transform to (16) yields

$$u(x, t) = u(x, 0) - \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} + u \frac{\partial^3 u}{\partial x^3} \right] \right].
 \tag{17}$$

Now, we apply the HPTM

$$\sum_{n=0}^{\infty} u_n(x, t) = x - z \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ \sum_{n=0}^{\infty} z^n H_n(u) + \sum_{i=0}^{\infty} z^i H_i(u) \right] \right],
 \tag{18}$$


where  $H_n(u)$  are the polynomials that represent the nonlinear terms defined in (11). The polynomials  $H_n(u)$  are calculated in the following form:

$$\begin{aligned}
 H_0(u) &= u_0 \frac{\partial}{\partial x}(u_0) + u_0 \frac{\partial^3}{\partial x^3}(u_0), \\
 H_1(u) &= \frac{\partial}{\partial z} \left[ (u_0 + zu_1) \frac{\partial}{\partial x}(u_0 + zu_1) + (u_0 + zu_1) \frac{\partial^3}{\partial x^3}(u_0 + zu_1) \right]_{z=0} \\
 &= u_0 \frac{\partial}{\partial x}(u_1) + u_1 \frac{\partial}{\partial x}(u_0) + u_0 \frac{\partial^3}{\partial x^3}(u_1) + u_1 \frac{\partial^3}{\partial x^3}(u_0), \\
 H_2(u) &= \frac{1}{2} \frac{\partial^2}{\partial z^2} \left[ (u_0 + zu_1 + z^2u_2) \frac{\partial}{\partial x}(u_0 + zu_1 + z^2u_2) \right]_{z=0} \\
 &\quad + \frac{1}{2} \frac{\partial^2}{\partial z^2} \left[ (u_0 + zu_1 + z^2u_2) \frac{\partial^3}{\partial x^3}(u_0 + zu_1 + z^2u_2) \right]_{z=0} \\
 &= u_0 \frac{\partial}{\partial x}(u_2) + u_1 \frac{\partial}{\partial x}(u_1) + u_2 \frac{\partial}{\partial x}(u_0) \\
 &\quad + u_0 \frac{\partial^3}{\partial x^3}(u_2) + u_1 \frac{\partial^3}{\partial x^3}(u_1) + u_2 \frac{\partial^3}{\partial x^3}(u_0) \\
 &\quad \vdots
 \end{aligned} \tag{19}$$

thus,  $H_0(u) = x$ .

Comparing the coefficients of  $z$  in equation (18), we have

$$\begin{aligned}
 z^0 : u_0(x, t) &= x, \\
 z^1 : u_1(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L}[H_0(u) + K_n(u)] \right] \\
 &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L}[x] \right] \\
 &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \frac{x}{s} \right] \\
 &= -\mathcal{L}^{-1} \left[ \frac{x\mu}{s^2} + \frac{(1-\mu)x}{s} \right] \\
 &= -x\mu t - x(1-\mu), \\
 z^2 : u_2(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L}[-2\mu xt - 2x(1-\mu)] \right] \\
 &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \left[ \frac{-2\mu x}{s^2} - \frac{2x(1-\mu)}{s} \right] \right] \\
 &= -2\mu x \mathcal{L}^{-1} \left[ \frac{s + \mu(1-s)}{s^3} \right] + 2x(1-\mu) \mathcal{L}^{-1} \left[ \frac{s + \mu(1-s)}{s^2} \right] \\
 &= -2\mu x \mathcal{L}^{-1} \left[ \frac{\mu}{s^3} + \frac{(1-\mu)}{s^2} \right] + 2x(1-\mu) \mathcal{L}^{-1} \left[ \frac{\mu}{s^2} + \frac{(1-\mu)}{s} \right] \\
 &= \mu^2 xt^2 + 4\mu(1-\mu)xt + 2(1-\mu)^2x \\
 &\quad \vdots
 \end{aligned} \tag{20}$$



so the approximate solution of  $u(x, t)$  is given by

$$\begin{aligned}
 u(x, t) &= \sum_{n=0}^{\infty} u_n(x, t) \\
 &= x(1 - \mu t - (1 - \mu) + \mu^2 t^2 + 4\mu(1 - \mu)t + 2(1 - \mu^2) + \dots). \tag{21}
 \end{aligned}$$

Therefore the analytical solution when  $\mu \rightarrow 1$ , is given by

$$\begin{aligned}
 u(x, t) &= x(1 - t + t^2 - t^3 + \dots) \\
 &= x \sum_{k=0}^{\infty} (-1)^k t^k \\
 &= \frac{x}{1 + t}. \tag{22}
 \end{aligned}$$

Figure 1 shows the numerical evaluation of (21).

**Example 2** Next, the following nonlinear KdV equation in the Caputo-Fabrizio sense is analyzed:

$${}_0^{\text{CF}}\mathcal{D}_t^\mu u(x, t) = -u \frac{\partial u}{\partial x} - \frac{\partial^3 u}{\partial x^3} + x^2 + 2x^3 t^2, \quad 0 < \mu \leq 1, \tag{23}$$

with the initial condition

$$u(x, 0) = 0. \tag{24}$$

Applying the Laplace transform to (23) and considering the condition (24), we obtain

$$\mathcal{L}[u(x, t)] = \left( \frac{s + \mu(1 - s)}{s} \right) \left[ \frac{x^2}{s} + \frac{4x^3}{s^3} \right] - \left( \frac{s + \mu(1 - s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} \right]. \tag{25}$$

Applying the inverse of the Laplace transform to the above equation yields

$$\begin{aligned}
 u(x, t) = & x^3 \left( \frac{2\mu t^3}{3} - 2\mu t^2 + 2t^2 \right) + x^2(-\mu + \mu t + 1) \\
 & - \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} + \frac{\partial^3 u}{\partial x^3} \right] \right].
 \end{aligned}
 \tag{26}$$

Now, we apply the HPTM to (26)

$$\begin{aligned}
 \sum_{n=0}^{\infty} u_n(x, t) = & x^3 \left( \frac{2\mu t^3}{3} - 2\mu t^2 + 2t^2 \right) + x^2(-\mu + \mu t + 1) \\
 & - z \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ \sum_{n=0}^{\infty} z^n H_n(u) + \left( \sum_{n=0}^{\infty} z^n u_n(x, t) \right)_{xxx} \right] \right],
 \end{aligned}
 \tag{27}$$

where the polynomials  $H_n(u)$  can be expressed in the following form:


$$\begin{aligned}
 H_0(u) &= u_0 \frac{\partial}{\partial x}(u_0), \\
 H_1(u) &= \frac{\partial}{\partial z} \left[ (u_0 + zu_1) \frac{\partial}{\partial x}(u_0 + zu_1) \right]_{z=0} \\
 &= u_0 \frac{\partial}{\partial x}(u_1) + u_1 \frac{\partial}{\partial x}(u_0), \\
 H_2(u) &= \frac{1}{2} \frac{\partial^2}{\partial z^2} \left[ (u_0 + zu_1 + z^2u_2) \frac{\partial}{\partial x}(u_0 + zu_1 + z^2u_2) \right]_{z=0} \\
 &= u_0 \frac{\partial}{\partial x}(u_2) + u_1 \frac{\partial}{\partial x}(u_1) + u_2 \frac{\partial}{\partial x}(u_0) \\
 &\vdots
 \end{aligned}
 \tag{28}$$

Comparing the coefficients of  $z$  in (27), we have

$$\begin{aligned}
 z^0 : u_0(x, t) &= x^3 \left( \frac{2\mu t^3}{3} - 2\mu t^2 + 2t^2 \right) + x^2(-\mu + \mu t + 1), \\
 z^1 : u_1(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ H_0(u) + \frac{\partial^3 u_0}{\partial x^3} \right] \right] \\
 &= -\frac{4\mu^3 t^7 x^5}{21} - 6t(\mu^3 x^3 - 2\mu^2 x^3 + \mu x^3) + 2(\mu^3 x^3 - 3\mu^2 x^3 + 3\mu x^3 - x^3) \\
 &\quad + 2t^2(-6\mu^2 + 12\mu + 5\mu^3 x^4 - 15\mu^2 x^4 + 15\mu x^4 - 5x^4 + 2\mu^3 x^3 - 2\mu^2 x^3 - 6) \\
 &\quad + \frac{8t^6}{3}(\mu^3 x^5 - \mu^2 x^5) \\
 &\quad - \frac{2t^3}{3}(-12\mu^2 + 12\mu + 25\mu^3 x^4 - 50\mu^2 x^4 + 25\mu x^4 + \mu^3 x^3) \\
 &\quad + \frac{t^4}{3}(-3\mu^2 + 36\mu^3 x^5 - 108\mu^2 x^5 + 108\mu x^5 - 36x^5 + 20\mu^3 x^4 - 20\mu^2 x^4) \\
 &\quad - \frac{2t^5}{15}(78\mu^3 x^5 - 156\mu^2 x^5 + 78\mu x^5 + 5\mu^3 x^4),
 \end{aligned}$$

$$\begin{aligned}
 z^2 : u_2(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ H_1(u) + \frac{\partial^3 u_1}{\partial x^3} \right] \right] \\
 &= -2 \left( -6\mu^4 + 24\mu^3 - 36\mu^2 + 24\mu - \frac{1}{2}\mu^4 t^4 + 6(\mu^4 - \mu^3)t^3 \right. \\
 &\quad \left. - 21(\mu^4 - 2\mu^3 + \mu^2)t^2 + 24\mu^4 t - 72\mu^3 t + 72\mu^2 t - 24\mu t - 6 \right) \\
 &\quad - 2x \left( -\frac{3}{2}\mu^4 t^6 + \frac{134}{5}(\mu^4 - \mu^3)t^5 - 144\mu^4 t^4 + 288\mu^3 t^4 - 144\mu^2 t^4 \right. \\
 &\quad \left. + 264(\mu^4 - 3\mu^3 + 3\mu^2 - \mu)t^3 - 132(\mu^4 - 4\mu^3 + 6\mu^2 - 4\mu + 1)t^2 \right) \\
 &\quad - 2x^2 \left( -\frac{47}{56}\mu^4 t^8 + \frac{138}{7}(\mu^4 - \mu^3)t^7 - 149\mu^4 t^6 + 298\mu^3 t^6 - 149\mu^2 t^6 \right. \\
 &\quad \left. + \frac{2,136}{5}(\mu^4 - 3\mu^3 + 3\mu^2 - \mu)t^5 \right. \\
 &\quad \left. - 396(\mu^4 t^4 - 4\mu^3 t^4 + 6\mu^2 t^4 - 4\mu t^4 + t^4) \right) \tag{29} \\
 &\quad - 2x^4 \left( 5\mu^5 - 25\mu^4 + 50\mu^3 - 50\mu^2 + 25\mu - \frac{1}{3}\mu^5 t^5 \right. \\
 &\quad \left. + \frac{55}{12}(\mu^5 t^4 - \mu^4 t^4) - 20(\mu^5 - 2\mu^4 + \mu^3)t^3 \right. \\
 &\quad \left. + 35(\mu^5 - 3\mu^4 + 3\mu^3 - \mu^2)t^2 - 25\mu^5 t + 100\mu^4 t \right. \\
 &\quad \left. - 150\mu^3 t + 100\mu^2 t - 25\mu t - 5 \right) \\
 &\quad - 2x^5 \left( -\frac{10}{21}\mu^5 t^7 + 9\mu^5 t^6 - 9\mu^4 t^6 - \frac{276}{5}(\mu^5 - 2\mu^4 + \mu^3)t^5 \right. \\
 &\quad \left. + 136\mu^5 t^4 - 408\mu^4 t^4 + 408\mu^3 t^4 \right. \\
 &\quad \left. - 136\mu^2 t^4 - 134(\mu^5 - 4\mu^4 + 6\mu^3 - 4\mu^2 + \mu)t^3 \right. \\
 &\quad \left. + 42(\mu^5 - 5\mu^4 + 10\mu^3 - 10\mu^2 + 5\mu - 1)t^2 \right) \\
 &\quad - 2x^6 \left( -\frac{20}{81}\mu^5 t^9 + 6\mu^5 t^8 - 6\mu^4 t^8 - \frac{2,204}{45}(\mu^5 - 2\mu^4 + \mu^3)t^7 \right. \\
 &\quad \left. + \frac{7,546}{45}(\mu^5 t^6 - 3\mu^4 t^6 + 3\mu^3 t^6 - \mu^2 t^6) \right. \\
 &\quad \left. - \frac{1,204}{5}(\mu^5 - 4\mu^4 + 6\mu^3 - 4\mu^2 + \mu)t^5 + 112\mu^5 t^4 - 560\mu^4 t^4 \right. \\
 &\quad \left. + 1,120\mu^3 t^4 - 1,120\mu^2 t^4 + 560\mu t^4 - 112t^4 \right) \\
 &\quad - 2x^7 \left( -\frac{32}{693}\mu^5 t^{11} + \frac{48}{35}(\mu^5 t^{10} - \mu^4 t^{10}) - \frac{13,312}{945}(\mu^5 t^9 - 2\mu^4 t^9 + \mu^3 t^9) \right. \\
 &\quad \left. + \frac{952}{15}(\mu^5 t^8 - 3\mu^4 t^8 + 3\mu^3 t^8 - \mu^2 t^8) \right. \\
 &\quad \left. - \frac{4,512}{35}(\mu^5 - 4\mu^4 + 6\mu^3 - 4\mu^2 + \mu)t^7 + 96\mu^5 t^6 - 480\mu^4 t^6 \right)
 \end{aligned}$$





$$\begin{aligned}
 &+ 960\mu^3 t^6 - 960\mu^2 t^6 + 480\mu t^6 - 96t^6) \\
 &\vdots
 \end{aligned}$$

so the approximate solution of  $u(x, t)$  is given by

$$u(x, t) = \sum_{n=0}^{\infty} u_n(x, t) = u_0 + u_1 + u_2 + \dots \tag{30}$$

Therefore the analytical solution when  $\mu \rightarrow 1$  is given by

$$u(x, t) = x^2 t. \tag{31}$$

Figure 2 shows the numerical evaluation of (30).

**Example 3** We consider the following nonlinear Burger equation in the Caputo-Fabrizio sense:

$${}_0^{\text{CF}}\mathcal{D}_t^\mu u(x, t) + u \frac{\partial u}{\partial x} = \eta \frac{\partial^2 u}{\partial x^2}, \quad 0 < \mu \leq 1, \tag{32}$$

subject to the initial condition

$$u(x, 0) = nx, \quad n \in \mathbb{N}. \tag{33}$$

Applying the Laplace transform to (32) and considering the condition (33), we have

$$\mathcal{L}[u(x, t)] = \frac{1}{s} u(x, 0) - \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} - \eta \frac{\partial^2 u}{\partial x^2} \right]. \tag{34}$$

Applying the inverse of the Laplace transform to the above equation yields

$$u(x, t) = u(x, 0) - \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ u \frac{\partial u}{\partial x} - \eta \frac{\partial^2 u}{\partial x^2} \right] \right]. \tag{35}$$

Now, we apply the HPTM to (35)

$$\sum_{n=0}^{\infty} u_n(x, t) = nx - z\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L} \left[ \sum_{n=0}^{\infty} z^n H_n(u) - \eta \frac{\partial^2 u}{\partial x^2} \sum_{i=0}^{\infty} z^i u_n(x, t) \right] \right], \tag{36}$$

where  $H_n(u)$  are the polynomials defined in equation (11), which represent the nonlinear terms. The polynomials  $H_n(u)$  are calculated in the following form:

$$\begin{aligned} H_0(u) &= u_0 \frac{\partial}{\partial x}(u_0), \\ H_1(u) &= \frac{\partial}{\partial z} \left[ (u_0 + zu_1) \frac{\partial}{\partial x}(u_0 + zu_1) \right] \\ &= u_0 \frac{\partial}{\partial x}(u_1) + u_1 \frac{\partial}{\partial x}(u_0), \\ H_2(u) &= \frac{1}{2} \frac{\partial^2}{\partial z^2} \left[ (u_0 + zu_1 + z^2u_2) \frac{\partial}{\partial x}(u_0 + zu_1 + z^2u_2) \right] \\ &= u_0 \frac{\partial}{\partial x}(u_2) + u_1 \frac{\partial}{\partial x}(u_1) + u_2 \frac{\partial}{\partial x}(u_0) \\ &\vdots \end{aligned} \tag{37}$$

thus,  $H_0(u) = n^2x$  and  $u_0(x, t) = 0$ .

Comparing the coefficients of  $z$  in (36), we have


$$\begin{aligned} z^0 : u_0(x, t) &= nx, \\ z^1 : u_1(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L}(nx) \right] \\ &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \left( \frac{nx}{s} \right) \right] \\ &= -n^2x(1 - \mu + \mu t), \end{aligned} \tag{38}$$

$$\begin{aligned} z^2 : u_2(x, t) &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \mathcal{L}[-2n^3x(-\mu + \mu t + 1)] \right] \\ &= -\mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s} \right) \left( -2n^3x \left( \frac{\mu}{s^2} - \frac{\mu}{s} + \frac{1}{s} \right) \right) \right] \\ &= 2n^3x \left( \mu^2 - 2\mu + \frac{\mu^2 t^2}{2} - 2(\mu^2 - \mu)t + 1 \right) \end{aligned} \tag{39}$$

$\vdots$

the approximate solution of  $u(x, t)$  is given by

$$\begin{aligned} u(x, t) &= \sum_{n=0}^{\infty} u_n(x, t) \\ &= u_0 + u_1 + u_2 + \dots \end{aligned} \tag{40}$$



Therefore the analytical solution when  $\mu \rightarrow 1$  is given by

$$\begin{aligned}
 u(x, t) &= nx[1 - nt + n^2t^2 - n^3t^3 + \dots] \\
 &= x \sum_{k=0}^{\infty} (-1)^k n^{k+1} t^k \\
 &= \frac{nx}{1 + nt}.
 \end{aligned}
 \tag{41}$$

Figure 3 shows the numerical evaluation of (40).

**Example 4** Consider the following nonlinear Klein-Gordon equation in the Caputo-Fabrizio sense:

$${}_0^{\text{CF}}\mathcal{D}_t^{\mu+1} u(x, t) = \frac{\partial^2 u}{\partial x^2} - u^2 + x^2 t^2, \quad 0 < \mu \leq 1,
 \tag{42}$$

subject to the initial condition

$$\begin{aligned}
 u(x, 0) &= 0, \\
 u_t(x, 0) &= x.
 \end{aligned}
 \tag{43}$$

Applying the Laplace transform to (42) and considering the condition (43), we have

$$\mathcal{L}[u(x, t)] = \frac{1}{s^2} u_t(x, 0) + 2 \left( \frac{s + \mu(1-s)}{s^5} \right) x^2 + \left( \frac{s + \mu(1-s)}{s^2} \right) \mathcal{L} \left[ \frac{\partial^2 u}{\partial x^2} - u^2 \right].
 \tag{44}$$

Applying the inverse of the Laplace transform to the above equation yields

$$u(x, t) = xt + \frac{\mu t^4 x^2}{12} + \frac{(1-\mu)t^3 x^2}{3} + \mathcal{L}^{-1} \left[ \left( \frac{s + \mu(1-s)}{s^2} \right) \mathcal{L} \left[ \frac{\partial^2 u}{\partial x^2} - u^2 \right] \right].
 \tag{45}$$

Now, we apply the HPTM to (45)

$$\sum_{n=0}^{\infty} u_n(x, t) = xt + \frac{\mu t^4 x^2}{12} + \frac{(1-\mu)t^3 x^2}{3} + z\mathcal{L}^{-1}\left[\left(\frac{s + \mu(1-s)}{s^2}\right)\mathcal{L}\left[\frac{\partial^2 u}{\partial x^2} \sum_{n=0}^{\infty} z^n u_n(x, t) - \sum_{i=0}^{\infty} z^i H_i(u)\right]\right], \tag{46}$$

where  $H_n(u)$  are the polynomials defined in equation (11), that represent the nonlinear terms. The polynomial  $H_n(u)$  are calculated in the following form:

$$\begin{aligned} H_0(u) &= (u_0)^2, \\ H_1(u) &= \frac{\partial}{\partial z} [(u_0 + zu_1)^2] \\ &= 2u_0u_1 \\ &\vdots \end{aligned} \tag{47}$$

thus,  $H_0(u) = (xt + \frac{\mu t^4 x^2}{12} + \frac{(1-\mu)t^3 x^2}{3})^2$ .

Comparing the coefficients of  $z$  in (46), we have

$$z^0 : u_0(x, t) = xt + \frac{\mu t^4 x^2}{12} + \frac{(1-\mu)t^3 x^2}{3},$$

$$z^1 : u_1(x, t)$$

$$\begin{aligned} &= \mathcal{L}^{-1}\left[\left(\frac{s + \mu(1-s)}{s^2}\right)\mathcal{L}\left(-\frac{2x^2}{s^3} + \frac{4(1-\mu)}{s^4} - \frac{16x^3(1-\mu)}{s^5} - \frac{80x^4(1-\mu)^2}{s^7} + \frac{4\mu}{s^5} - \frac{20x^3\mu}{s^6} - \frac{280x^4(1-\mu)\mu}{s^8} - \frac{280x^4\mu^2}{s^9}\right)\right] \\ &= -\frac{t^4 x^2 \mu}{12} + \frac{t^5(1-\mu)\mu}{30} - \frac{t^6 x^3(1-\mu)\mu}{45} - \frac{t^8 x^4(1-\mu)^2 \mu}{504} \\ &\quad + \frac{t^6 \mu^2}{180} - \frac{x^3 \mu^2 t^7}{252} - \frac{t^9 x^4(1-\mu)\mu^2}{1,296} - \frac{t^{10} x^4 \mu^2}{12,960} \\ &\quad - \frac{t^3 x^2(1-\mu)}{3} + \frac{t^4(1-\mu)^2}{6} - \frac{2t^5 x^3(1-\mu)^2}{15} - \frac{t^7 x^4(1-\mu)^3}{63} \\ &\quad + \frac{t^5 \mu(1-\mu)x^4}{30} - \frac{t^6 x^3 \mu(1-\mu)}{36} - \frac{t^8 x^4(1-\mu)^2 \mu}{144} - \frac{t^9 x^4(1-\mu)\mu^2}{1,296}, \end{aligned}$$

$$z^2 : u_2(x, t)$$

$$\begin{aligned} &= -\mathcal{L}^{-1}\left[\left(\frac{s + \mu(1-s)}{s^2}\right)\mathcal{L}[(u_1)_{xx} - H_1(u)]\right] \\ &= \mu\left[\frac{\mu^4 t^{16} x^6}{18,662,400} - \frac{t^6}{180}(-\mu^2 + \mu + 4\mu x^3 - 4x^3) - \frac{t^5}{30}(\mu^2 - 2\mu + 1) + \frac{t^8(44\mu^2 x^4 - 88\mu x^4 + 44x^4 - 27\mu^3 x + 66\mu^2 x - 39\mu x)}{5,040} + \frac{t^7(5\mu x^3 + 24\mu^3 x - 82\mu^2 x + 92\mu x - 34x)}{1,260}\right] \end{aligned} \tag{48}$$




Figure 4 Numerical evaluation of (49).

$$\begin{aligned}
 & -t^{10}(304\mu^3x^5 - 912\mu^2x^5 + 912\mu x^5 - 304x^5 - 55\mu^2x^4 - 270\mu^4x^2 \\
 & + 992\mu^3x^2 - 1,174\mu^2x^2 + 452\mu x^2)/226,800 \\
 & + \frac{t^{11}(555\mu^3x^5 - 1,110\mu^2x^5 + 555\mu x^5 - 140\mu^4x^2 + 392\mu^3x^2 - 252\mu^2x^2)}{831,600} \\
 & -t^9(133\mu^2x^4 - 133\mu x^4 + 120\mu^4x^2 - 550\mu^3x^2 \\
 & + 930\mu^2x^2 - 690\mu x^2 + 190x^2 - 15\mu^3x + 22\mu^2x)/45,360 \\
 & + \frac{137t^{14}(\mu^4x^6 - 2\mu^3x^6 + \mu^2x^6)}{9,906,624} - \frac{t^{15}(\mu^4x^6 - \mu^3x^6)}{680,400} \\
 & + t^{12}(240\mu^4x^6 - 960\mu^3x^6 + 1,440\mu^2x^6 - 960\mu x^6 + 240x^6 \\
 & - 319\mu^3x^5 + 319\mu^2x^5 + 21\mu^4x^2 - 42\mu^3x^2)/2,993,760 \\
 & - \frac{t^{13}(390\mu^4x^6 - 1,170\mu^3x^6 + 1,170\mu^2x^6 - 390\mu x^6 - 37\mu^3x^5)}{7,076,160} \Big] \\
 & \vdots
 \end{aligned}$$

and the approximate solution of  $u(x, t)$  is

$$u(x, t) = \sum_{n=0}^{\infty} u_n(x, t) = u_0 + u_1 + u_2 + u_3 + \dots \tag{49}$$

Therefore the analytical solution when  $\mu \rightarrow 1$  is given by

$$u(x, t) = xt. \tag{50}$$

Figure 4 shows the numerical evaluation of (49).

**Example 5** Consider the following nonlinear Klein-Gordon equation in the Caputo-Fabrizio sense:

$${}^{\text{CF}}\mathcal{D}_t^{\mu+1}u(x, t) = \frac{\partial^2 u}{\partial x^2} - u^2 + t \sin(x) + t^2 \sin^2(x), \quad 0 < \mu \leq 1, \tag{51}$$

subject to the initial condition

$$\begin{aligned} u(x, 0) &= 0, \\ u_t(x, 0) &= \sin(x). \end{aligned} \tag{52}$$

Applying the Laplace transform to (51) and considering the condition (52), we have

$$\begin{aligned} \mathcal{L}[u(x, t)] &= \frac{\sin(x)}{s^2} + \left(\frac{s + \mu(1-s)}{s^2}\right) \left[\frac{\sin(x)}{s^2} + \frac{2 \sin^2(x)}{s^3}\right] \\ &\quad + \left(\frac{s + \mu(1-s)}{s^2}\right) \mathcal{L}\left[\frac{\partial^2 u}{\partial x^2} - u^2\right]. \end{aligned} \tag{53}$$

Applying the inverse of the Laplace transform to the above equation yields

$$\begin{aligned} u(x, t) &= \sin(x)t + \frac{\mu t^4 \sin^2(x)}{12} + \frac{(1-\mu)t^3 \sin^2(x)}{3} + \frac{\mu t^3 \sin(x)}{6} \\ &\quad + \frac{(1-\mu)t^2 \sin(x)}{2} + \mathcal{L}^{-1}\left[\left(\frac{s + \mu(1-s)}{s^2}\right) \mathcal{L}\left[\frac{\partial^2 u}{\partial x^2} - u^2\right]\right]. \end{aligned} \tag{54}$$

Now, we apply the HPTM to (54),

$$\begin{aligned} \sum_{n=0}^{\infty} u_n(x, t) &= \sin(x)t + \frac{\mu t^4 \sin^2(x)}{12} + \frac{(1-\mu)t^3 \sin^2(x)}{3} + \frac{\mu t^3 \sin(x)}{6} + \frac{(1-\mu)t^2 \sin(x)}{2} \\ &\quad + z \mathcal{L}^{-1}\left[\left(\frac{s + \mu(1-s)}{s^2}\right) \mathcal{L}\left[\sum_{n=0}^{\infty} z^n u_n(x, t) - \sum_{n=0}^{\infty} z^n H_n(x, t)\right]\right], \end{aligned} \tag{55}$$

where  $H_n(u)$  are the polynomials defined in (11), which represent the nonlinear terms. The polynomial  $H_n(u)$  are calculated in the following form:

$$\begin{aligned} H_0(u) &= (u_0)^2, \\ H_1(u) &= \frac{\partial}{\partial z} [(u_0 + zu_1)^2] \\ &= 2u_0u_1 \\ &\vdots \end{aligned} \tag{56}$$

Comparing the coefficients of the same order in powers of  $p$  in (55), we have

$$\begin{aligned} z^0 : u_0(x, t) &= \sin(x)t + \frac{\mu t^4 \sin^2(x)}{12} + \frac{(1-\mu)t^3 \sin^2(x)}{3} + \frac{\mu t^3 \sin(x)}{6} + \frac{(1-\mu)t^2 \sin(x)}{2}, \\ z^1 : u_1(x, t) &= \mathcal{L}^{-1}\left[\left[\frac{s + \mu(1-s)}{s^2}\right] \mathcal{L}\left[\frac{\partial^2 u_0}{\partial x^2} - H_0(u)\right]\right] \end{aligned}$$

$$\begin{aligned}
 &= -\frac{\mu^2 t^5 \cos^2(x)}{15} + \frac{\mu t^5 \cos^2(x)}{15} + \frac{\mu^2 t^4 \cos^2(x)}{6} - \frac{\mu t^4 \cos^2(x)}{3} + \frac{t^4 \cos^2(x)}{6} \\
 &+ \frac{\mu^2 t^6 \cos^2(x)}{180} + \frac{\mu t^3 \sin(x)}{6} - \frac{t^3 \sin(x)}{6} + \frac{\mu t^2 \sin(x)}{2} - \frac{t^2 \sin(x)}{2} \\
 &- \frac{\mu^2 t^5 \sin(x)}{120} + \frac{\mu^2 t^4 \sin(x)}{12} - \frac{\mu t^4 \sin(x)}{12} - \frac{\mu^2 t^3 \sin(x)}{6} \\
 &- \frac{t^5 \sin^2(x)}{20} + \frac{3\mu t^4 \sin^2(x)}{4} - \frac{5t^4 \sin^2(x)}{12} + \frac{\mu t^3 \sin^2(x)}{3} - \frac{t^3 \sin^2(x)}{3} \\
 &- \frac{13\mu t^6 \sin^2(x)}{360} + \frac{\mu^2 t^5 \sin^2(x)}{30} - \frac{\mu t^5 \sin^2(x)}{30} - \frac{5\mu^2 t^4 \sin^2(x)}{12} \\
 &- \frac{\mu^2 t^7 \sin^2(x)}{126} - \frac{13\mu^3 t^6 \sin^2(x)}{360} + \frac{\mu^2 t^6 \sin^2(x)}{18} + \frac{\mu^3 t^5 \sin^2(x)}{20} \tag{57} \\
 &- \frac{\mu^3 t^8 \sin^2(x)}{2,016} + \frac{\mu^3 t^7 \sin^2(x)}{126} - \frac{t^6 \sin^3(x)}{18} + \frac{4\mu t^5 \sin^3(x)}{15} - \frac{2t^5 \sin^3(x)}{15} \\
 &- \frac{\mu t^7 \sin^3(x)}{28} - \frac{7\mu^2 t^6 \sin^3(x)}{60} + \frac{7\mu t^6 \sin^3(x)}{60} - \frac{2\mu^2 t^5 \sin^3(x)}{15} \\
 &- \frac{\mu^2 t^8 \sin^3(x)}{144} - \frac{\mu^3 t^7 \sin^3(x)}{28} + \frac{17\mu^2 t^7 \sin^3(x)}{252} + \frac{\mu^3 t^6 \sin^3(x)}{18} \\
 &- \frac{\mu^3 t^9 \sin^3(x)}{2,592} + \frac{\mu^3 t^8 \sin^3(x)}{144} + \frac{\mu t^7 \sin^4(x)}{21} - \frac{t^7 \sin^4(x)}{63} \\
 &- \frac{\mu^2 t^9 \sin^4(x)}{648} + \frac{\mu^2 t^8 \sin^4(x)}{56} - \frac{\mu t^8 \sin^4(x)}{112} - \frac{\mu^2 t^7 \sin^4(x)}{21} \\
 &- \frac{\mu^3 t^{10} \sin^4(x)}{12,960} + \frac{\mu^3 t^9 \sin^4(x)}{648} - \frac{\mu^3 t^8 \sin^4(x)}{112} + \frac{\mu^3 t^7 \sin^4(x)}{63} \\
 &\vdots
 \end{aligned}$$

and the approximate solution of  $u(x, t)$  is given by

$$\begin{aligned}
 u(x, t) &= \sum_{n=0}^{\infty} u_n(x, t) \\
 &= u_0 + u_1 + u_2 + \dots \tag{58}
 \end{aligned}$$


Therefore the analytical solution when  $\mu \rightarrow 1$  is

$$u(x, t) = \sin(x)t. \tag{59}$$

Figure 5 shows the numerical evaluation of (58).

#### 4.1 Convergence and stability analysis

If the series (12) converges ( $n = 0, 1, 2, \dots, n$ ), where  $\Theta(x, s)$  is governed by (7), it must be the solution of equation (4). Overall, the results show that the proposed approach is unconditionally stable and convergent. The method provides a simple way to control the convergence region of the solution by introducing (11) and our approximate results agree well with exact solutions and numerical ones.



Madani in [53] have compared the approximate solutions obtained by means of HPTM in a wide range of the problem's domain with those results obtained from the exact analytical solutions and the HAM. This comparison shows precise agreement between the HPTM and exact results. The HPTM solution is valid for a wide range of time and this suggests that the HPTM method can solve non-homogeneous equations with a high degree of accuracy by considering only few terms in the perturbed solution. On the other hand the relative error for the HAM is dramatically increased as the time value  $t$  increases, so the HAM solution validity range is restricted to a short region.

Therefore the HPTM method is a powerful new method which needs less computation time and is much easier and more convenient than the HAM, because the Laplace transform allows one in many situations to overcome the deficiency mainly caused by unsatisfied boundary or initial conditions that appear in other semi-analytical methods such as HAM [53].

## 5 Conclusions

In this paper, the HPTM method was developed to solve fractional nonlinear differential equations using the Caputo-Fabrizio operator. With the polynomials expansion considered in the HTPM method we obtained an infinite series solution for the fractional partial differential equations. Based on the HPTM, a general scheme was developed to obtain approximate solutions of fractional equations and the solutions are given in a series form, which converges rapidly. The methodology presented has become an important mathematical tool, motivated by the potential use for physicists and engineers working in various areas of the natural sciences.

This work shows that the HPTM method is an efficient tool for solving nonlinear fractional partial differential equations considering the fractional operator of Caputo-Fabrizio type. The HPTM yields a rapidly convergent series solution by using a few iterations [53, 54]. In this paper a Mathematica program has been used for computations and programming.

### Competing interests

The authors declare that there is no conflict of interests regarding the publication of this paper.



**Authors' contributions**

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

**Author details**

<sup>1</sup>CONACyT-Centro Nacional de Investigación y Desarrollo Tecnológico, Tecnológico Nacional de México, Interior Internado Palmira S/N, Col. Palmira, Cuernavaca, Morelos 62490, México. <sup>2</sup>Universidad Autónoma de la Ciudad de México, Prolongación San Isidro 151, Col. San Lorenzo Tezonco, Del. Iztapalapa, México D.F. 09790, México. <sup>3</sup>Departamento de Electromecánica, Instituto Tecnológico Superior de Irapuato, Irapuato, Guanajuato, México. <sup>4</sup>Departamento de Ingeniería Física, División de Ciencias e Ingenierías Campus León, Universidad de Guanajuato, León, Guanajuato, México. <sup>5</sup>Centro Nacional de Investigación y Desarrollo Tecnológico, Tecnológico Nacional de México, Interior Internado Palmira S/N, Col. Palmira, Cuernavaca, Morelos 62490, México.

**Acknowledgements**

We would like to thank Mayra Martínez for the interesting discussions. José Francisco Gómez Aguilar acknowledges the support provided by CONACyT: Cátedras CONACyT para jóvenes investigadores 2014.

Received: 24 November 2016 Accepted: 17 February 2017 Published online: 28 February 2017

**References**

- Cattani, C, Srivastava, HM, Yang, XJ: Fractional Dynamics. de Gruyter, Berlin (2016)
- Yao, JJ, Kumar, A, Kumar, S: A fractional model to describe the Brownian motion of particles and its analytical solution. *Adv. Mech. Eng.* **7**(12), 1687814015618874 (2015)
- Gómez-Aguilar, JF, Miranda-Hernández, M, López-López, MG, Alvarado-Martínez, VM, Baleanu, D: Modeling and simulation of the fractional space-time diffusion equation. *Commun. Nonlinear Sci. Numer. Simul.* **30**(1), 115-127 (2016)
- Kumar, D, Singh, J, Baleanu, D: A hybrid computational approach for Klein-Gordon equations on Cantor sets. *Nonlinear Dyn.* **87**(1), 511-517 (2017)
- Kumar, D, Singh, J, Baleanu, D: Numerical computation of a fractional model of differential-difference equation. *J. Comput. Nonlinear Dyn.* **11**(6), 061004 (2016)
- Singh, J, Kumar, D, Nieto, JJ: A reliable algorithm for a local fractional tricomi equation arising in fractal transonic flow. *Entropy* **18**(6), 206 (2016)
- Kumar, D, Singh, J, Kumar, S, Singh, BP: Numerical computation of nonlinear shock wave equation of fractional order. *Ain Shams Eng. J.* **6**(2), 605-611 (2015)
- Kumar, D, Singh, J, Kılıçman, A: An efficient approach for fractional Harry Dym equation by using Sumudu transform. *Abstr. Appl. Anal.* **2013**, Article ID 608943 (2013)
- Yang, XJ, Tenreiro Machado, JA, Baleanu, D, Cattani, C: On exact traveling-wave solutions for local fractional Korteweg-de Vries equation. *Chaos, Interdiscip. J. Nonlinear Sci.* **26**(8), 084312 (2016)
- Yang, XJ, Baleanu, D, Khan, Y, Mohyud-Din, ST: Local fractional variational iteration method for diffusion and wave equations on Cantor sets. *Rom. J. Phys.* **59**(1-2), 36-48 (2014)
- Baleanu, D, Srivastava, HM, Yang, XJ: Local fractional variational iteration algorithms for the parabolic Fokker-Planck equation defined on Cantor sets. *Prog. Fract. Differ. Appl.* **1**(1), 1-11 (2015)
- Yang, XJ: Fractional derivatives of constant and variable orders applied to anomalous relaxation models in heat-transfer problems. (2016) arXiv:1612.03202
- Xiao-Jun, XJ, Srivastava, HM, Machado, JT: A new fractional derivative without singular kernel. *Therm. Sci.* **20**(2), 753-756 (2016)
- Hristov, J: Transient heat diffusion with a non-singular fading memory: from the Cattaneo constitutive equation with Jeffrey's kernel to the Caputo-Fabrizio time-fractional derivative. *Therm. Sci.* **20**(2), 757-762 (2016)
- Spasic, DT, Kovincic, NI, Dankuc, DV: A new material identification pattern for the fractional Kelvin-Zener model describing biomaterials and human tissues. *Commun. Nonlinear Sci. Numer. Simul.* **37** 193-199 (2016)
- Ma, HC, Yao, DD, Peng, XF: Exact solutions of non-linear fractional partial differential equations by fractional sub-equation method. *Therm. Sci.* **19**(4), 1239-1244 (2015)
- Guo, S, Mei, L: Exact solutions of space-time fractional variant Boussinesq equations. *Adv. Sci. Lett.* **10**(1), 700-702 (2012)
- Mohyud-Din, ST, Nawaz, T, Azhar, E, Akbar, MA: Fractional sub-equation method to space-time fractional Calogero-Degasperis and potential Kadomtsev-Petviashvili equations. *Journal of Taibah University for Science* (2015)
- Jafari, H, Ghorbani, M, Ghasempour, S: A note on exact solutions for nonlinear integral equations by a modified homotopy perturbation method. *New Trends Math. Sci.* **1**(2), 22-26 (2013)
- Nino, UF, Leal, HV, Khan, Y, Díaz, DP, Sesma, AP, Fernández, VJ, Orea, JS: Modified nonlinearities distribution homotopy perturbation method as a tool to find power series solutions to ordinary differential equations. *Nova Scientia* **12**, 13-38 (2014)
- Sayevand, K, Jafari, H: On systems of nonlinear equations: some modified iteration formulas by the homotopy perturbation method with accelerated fourth-and fifth-order convergence. *Appl. Math. Model.* **40**(2), 1467-1476 (2016)
- Das, N, Singh, R, Wazwaz, AM, Kumar, J: An algorithm based on the variational iteration technique for the Bratu-type and the Lane-Emden problems. *J. Math. Chem.* **54**(2), 527-551 (2016)
- Mistry, PR, Pradhan, VH: Approximate analytical solution of non-linear equation in one dimensional instability phenomenon in homogeneous porous media in horizontal direction by variational iteration method. *Proc. Eng.* **127**, 970-977 (2015)
- Wu, GC, Baleanu, D, Deng, ZG: Variational iteration method as a kernel constructive technique. *Appl. Math. Model.* **39**(15), 4378-4384 (2015)
- Wu, GC, Baleanu, D: Variational iteration method for the burgers' flow with fractional derivatives-new Lagrange multipliers. *Appl. Math. Model.* **37**(9), 6183-6190 (2013)

26. Wu, GC, Baleanu, D: New applications of the variational iteration method-from differential equations to q-fractional difference equations. *Adv. Differ. Equ.* **2013**(1), 21 (2013)
27. Duan, JS, Rach, R, Wazwaz, AM: Higher order numeric solutions of the Lane-Emden-type equations derived from the multi-stage modified Adomian decomposition method. *Int. J. Comput. Math.* **94** 197-215 (2015)
28. Kumar, D, Singh, J, Kumar, S: Analytic and approximate solutions of space-time fractional telegraph equations via Laplace transform. *Walailak J. Sci. Technol.* **11**(8), 711-728 (2013)
29. Kumar, D, Singh, J, Kumar, S: Numerical computation of nonlinear fractional Zakharov-Kuznetsov equation arising in ion-acoustic waves. *J. Egypt. Math. Soc.* **22**(3), 373-378 (2014)
30. Duan, JS, Rach, R, Wazwaz, AM: Steady-state concentrations of carbon dioxide absorbed into phenyl glycidyl ether solutions by the Adomian decomposition method. *J. Math. Chem.* **53**(4), 1054-1067 (2015)
31. Tsai, PY, Chen, COK: Free vibration of the nonlinear pendulum using hybrid Laplace Adomian decomposition method. *Int. J. Numer. Methods Biomed. Eng.* **27**(2), 262-272 (2011)
32. Tsai, PY: An approximate analytic solution of the nonlinear Riccati differential equation. *J. Franklin Inst.* **347**(10), 1850-1862 (2010)
33. Lu, L, Duan, J, Fan, L: Solution of the magnetohydrodynamics Jeffery-Hamel flow equations by the modified Adomian decomposition method. *Adv. Appl. Math. Mech.* **7**(5), 675-686 (2015)
34. Yousefi, SA, Dehghan, M, Lotfi, A: Generalized Euler-Lagrange equations for fractional variational problems with free boundary conditions. *Comput. Math. Appl.* **62**(3), 987-995 (2011)
35. Arqub, OA, El-Ajou, A: Solution of the fractional epidemic model by homotopy analysis method. *J. King Saud Univ., Sci.* **25**(1), 73-81 (2013)
36. Vong, S, Wang, Z: A compact difference scheme for a two dimensional fractional Klein-Gordon equation with Neumann boundary conditions. *J. Comput. Phys.* **274**, 268-282 (2014)
37. Atangana, A: On the new fractional derivative and application to nonlinear Fisher's reaction-diffusion equation. *Appl. Math. Comput.* **273**, 948-956 (2016)
38. Mohebbi, A, Abbaszadeh, M, Dehghan, M: High-order difference scheme for the solution of linear time fractional Klein-Gordon equations. *Numer. Methods Partial Differ. Equ.* **30**(4), 1234-1253 (2014)
39. Atangana, A, Secer, A: A note on fractional order derivatives and table of fractional derivatives of some special functions. *Abstr. Appl. Anal.* **2013**, Article ID 279681 (2013)
40. Caputo, M, Fabrizio, M: A new definition of fractional derivative without singular kernel. *Prog. Fract. Differ. Appl.* **1**(2), 73-85 (2015)
41. Lozada, J, Nieto, JJ: Properties of a new fractional derivative without singular kernel. *Prog. Fract. Differ. Appl.* **1**(2), 87-92 (2015)
42. Gómez-Aguilar, JF, Córdova-Fraga, T, Escalante-Martínez, JE, Calderón-Ramón, C, Escobar-Jiménez, RF: Electrical circuits described by a fractional derivative with regular kernel. *Rev. Mex. Fis.* **62**, 144-154 (2016)
43. Gómez-Aguilar, JF, Yépez-Martínez, H, Calderón-Ramón, C, Cruz-Orduña, I, Escobar-Jiménez, RF, Olivares-Peregrino, VH: Modeling of a mass-spring-damper system by fractional derivatives with and without a singular kernel. *Entropy* **17**(9), 6289-6303 (2015)
44. Atangana, A, Alkahtani, BST: Extension of the resistance, inductance, capacitance electrical circuit to fractional derivative without singular kernel. *Adv. Mech. Eng.* **7**(6), 1-6 (2015)
45. Liao, S: *Homotopy Analysis Method in Nonlinear Differential Equations*. Springer, Berlin (2012)
46. Yin, XB, Kumar, S, Kumar, D: A modified homotopy analysis method for solution of fractional wave equations. *Adv. Mech. Eng.* **7**(12), 1687814015620330 (2015)
47. Kumar, S: A new analytical modelling for fractional telegraph equation via Laplace transform. *Appl. Math. Model.* **38**(13), 3154-3163 (2014)
48. Kumar, S, Rashidi, MM: New analytical method for gas dynamics equation arising in shock fronts. *Comput. Phys. Commun.* **185**(7), 1947-1954 (2014)
49. Mohamed, MS, Gepreel, KA, Al-Malki, FA, Al-Humyani, M: Approximate solutions of the generalized Abel's integral equations using the extension Khan's homotopy analysis transformation method. *J. Appl. Math.* **2015**, Article ID 357861 (2015)
50. Gupta, VG, Kumar, P: Approximate solutions of fractional linear and nonlinear differential equations using Laplace homotopy analysis method. *Int. J. Nonlinear Sci.* **19**(2), 113-120 (2015)
51. Khan, Y, Wu, Q: Homotopy perturbation transform method for nonlinear equations using He's polynomials. *Comput. Math. Appl.* **61**(8), 1963-1967 (2011)
52. Ghorbani, A: Beyond Adomian's polynomials: He's polynomials. *Chaos Solitons Fractals* **39**, 1486-1492 (2009)
53. Madani, M, Fathizadeh, M, Khan, Y, Yildirim, A: On the coupling of the homotopy perturbation method and Laplace transformation. *Math. Comput. Model.* **53**(9-10), 1937-1945 (2011)
54. Liao, S: *Beyond Perturbation: Introduction to the Homotopy Analysis Method*. CRC press, New York (2003)