



## Correction to: Prevalence and determinants of double burden of malnutrition in Bangladesh: evidence from a nationwide cross-sectional survey

Md. Ashfikur Rahman<sup>1</sup> · Henry Ratul Halder<sup>2,4</sup> · Tanjim Siddiquee<sup>3,4</sup> · Syeda Afia Farjana<sup>4</sup> · Harun Or Roshid<sup>4</sup> · Bayezid Khan<sup>1</sup> · Nuzhat Fatema<sup>1</sup> · Md. Hasan Howlader<sup>1</sup> · Shaharior Rahman Razu<sup>5</sup>

Published online: 30 August 2022  
© Springer Nature Switzerland AG 2022

### Correction to: Nutrire (2021) 46:11

<https://doi.org/10.1186/s41110-021-00140-w>

In the original publication of the article, the second sentence of the 4<sup>th</sup> paragraph on page 11 should read:

OWOBM/WSC had a higher likelihood to respondents who were non-exposed to media (AOR: 1.14; 95% CI: 1.11-2.13).

Also, Table 5 was incorrect and should be replaced with:

---

The original article can be found online at <https://doi.org/10.1186/s41110-021-00140-w>.

---

✉ Md. Ashfikur Rahman  
ashfikur@ku.ac.bd

Henry Ratul Halder  
halderhr@myumanitoba.ca

Tanjim Siddiquee  
ts19em@brocku.ca

Syeda Afia Farjana  
afafarjanatisha@gmail.com

Harun Or Roshid  
harunurasid97@gmail.com

Bayezid Khan  
bayezidkhan@ku.ac.bd

Nuzhat Fatema  
fatema.nuzhat@yahoo.com

Md. Hasan Howlader  
hasan\_ds@ku.ac.bd

Shaharior Rahman Razu  
razusocku@gmail.com

- <sup>1</sup> Development Studies Discipline, Social Science School, Khulna University, Khulna, Bangladesh
- <sup>2</sup> Rady Faculty of Health Sciences, Department of Community Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada
- <sup>3</sup> Faculty of Mathematics and Science, Department of Mathematics and Statistics, Brock University, St. Catharines, Ontario, Canada
- <sup>4</sup> Statistics Discipline, Science, Engineering and Technology School, Khulna University, Khulna, Bangladesh
- <sup>5</sup> Sociology Discipline, Social Science School, Khulna University, Khulna, Bangladesh

**Table 5** Estimation of adjusted odds ratio with 95% confidence interval for different forms of double burden of malnutrition using binary logistic regression model

Variables	OWOBM/STC AOR (95% CI)	OWOBM/WSC AOR (95% CI)	OWOBM/UWC AOR (95% CI)	OWOBM/SWUC AOR (95% CI)
<b>Residence</b>				
Rural	Ref.	Ref.	Ref.	Ref.
Urban	1.23 (0.84-1.81)	0.77 (0.43-1.36)	1.13 (0.72-1.77)	1.12 (0.81-1.54)
<b>Wealth index</b>				
Poor	1.44** (1.26-1.75)	1.47** (1.16-3.27)	1.64** (1.35-2.22)	1.59 (1.38-1.92)
Middle	Ref.	Ref.	Ref.	Ref.
Rich	1.27 (0.79-2.05)	1.17 (0.54-2.50)	1.23 (0.57-1.76)	1.13** (1.07-1.70)
<b>Media exposure</b>				
Exposed	Ref.	Ref.	Ref.	Ref.
Non-exposed	1.77*** (1.51-2.16)	1.14* (1.11-2.13)	1.98** (1.60-2.58)	1.86** (1.61-2.22)
<b>Drinking water source</b>				
Protected	Ref.	Ref.	Ref.	Ref.
Unprotected	0.70 (0.36-1.35)	0.48 (0.15-1.50)	0.46 (0.20-1.08)	0.66 (0.38-1.16)
<b>Toilet facility</b>				
Hygienic	Ref.	Ref.	Ref.	Ref.
Unhygienic	1.43* (1.09-1.97)	1.02 (0.52-2.00)	1.40** (1.04-1.92)	1.33*** (1.10-1.95)
<b>Mother's age (years)</b>				
<20	0.75*** (0.42-0.95)	0.43*** (0.14-0.83)	0.32** (0.13-0.79)	0.66*** (0.39-0.91)
20-35	Ref.	Ref.	Ref.	Ref.
>35	2.03* (1.12-3.67)	2.76** (1.29-2.93)	2.40** (1.29-2.86)	2.20** (1.35-2.90)
<b>Mother's education</b>				
No education	Ref.	Ref.	Ref.	Ref.
Primary	0.87 (0.48-1.57)	1.59 (0.60-2.28)	0.88 (0.46-1.69)	0.94 (0.57-1.55)
Secondary	0.90 (0.49-1.66)	1.50 (0.53- 2.25)	0.99 (0.50-1.95)	0.92 (0.55-1.55)
Higher	0.57 (0.25-1.34)	1.86 (0.53-2.50)	0.76 (0.29-1.98)	0.76 (0.38-1.53)
<b>C-section delivery</b>				
No	Ref.	Ref.	Ref.	Ref.
Yes	1.04 (0.66-1.64)	1.25 (0.66-1.41)	0.72 (0.43-1.21)	1.20* (1.08-1.57)
<b>Breastfeeding</b>				
No	1.78** (1.52-2.18)	1.68* (1.33-2.40)	1.53*** (1.06-1.93)	1.89* (1.08-2.56)
Yes	Ref.	Ref.	Ref.	Ref.
<b>Birth order</b>				
First order	Ref.	Ref.	Ref.	Ref.
Other orders	1.68* (1.08-2.60)	2.40* (1.25-2.94)	1.76** (1.05-2.95)	1.92** (1.33-2.79)
<b>Father's education</b>				
No education	Ref.	Ref.	Ref.	Ref.
Primary	1.71* (1.01-2.91)	1.16* (1.04-2.27)	1.53*** (1.08-1.75)	1.63* (1.03-2.56)
Secondary	1.15 (0.63-2.07)	2.26** (1.95-3.37)	1.28*** (1.06-2.04)	1.51* (1.09-2.47)
Higher	1.57 (0.77-3.19)	2.51 (1.88-3.13)	1.65*** (1.13-1.97)	1.75** (1.09-2.19)
<b>Father's occupation</b>				
Service holder	Ref.	Ref.	Ref.	Ref.
Agricultural services	0.95 (0.58-1.55)	1.02 (0.50-2.06)	1.14*** (1.06-1.95)	1.91* (1.61-2.39)
Laborer	0.88 (0.60-1.29)	1.09 (0.62-1.92)	1.87*** (1.55-2.18)	1.98** (1.71-2.35)

OWOBM/STC: Overweight/obese mother and stunted child; OWOBM/WSC: Overweight/obese mother and wasted child; OWOBM/UWC: Overweight/obese mother and underweight child; OWOBM/SWUC: Overweight/obese mother and stunted/wasted/underweight child; C-section: Cesarean delivery; CI: Confidence interval; Ref.: Reference category.

\*\*\*p>0.000; \*\*p>0.001; \*p>0.05.